Bound on nonlocal scale from LEP

N.V.Krasnikov *

Institute for Nuclear Research 60-th October Anniversary Prospect 7a, Moscow 117312, Russia

February 1995

Abstract

Leptons, quarks and gauge bosons are assumed to be described by local field theory. Stringent bound on nonlocal scale can be derived from the high-precision LEP measurements. We find a bound on nonlocal scale $\Lambda_{nl} > 1020 \text{ Gev}$, C.L. 95% from LEP data.

^{*}E-mail address: KRASNIKO@MS2.INR.AC.RU

At present the standard explanation of the gauge hierarchy problem is the supersymmetry with the supersymmetry breaking scale $M_{SUSSY} \leq O(1)Tev$ [1]. However it is possible to explain the gauge hierarchy problem by postulation that there is a nonlocal scale $\Lambda_{nl} \leq O(10)Tev$ [2] in nature. The main idea of the nonlocal field theory [3] consists in the use of nonlocal propagator

$$D_{nl}(p^2) = V_{nl}(p^2)(p^2 - m^2 + i\epsilon)^{-1}$$
(1)

instead of local propagator $D_l(p^2) = (p^2 - m^2 + i\epsilon)^{-1}$. Here $V_{nl}(p^2)$ is an entire function on p^2 decreasing in the euclidean region $p^2 < 0$ that leads to the ultraviolet convergence of the perturbation theory in the case of nonlocal field theory [3]. Typical example of nonlocal formfactor is $V_{nl}(p^2) = \exp(\frac{p^2}{\Lambda_{nl}^2})$. For one-loop correction to the Higgs boson mass we have the following estimate [2]:

$$\delta m_H^2 = O(\frac{g_2^2}{16\pi^2})\Lambda_{nl}^2 \tag{2}$$

Using the well known upper bound [4] $m_H \leq 1 \text{Gev}$ on the Higgs boson mass and requiring that $\delta m_H^2 \leq m_H^2$ one can find an upper bound for nonlocal scale [2]

$$\Lambda_{nl} \le O(10)Tev \tag{3}$$

So if quantum field theory with nonlocal scale is responsible for the solution of the gauge hierarchy problem we should expect the existence of very low nonlocal scale compared to the Planck scale $M_{PL} = 1.2 \cdot 10^{19} \, Gev$.

In this note we find a bound on nonlocal scale $\Lambda_{nl} > 1020 \, Gev$, C.L. 95% from highprecision LEP data. As it has been mentioned before the main idea of nonlocal field theory consists in the use of nonlocal propagator (1) instead of local propagator. At the language of Lagrangians nonlocal propagator (1) corresponds to nonlocal Lagrangian

$$L_{nl} = \frac{1}{2}\phi(x)[V^{-1}(-\Box_x)(\Box_x + m^2)]\phi(x)$$
(4)

for scalar particles. Here $\Box_x = \partial^{\mu}\partial_{\mu}$. For the gauge fields the corresponding nonlocal Lagrangian has the form

$$L_{nl,g} = -\frac{1}{2} Tr[F_{\mu\nu}(x)V^{-1}(-\Box_x)F^{\mu\nu}(x)], \qquad (5)$$

where

$$F_{\mu\nu(x)} = \partial_{\mu}A_{\nu}(x) - \partial_{\nu}A_{\mu}(x) + ig[A_{\mu}(x), A_{\nu}(x)], \qquad (6)$$

$$\Box_x = (\partial_\mu - igA_\mu(x))(\partial^\mu - igA^\mu(x)).$$
(7)

In this paper we shall assume that nonlocal formfactor $V(p^2)$ is the same for SU(3), SU(2) and U(1) gauge groups. It appears that in the first approximation the nonlocale Lagrangian of the type (4) for quarks and leptons does not lead to the observable effects for the LEP measurements so we neglect this type of nonlocality. Typical example for the nonlocal formfactor $V(p^2)$ is [3]

$$V_{nl}(p^2) = \left[\frac{\sin(-p^2l^2)^{1/2}}{(-p^2l^2)^{1/2}}\right]^2.$$
(8)

The formfactor (5) has the meaning of the uniformly distributed charge with the radius 1 [3]. The introduction of the nonlocal formfactor $V_{nl}(p^2)$ leads to the propagator for the Z-boson

$$D_{\mu\nu}(p^2) = \frac{g_{\mu\nu} - \frac{p_{\mu}p_{\nu}}{p^2}}{p^2 V_{nl}^{-1}(p^2) - M_{Z_o}^2}$$
(9)

At present the most accurate and "theoretically clean" LEP observables are the leptonic widths and leptonic forward-backward asymmetries. We have the following formulae for the leptonic widths and asymmetries [5, 6]

$$\Gamma_l = 2^{0.5} G_\mu M_z^3 (g_V^2 + g_A^2) (1 + \frac{3\alpha}{4\pi}) \frac{1}{12\pi} , \qquad (10)$$

$$A_{FB} = \frac{3g_A^2 g_V^2}{(g_V^2 + g_A^2)^2},$$
(11)

where at the tree level

$$g_A = -\frac{1}{2} , \qquad (12)$$

$$g_V = -\frac{1}{2} + 2\sin^2(\theta_w) \,. \tag{13}$$

The corresponding one-loop formulae for vector and axial coupling constants are well known [5, 6]. For instance, for the axial coupling constant in notations of ref.[6] we have

$$g_A = \left(1 - \frac{1}{2}\Pi_w(0) + \frac{1}{2}\Pi_Z(M_Z^2) - \frac{1}{2}D - \frac{1}{2}\Sigma'_Z(M_Z^2)\right)\left(-\frac{1}{2} + 2csF_A\right).$$
(14)

Analogous formula takes place for the vector coupling constant g_V . For the nonlocal formfactor $V_{nl}(p^2)$ we have the asymptotic expansion

$$V_{nl}(p^2) = 1 + \frac{p^2}{\Lambda_{nl}} + O((\frac{p^2}{\Lambda_{nl}^2})^2) .$$
(15)

One can find using the formula (15) that an account of nonlocal propagator (9) for Z-boson leads to an additional factor

$$K = 1 + \frac{M_Z^2}{2\Lambda_{nl}^2} + O((\frac{M_Z^2}{\Lambda_{nl}^2})^2)$$
(16)

for axial and vector lepton coupling constants. We have used the following LEP measurements as input [7]: the mass of the Z, $M_Z = 91.197 \pm 0.007$ Gev, the total width of the Z, $\Gamma_Z = 2.490 \pm 0.007$ Gev, the ratio of the hadronic partial width to the partial width for Z decay into electron, muon and tau pairs $R_e = 20.76 \pm 0.08$, $R_{\mu} = 20.76 \pm 0.07$, $R_{\tau} = 20.80 \pm 0.08$ and the effective leptonic axial and vector couplings with Z-boson $g_A^l = -0.5008 \pm 0.0008$, $g_V^l = 0.0377 \pm 0.0016$. In our analysis we used the formulae of ref.[6] to estimate the standard one-loop corrections. We took the top quark and the Higgs boson masses in the interval $130 Gev \leq m_t \leq 190 Gev$, $60 Gev \leq m_H \leq 500 Gev$. For $\Lambda_{nl}^2 > 0$ we have found that

$$\Lambda_{nl} > 1020(1320) \, Gev, \, C.L.95\%(C.L.68\%) \,. \tag{17}$$

For the less probable from the theoretical point of view case $\Lambda_{nl}^2 < 0$ (typical formfactor for this case is $V_{nl}(p^2) = \exp(-\frac{p^2}{\Lambda^2} - c_1(\frac{p^2}{\Lambda^2})^2)$) we have found that

$$-\Lambda_{nl}^2 > (930Gev)^2((1140Gev)^2), \ C.L.95\%(C.L.68\%)$$
(18)

For the nonlocal formfactor (8) bound (17) corresponds to the bound

$$l < 2.4(1.9) \cdot 10^{-17} cm, \ C.L.95\%(C.L.68\%)$$
 (19)

for nonlocal radius. Note that in ref.[8] bounds on the radii of quarks and leptons and their weak anomalous magnetic moments have been derived from the high-precision measurements at LEP and SLC. The authors of ref.[8] have found bound $R \ge 10^{-17} cm$ for quark and lepton radii that is very closed to our bound (19). To conclude, in this note we have found a bound on nonlocal scale $\Lambda_{nl} > 1020 \, Gev$ (C.L.95%) from LEP measurements. If the nonlocal scale is responsible for the solution of the gauge hierarchy problem then it could be discovered at LHC provided that $\Lambda_{nl} \leq$ $10 \, Tev$ (the nonlocal glueon propagator will change the prediction for the two jet cross sections compared to standard QCD case).

I am indebted to the collaborators of the INR theoretical department for discussions and critical comments. I thank prof. P.M.Zervas for sending me preprint [8]. The research described in this publication was made possible in part by Grant N6G000 from the International Science Foundation and by Grant 94-02-04474-a of the Russian Scientific Foundation.

References

- For reviews, see: H.P.Nilles, Phys.Rep.110(1984)1; G.G.Ross, Grand Unified Theories (Benjamin, New York 1984); R.N.Mohapatra, Unification and Supersymmetry (Springer, New York 1992).
- N.V.Krasnikov, Soviet Journal Theoretical and Mathematical Physics, 73(1987)325;
 N.V.Krasnikov, Phys.Lett.B214(1988)363.
- G.V.Efimov, Non-local Interactions of Quantum Fields, Nauka, Moscow (1977) (in russian); V.A.Alebastrov and G.V.Efimov, Commun. Math. Phys.38(1974)11;
 G.V.Efimov, Annals of Phys.7191972)466.
- [4] B.W.Lee, C.Quigg and H.B.Thacker, Phys.Rev.Lett.38(1977) 1330.
- [5] For reviews, see: G.Altarelli, R.Kleiss and C.Verzegnassi,eds., Physics at LEP 1, vol.1, CERN report CERN 89-08(CERN, Geneva, 1989); R.Barbieri, Lectures given at the Cargese Summer School, july 1992, CERN Preprint CERRN-TH.6659/92.
- [6] V.A.Novikov, L.B.Okun and M.I.Vysotsky, Nucl. Phys. B397 (1993)35.
- [7] Particle Data Group, Phys.Rev.**D50**(1994)1173.
- [8] G.Kopp, D.Schaile, M.Spira and P.M.Zervas, Bounds on Radii and Magnetic Dipole Moments of Quarks and Leptons from LEP, SLC and HERA, DESY Preprint, DESY 94-148.