

Calculation of Power Corrections to Hadronic Event Shapes*

Yu.L. Dokshitzer[†]

Department of Theoretical Physics, University of Lund,
Sölvegatan 14 A, S-223 62 Lund, Sweden

and

B.R. Webber

Cavendish Laboratory, University of Cambridge,
Madingley Road, Cambridge CB3 0HE, U.K.

Abstract

We compute power corrections to hadronic event shapes in e^+e^- annihilation, assuming an infrared regular behaviour of the effective coupling α_s . With the integral of α_s over the infrared region as the only non-perturbative parameter, also measured in heavy quark physics, we can account for the empirical features of $1/Q$ corrections to the mean values of various event shapes.

Cavendish-HEP-95/2
LU TP 95-8
March 1995

*Research supported in part by the U.K. Particle Physics and Astronomy Research Council and by the EC Programme “Human Capital and Mobility”, Network “Physics at High Energy Colliders”, contract CHRX-CT93-0357 (DG 12 COMA).

[†]Permanent address: St Petersburg Nuclear Physics Institute, Gatchina, St Petersburg 188350, Russian Federation

1. Introduction

Infrared-safe shape measures for hadronic final states in e^+e^- annihilation would appear in principle to be an ideal testing ground for perturbative QCD. Such quantities are asymptotically insensitive to long-distance non-perturbative physics and can thus be computed order-by-order in perturbation theory. The large momentum scale $Q \sim M_Z$ available at existing e^+e^- machines implies a small value of the running coupling $\alpha_s(Q)$ and so the perturbation series should be relatively well-behaved. Non-perturbative effects should be suppressed by inverse powers of Q . Thus hadronic event shapes should be capable of providing a high-precision measurement of α_s .

Unfortunately the hoped-for precision has not yet been achieved, partly because $\mathcal{O}(\alpha_s^3)$ calculations of event shapes are still lacking, but also because non-perturbative effects turn out to be significant even at $Q \sim M_Z$. This is because they are in fact suppressed by only a single inverse power of Q in many cases. Bearing in mind that $\alpha_s(M_Z) \sim 0.12$ and the non-perturbative scale is $\mathcal{O}(1 \text{ GeV})$, we see that the power correction may easily be comparable with the $\mathcal{O}(\alpha_s^2)$ next-to-leading term at present energies. Consequently it becomes essential to achieve some understanding of power corrections before embarking on any $\mathcal{O}(\alpha_s^3)$ calculations of event shapes.

In the present paper we adopt the approach, advocated in Refs. [1,2], of trying to deduce as much as possible about power corrections from perturbation theory. In particular we explore the consequences of assuming that α_s , defined in some appropriate way, does not grow indefinitely at low scales but instead has an infrared-regular effective form. Then various moments of α_s , integrated over the infrared region, play the rôle of non-perturbative parameters which determine the form and magnitude of power corrections. Since α_s is supposed to be universal, we obtain relations between the power corrections to various observables.

Our approach is related to that of Korchemsky and Sterman [3], and also to several other recent papers that discuss power corrections and the ambiguities of perturbation theory in terms of infrared renormalons [4], in the context of the Drell-Yan process [5], event shapes [6], deep inelastic scattering [7], heavy quark effective theory [8] or quark confinement [9]. From our viewpoint, infrared renormalons arise from the divergence of the perturbative expression for α_s at low scales, and the ambiguities associated with different ways of avoiding the renormalon poles in the Borel transform plane are resolved by specifying the infrared behaviour of α_s . This approach implies relationships between the contributions of a given renormalon to different processes.

The quantitative results we obtain look very good in the case of the mean value of the thrust parameter [10]. The required value of the relevant moment of α_s is consistent with that obtained from a similar approach to heavy quark fragmentation [1]. For the other shape variables discussed here, the mean value of the C -parameter [11] and the longitudinal cross section [12], a comparison with LEP data is encouraging, but detailed tests must await the re-analysis of lower-energy data to establish the energy dependence of the leading power correction.

2. Calculations

Consider a quantity of the form

$$F = \int_0^Q dk f(k) \quad (1)$$

where $f(k)$ behaves like $\alpha_s(k) k^p$ at $k \ll Q$, say

$$f(k) \sim a_F \alpha_s(k) k^p / Q^{p+1} \quad (k \ll Q) \quad (2)$$

where we have included the appropriate Q dependence assuming F is dimensionless. Suppose that F has the perturbative expansion

$$F^{\text{pert}} = F_1 \alpha_s + F_2 \alpha_s^2 + \dots \quad (3)$$

More precisely, if the coefficients F_n are computed in the $\overline{\text{MS}}$ renormalization scheme at scale Q , then in terms of the $\overline{\text{MS}}$ coupling at scale μ_R we have

$$F^{\text{pert}} = F_1 \alpha_s(\mu_R) + \left(F_2 + \frac{\beta_0}{2\pi} \ln \frac{\mu_R}{Q} F_1 \right) \alpha_s^2(\mu_R) + \dots \quad (4)$$

where $\beta_0 = (11C_A - 2N_f)/3$, with $C_A = 3$, for N_f active flavours.

In part of the integration region of Eq. (1) the perturbative expression for $\alpha_s(k)$ is not appropriate. We may however choose an infrared matching scale μ_I such that $\Lambda \ll \mu_I \ll Q$ and assume that perturbation theory is valid above that scale. We can then introduce a non-perturbative parameter $\bar{\alpha}_p(\mu_I)$ to represent the portion of the integral below μ_I :

$$\int_0^{\mu_I} dk \alpha_s(k) k^p \equiv \frac{\mu_I^{p+1}}{p+1} \bar{\alpha}_p(\mu_I) \quad (5)$$

Before adding this contribution to F^{pert} , we have to subtract the perturbative value of this integral, which is, to second order,

$$\frac{\mu_I^{p+1}}{p+1} \left[\alpha_s(\mu_R) + \frac{\beta_0}{2\pi} \left(\ln \frac{\mu_R}{\mu_I} + \frac{1}{p+1} \right) \alpha_s^2(\mu_R) \right] \quad (6)$$

As a refinement, and for consistency with Ref. [1], we shall assume that the parameter $\bar{\alpha}_p(\mu_I)$ refers not to the coupling in the $\overline{\text{MS}}$ scheme but rather to the scheme proposed in Ref. [13], which is expected to be more physical in the region under consideration. Thus α_s in Eq. (5) (only) is to be interpreted as α_s^{eff} where in terms of the $\overline{\text{MS}}$ coupling

$$\alpha_s^{\text{eff}} = \alpha_s + \frac{K}{2\pi} \alpha_s^2 \quad (7)$$

with

$$K = \left(\frac{67}{18} - \frac{\pi^2}{6} \right) C_A - \frac{5}{9} N_f \quad (8)$$

The only effect on Eq. (6) is that the term $\ln(\mu_R/\mu_I)$ becomes $\ln(\mu_R/\mu_I) + K/\beta_0$. The full expression for F is thus

$$F = F^{\text{pert}} + F^{\text{pow}} \quad (9)$$

where F^{pert} is as given in Eq. (4) while

$$F^{\text{pow}} = \frac{a_F}{p+1} \left(\frac{\mu_1}{Q} \right)^{p+1} \left[\bar{\alpha}_p(\mu_1) - \alpha_s(\mu_R) - \frac{\beta_0}{2\pi} \left(\ln \frac{\mu_R}{\mu_1} + \frac{K}{\beta_0} + \frac{1}{p+1} \right) \alpha_s^2(\mu_R) \right]. \quad (10)$$

The dependence of F^{pert} on the renormalization scale μ_R is one order higher in α_s than that calculated, i.e. third-order in this case. Similarly, the dependence of the power correction F^{pow} on both μ_R and the infrared matching scale μ_1 is third-order, provided μ_1 is sufficiently large for $\alpha_s(\mu_1)$ to have reached its perturbative form. Of course, the value obtained for $\bar{\alpha}_p(\mu_1)$ depends on μ_1 , but this is mostly compensated by the other μ_1 -dependent term.

The value of the power p and the coefficient a_F may be found from the infrared cutoff dependence of the lowest-order perturbative result. In this connection, it is crucial that the appropriate argument of α_s for soft and/or collinear gluon emission is the gluon transverse momentum k_\perp [14]. Thus the cutoff should be a k_\perp -cutoff.

Consider for example the mean value of the thrust T . The contribution to this quantity from the region $k_\perp < \mu_1$ is

$$\delta \langle T \rangle = -\frac{C_F}{2\pi} \int_{k_\perp < \mu_1} dx_1 dx_2 \alpha_s(k_\perp) \frac{x_1^2 + x_2^2}{(1-x_1)(1-x_2)} \min\{(1-x_1), (1-x_2)\} \quad (11)$$

where $C_F = 4/3$. Setting $1-x_{1,2} = y_{1,2}$ and considering the region $y_1 < y_2 \ll 1$, we have $k_\perp = \sqrt{y_1 y_2} Q$ and hence

$$\begin{aligned} \delta \langle T \rangle &= -4 \frac{C_F}{\pi} \int_0^{\mu_1/Q} dy_1 \int_{y_1 Q}^{\mu_1} \frac{dk_\perp}{k_\perp} \alpha_s(k_\perp) \\ &= -\frac{4C_F}{\pi Q} \int_0^{\mu_1} dk_\perp \alpha_s(k_\perp) \equiv -4 \frac{C_F}{\pi} \frac{\mu_1}{Q} \bar{\alpha}_0(\mu_1). \end{aligned} \quad (12)$$

Thus in this case $p = 0$ and we obtain a $1/Q$ correction, with a coefficient in Eq. (10) of $a_F = a_T$ where

$$a_T = -4 \frac{C_F}{\pi} = -1.70. \quad (13)$$

As shown by the solid curve in Fig. 1, an excellent fit to the data on $\langle T \rangle$ over the range $14 < Q < 92$ GeV can be obtained using the perturbative prediction [15]

$$\langle T \rangle^{\text{pert}} = 1 - 0.335 \alpha_s - 1.02 \alpha_s^2 \quad (14)$$

with $\mu_R = Q$ and $\alpha_s(M_Z) = 0.117 \pm 0.005$ [16], plus a power correction of the form (10). For $\mu_1 = 2$ GeV, the fitted value of the non-perturbative parameter $\bar{\alpha}_0$ is

$$\bar{\alpha}_0(2 \text{ GeV}) \equiv (2 \text{ GeV})^{-1} \int_0^{2 \text{ GeV}} dk \alpha_s^{\text{eff}}(k) = 0.53 \pm 0.04, \quad (15)$$

with $\chi^2 = 4.5$ for 8 degrees of freedom. Allowing both $\alpha_s(M_Z)$ and $\bar{\alpha}_0(2 \text{ GeV})$ to be free parameters gives

$$\alpha_s(M_Z) = 0.120 \pm 0.004, \quad \bar{\alpha}_0(2 \text{ GeV}) = 0.52 \pm 0.03, \quad (16)$$

with $\chi^2/\text{d.o.f.} = 3.7/7$.




Figure 1: Mean value of $1 - T$, where T is the thrust.

We also obtain good fits for other values of the arbitrary infrared matching parameter μ_t . At $\mu_t = 3$ GeV, for example, we find

$$\alpha_s(M_Z) = 0.118 \pm 0.004, \quad \bar{\alpha}_0(3 \text{ GeV}) = 0.42 \pm 0.03, \quad (17)$$

with $\chi^2/\text{d.o.f.} = 4.0/7$. The change in $\bar{\alpha}_0$ implies that $\alpha_s^{\text{eff}}(2.5 \text{ GeV}) \simeq 0.2 \pm 0.1$, which is reasonable, the perturbative value being around 0.3. The change in the overall power correction is small (about 5%), since, as explained above, the μ_t -dependence mostly cancels in Eq. (10).

Using the value (15) of $\bar{\alpha}_0$, obtained by fitting the thrust data, one can now predict the power corrections to other event shapes. For the mean value of the C -parameter, for example, we find that the coefficient a_C in Eq. (10) is

$$a_C = 6 C_F = 8. \quad (18)$$

At present there are only data on $\langle C \rangle$ at $Q = M_Z$, where Eqs. (15) and (18) imply

$$\langle C \rangle^{\text{pow}} = 0.057 \pm 0.008. \quad (19)$$

The second-order perturbative prediction is [15]

$$\langle C \rangle^{\text{pert}} = 1.375 \alpha_s + 3.88 \alpha_s^2 = 0.214 \pm 0.011 \quad (20)$$

for $\alpha_s = 0.117 \pm 0.005$. Thus the full theoretical prediction is

$$\langle C \rangle^{\text{th}} = \langle C \rangle^{\text{pert}} + \langle C \rangle^{\text{pow}} = 0.271 \pm 0.014, \quad (21)$$

which is consistent with the experimental result [17]

$$\langle C \rangle^{\text{exp}} = 0.2587 \pm 0.0013 \pm 0.0018 . \quad (22)$$

Note that the power correction represents over 20% of the value of this quantity.

Finally, for the longitudinal cross section fraction $\sigma_L/\sigma_{\text{tot}}$ we predict a coefficient

$$a_L = C_F = 1.33 , \quad (23)$$

leading to the power correction at $Q = M_Z$

$$(\sigma_L/\sigma_{\text{tot}})^{\text{pow}} = 0.010 \pm 0.001 . \quad (24)$$

The first-order perturbative prediction is $\alpha_s/\pi = 0.037$. However, the second-order correction is not yet known. The preliminary OPAL measurement [18] is

$$(\sigma_L/\sigma_{\text{tot}})^{\text{exp}} = 0.067 \pm 0.008 . \quad (25)$$

Clearly the second-order perturbative correction should be significant if there is to be satisfactory agreement between theory and experiment.

The values for $\bar{\alpha}_0$ obtained above from event shapes are in reasonable agreement with those deduced from heavy quark fragmentation spectra. In Ref. [1] the value $\bar{\alpha}_0(2 \text{ GeV}) \simeq 0.6$ was obtained from fits to heavy quark energy losses in e^+e^- annihilation. The same conclusion follows from an analysis of the quantity $\xi_H = -\ln \langle x_H \rangle$, where x_H is the energy fraction carried by the heavy quark H , using the approach of the present paper. We find a quark mass ($1/M$) correction of the form (10), with Q replaced by M , $\mu_R \sim M$, and coefficient $a_H = C_F/2$. The perturbative prediction deduced from Ref. [1] is

$$\begin{aligned} \xi_H^{\text{pert}} = & \frac{4C_F}{3\pi} \left\{ \int_M^Q \frac{dk}{k} \alpha_s(k) - \frac{35}{24} \alpha_s(Q) + \frac{13}{24} \alpha_s(M) \right. \\ & \left. + \frac{1}{\beta_0} (K + \delta_2) [\alpha_s(M) - \alpha_s(Q)] \right\} , \end{aligned} \quad (26)$$

where δ_2 is the (numerically negligible) 2-loop anomalous dimension correction

$$\delta_2 = \left(\frac{53}{18} - \frac{\pi^2}{3} \right) C_F + \frac{31}{36} (C_A - 2C_F) = -0.173 . \quad (27)$$

The expression (26), which accounts for the $\alpha_s \ln(Q/M)$, α_s and $\alpha_s^2 \ln(Q/M)$ terms in ξ_H , but neglects α_s^2 terms with no large logarithm, gives $\xi_b^{\text{pert}} = 0.26 \pm 0.02$ for b -quarks at $Q = M_Z$. Comparing with the experimental value of 0.36 ± 0.02 deduced from lepton spectra [19], this implies that $\xi_b^{\text{pow}} = 0.10 \pm 0.03$ and hence that $\bar{\alpha}_0(2 \text{ GeV}) \simeq 0.6 \pm 0.1$. The errors are estimated conservatively, taking into account the small scale $M_b \sim 5 \text{ GeV}$ and the lack of a complete $\mathcal{O}(\alpha_s^2)$ calculation of ξ_b^{pert} .

3. Conclusions

Note that the power correction coefficients a_T , a_C and a_L deduced above using a k_\perp cutoff are identical to those obtained in Ref. [2] with a gluon mass cutoff. With a k_\perp cutoff, however, these coefficients have a physical interpretation: they measure the contribution of the low-scale region in which α_s departs significantly from its perturbative form. After being used to calculate the coefficients, the cutoff is replaced by an infrared matching parameter μ_\perp , which represents the scale below which we switch from the perturbative to the non-perturbative description of α_s . As long as μ_\perp is not too small (larger than about 1 GeV) the predictions are quite insensitive to its value, indicating that the perturbative behaviour has set in at that scale.

The divergence in the perturbative expression for α_s at low scales is responsible for the divergence of the perturbation series for quantities like those considered here, giving rise to the so-called “renormalon ambiguity”. By assuming an infrared regular form for the effective coupling, we resolve this ambiguity, at the price of introducing the non-perturbative parameters $\bar{\alpha}_p$. These parameters are, however, universal, and can be measured experimentally, like $\bar{\alpha}_0$ in Eq. (15).

Combined fits to the non-perturbative parameters $\bar{\alpha}_p$ and the perturbative parameter α_s , using data on several different event shapes, provide the possibility of understanding something new about QCD at low scales and at the same time measuring α_s with improved precision. This would be useful not only for QCD but also in constraining physics beyond the Standard Model.

References

1. Yu.L. Dokshitzer, in *Proc. XXVII Recontre de Moriond: Perturbative QCD and Hadronic Interactions, March, 1992*, ed. J. Tran Than Van (Editions Frontieres, 1992); Yu.L. Dokshitzer, V.A. Khoze and S.I. Troyan, Lund preprints LU TP 92–10, 94–23; Yu.L. Dokshitzer, in *Proc. Int. School of Subnuclear Physics, Erice, 1993*.
2. B.R. Webber, *Phys. Lett.* B339 (1994) 148; see also *Proc. Summer School on Hadronic Aspects of Collider Physics, Zuoz, Switzerland, August 1994*, ed. M.P. Locher (PSI, Villigen, 1994).
3. G.P. Korchemsky and G. Sterman, *Nucl. Phys.* B437 (1995) 415.
4. A.H. Mueller, in *QCD 20 Years Later*, vol. 1 (World Scientific, Singapore, 1993).
5. H. Contopanagos and G. Sterman, *Nucl. Phys.* B419 (1994) 77.
6. A.V. Manohar and M.B. Wise, *Phys. Lett.* 344B (1995) 407.
7. X. Ji, preprint MIT-CTP-2381 (hep-ph/9411312).
8. I.I. Bigi, M.A. Shifman, N.G. Uraltsev and A.I. Vainshtein, *Phys. Rev.* D50 (1994) 2234; M. Beneke and V.M. Braun, *Nucl. Phys.* B426 (1994) 301; M. Neubert and C.T. Sachrajda, preprint CERN-TH.7312/94 (hep-ph/9407394).
9. U. Aglietti and Z. Ligeti, preprint CALT-68-1982 (hep-ph/9503209).
10. E. Farhi, *Phys. Rev. Lett.* 39 (1977) 1587.
11. R.K. Ellis, D.A. Ross and A.E. Terrano, *Nucl. Phys.* B178 (1981) 421.
12. P. Nason and B.R. Webber, *Nucl. Phys.* B421 (1994) 473.
13. S. Catani, G. Marchesini and B.R. Webber, *Nucl. Phys.* B349 (1991) 635.
14. Yu.L. Dokshitzer, D.I. Dyakonov and S.I. Troyan, *Phys. Reports* 58 (1980) 270; D. Amati, A. Bassetto, M. Ciafaloni, G. Marchesini and G. Veneziano, *Nucl. Phys.* B173 (1980) 429.
15. Z. Kunszt, P. Nason, G. Marchesini and B.R. Webber, in *Z Physics at LEP 1*, CERN Yellow Book 89-08.
16. B.R. Webber, in *Proc. XXVII Int. Conf. on High Energy Physics, Glasgow, 1994*, ed. P.J. Bussey and I.G. Knowles (Institute of Physics, 1995).
17. ALEPH Collaboration, D. Buskulic et al., *Z. Phys.* C55 (1992) 209.
18. OPAL Collaboration, D.R. Ward, *Nucl. Phys. B (Proc. Suppl.)* 39B, C (1995) 134.
19. T. Behnke, in *Proc. XXVI Int. Conf. on High Energy Physics, Dallas, 1992*, ed. J.R. Sanford (AIP, 1993).