

A Self-consistent Inclusion of Magnetic Screening
for the Quark-Gluon Plasma

G. ALEXANIAN and V. P. NAIK *

Physics Department

City College of the City University of New York

New York, New York 10031.

Abstract

We discuss how magnetic screening can be systematically included in a self-consistent way for Chromodynamics at high temperatures. The resulting gap equation, which gives the summation of selfenergy insertions, is calculated to one-loop order and leads to a nonzero value for the magnetic mass.

* E-mail: gamik@scisun.sciccnycuny.edu, vpn@ajanta.sciccnycuny.edu

The generating functional for hard thermal loops in Quantum Chromodynamics (QCD) is closely related to the eikonal for a Chern-Simons theory [1]. The importance and various properties of hard thermal loops have been the subject of many recent investigations [2-5]. The generalization of the Chern-Simons eikonal to a moving plasma suggests that there is another closely related gauge-invariant mass term which gives screening of static magnetic interactions, in other words, a magnetic mass term [6]. Although nonlocal, the equations of motion for this term can be written in a local way by using auxiliary fields and without introducing additional degrees of freedom. Because of this last property, the ultraviolet behaviour of the theory is unchanged. On this basis, it was suggested that this mass term could be used as a gauge-invariant infrared cutoff for loop calculations in QCD at high temperatures. In this paper we discuss how this can be carried out systematically. Some recent related papers are listed as ref. [7].

A consistent implementation of a gauge-invariant infrared cutoff will lead to a gap equation for the value of the magnetic mass. We obtain this equation to one-loop order; as usual, this means a self-consistent summation of one loop self-energy insertions. The magnetic mass to this order is obtained as $m_m^2 = \frac{3}{8} C g^2 T$ where g is coupling constant, T is the temperature and C is defined by $C^{ab} = f^{am n} f^{bn}$, $f^{am n}$ being the structure constants of the gauge group. ($C = N$ for an $SU(N)$ -gauge theory.) An immediate question is whether the two- and higher loop contributions are smaller than the one-loop terms. Purely based on counting dimensions of momentum integrals and powers of g , we cannot conclude whether higher loop effects are smaller or not. It is really an issue of numerical factors and possible logarithms of g . Now, the magnetic mass, by the standard arguments of dimensional reduction at high temperatures is related to the mass gap of the three-dimensional QCD. For the mass gap of three-dimensional gauge theories, the one-loop calculations can hardly be adequate. However, for the quark-gluon plasma, perturbative loop expansion, perhaps with resummations, is expected to be reasonable at high temperatures, at least for a number of processes of interest and our approach is applicable. In any case, qualitatively, it is interesting that a nonzero value can be obtained to this order; computationally, it is useful to have a loopwise implementation of gauge-invariant magnetic screening, irrespective of the specific numerical value.

In the rest frame of the plasma, the magnetic mass term considered in [6] has the following form

$$\tilde{S}_m = M^2 S_m \quad (1a)$$

$$S_m = \int d^3x \int d^3n K(A_n; A_n) \quad (1b)$$

where $A_n = \frac{1}{2}A_i n_i; A_n = \frac{1}{2}A_i n_i$. n_i is a (complex) three-dimensional null vector of the form

$$n_i = (\cos \alpha \cos \alpha' \quad i \sin \alpha'; \cos \alpha \sin \alpha' + i \cos \alpha'; \sin \alpha) \quad (2)$$

In Eq.(1), $d^3n = \sin \alpha \, d\alpha \, d\alpha'$ and denotes integration over the angles of n_i . $K(A_n; A_n)$ is given by

$$K(A_n; A_n) = \frac{1}{2} \int d^2x^T \int d^2z \text{Tr}(A_n; A_n) + i \int I(A_n) + i \int I(A_n) \quad (3)$$

$z = n \times x; z = n \times x^T$ denotes coordinates transverse to n , i.e., $x^T n = 0$: Also

$$I(A_n) = i \int \frac{d^2x^T}{2} \frac{(1)^m}{m} \int \frac{d^2z_1}{2} \dots \frac{d^2z_m}{2} \frac{\text{Tr}(A_n(x_1) \dots A_n(x_m))}{z_{12} z_{23} \dots z_{m-1m} z_{m1}} \quad (4)$$

$z_{ij} = z_i - z_j$. The argument of all A 's in Eq.(4) is the same for the transverse coordinates x^T . The lowest order term in S_m was shown to be

$$S_m = \int \frac{d^4k}{(2\pi)^4} \frac{1}{2} A_i^a(k) A_j^a(k) \delta_{ij} \frac{k_i k_j}{k^2} + O(A^3) \quad (5)$$

This term involves only the transverse potentials, as expected for magnetic screening and on account of gauge invariance. The terms with higher powers of A make S_m invariant under the full non-Abelian gauge transformations.

The general strategy for the inclusion of this term is as follows. We write the action as

$$S = S_0 + M^{-2} S_m \quad (6)$$

S_0 is the standard quark and gluon part of the action. Below we shall not consider the quark terms since their effects are small and have the same general qualitative features. S_0 will be just the Yang-Mills action. S is taken to have a loop expansion, $S = S^{(0)} + S^{(1)} + S^{(2)} + \dots$. Calculations can be done in a loop expansion. We require the pole of the propagator to remain at $k_0^2 = k^2 = M^2$ (for the transverse potentials) as loop corrections are added. This requires choosing $S^{(1)}$ to cancel the one-loop shift of the pole, $S^{(2)}$ to cancel the two-loop shift of the pole, etc., as is usually done for mass renormalization. Of course, we do not want to change the theory, only rearrange and resum various terms. Thus we should impose the condition

$$S = S^{(0)} + S^{(1)} + S^{(2)} + \dots = M^2 \quad (7)$$

This condition is the gap equation determining M in terms of g^2 and T . This procedure of adding and subtracting a mass term, with a gap equation required for consistency, is very standard, for example, in the Nambu-Jona-Lasinio model. It amounts to the self-consistent summation of self-energy corrections. The difference in the present case is that, for reasons of gauge invariance, the mass term involves an infinite number of interaction vertices as well.

We shall now turn to the explicit one-loop calculations. For the plasma at high temperatures, the effects of magnetic screening will be insignificant for high momentum processes. The regime of interest involves momenta small compared to T . The thermal part of the gluon propagator simplifies as

$$\frac{(k^2 - M^2)}{e^{ik_0 T}} \frac{1}{k^2 + M^2} \frac{T}{2k^2} ((k_0 - k) + (k_0 + k)) \quad (8)$$

with a coupling constant $e = \sqrt{g^2 T}$; the thermal part of the loop contributions can be done in a three-dimensional theory. This is, of course, the standard dimensional reduction argument. The electrostatic field, with a Debye mass of order gT , will also be neglected for low momentum calculations. The relevant momenta for which this approximation is valid will be of order $g^2 T$; the coupling constant is also self-consistently evaluated at a scale of order $g^2 T$. This is all in keeping with the assumed hierarchy of $Q_{CD} \ll g^2 T \ll gT \ll T$ for the hot quark-gluon plasma.

The action for momenta small compared to T and gT can be written as

$$S = \int d^3x \left[\frac{1}{4e^2} F^a F^a + \frac{M^2}{e^2} L_m \right] \quad (9)$$

$F^a = \partial A^a - \partial A^a + f^{abc} A^b A^c$. The integral of L_m is the three-dimensional Euclidean version of Eqs.(1b,3,4). There is now only one transverse coordinate x^T . A convenient gauge-fixing term is $\frac{1}{2} \partial A (1 - M^2 \frac{1}{\partial^2}) \partial A$. This gives the gluon propagator as $\delta^{ab} \delta_{ij} (k^2 + M^2)^{-1}$. The sum of all one-loop contributions to the gluon polarization is finite. The functional integral is thus

$$Z = \int [dA] \det(\partial D) e^S \quad (10)$$

$D^{ab} = \partial^{ab} + f^{acb} A^c$. The action simplifies as

$$S = \int \frac{1}{2} A_i^a (\partial^2 + M^2) A_i^a + \int A_i^a A_j^b A_k^c f^{abc} (v_{ijk} + V_{ijk}) + \frac{1}{4} f^{am n} f^{abc} A_i^m A_j^n A_i^b A_j^c + \dots \quad (11)$$

$$v_{ijk}(k; q; (k+q)) = \frac{i}{6} [(2k+q)_j{}_{ik} (2q+k)_i{}_{jk} + (q-k)_k{}_{ij}] \quad (12a)$$

$$V_{ijk}(k; q; (k+q)) = i \frac{M^2}{24} \int d^4n \frac{n_i n_j n_k}{k \cdot n} \frac{q \cdot n}{q \cdot n} \frac{(q+k) \cdot n}{(q+k) \cdot n} \quad (12b)$$

This, after angular integration, becomes

$$V_{ijk}(k; q; (k+q)) = i \frac{M^2}{6} \frac{1}{k^2 q^2} \frac{1}{(q-k)^2} \frac{q \cdot k}{k^2} \frac{q \cdot (q+k)}{(q+k)^2} k_i k_j k_k + \frac{k \cdot (q+k)}{(q+k)^2} (q_i q_j k_k + q_k q_i k_j + q_j q_k k_i) \quad (q \cdot k) \quad (12c)$$

Expression (12c) is the contribution from S_m . The four-point vertex involves, in addition to the standard Yang-Mills vertex of Eq.(11), a term from S_m of the form $\int d^4n (n_i n_j n_k n_l)$. Since n_i is a null vector we get zero from this term to one loop order. This is one of the advantages of our form of S_m ; because n_i is null, Wick contractions at the same point, with a propagator of the form $\delta_{ij} (k^2 + M^2)^{-1}$, give zero. Vertices of higher than quartic order do not contribute at one-loop level. The relevant Feynman diagrams are shown in Fig. 1. The calculations are straightforward although somewhat tedious. Despite the algebraic complexity of V_{ijk} , there are many cancellations and for the one loop part of the effective action we find

$$(\text{1 loop}) = \int \frac{d^3k}{(2\pi)^3} \frac{1}{2} A_i^a(k) \Gamma_{ij}(k) A_j^a(k) + \dots \quad (13)$$

$$\Gamma_{ij} = \Gamma_{ij}^{(1)} + \Gamma_{ij}^{(2)} = \frac{\Gamma_{ij}^{(1)}}{e^2} = \frac{k_i k_j}{k^2} \quad (14a)$$

$$\Gamma_{ij}^{(1)} = C \Gamma_{ij} = \frac{M}{2} \int_0^1 d^4p \frac{3}{4} \frac{p \cdot (k^2 - p^2) + M^2}{k^2 (k^2 - p^2) + M^2} \frac{k \cdot p}{8} \frac{p \cdot (k^2 - p^2)}{k^2 (k^2 - p^2) + M^2} + \frac{k^2}{16} \frac{5}{k^2 (k^2 - p^2) + M^2} + \frac{2}{16} \frac{p \cdot (k^2 - p^2)}{k^2 (k^2 - p^2) + M^2} + C \frac{k_i k_j}{16} \int_0^1 d^4p \frac{3 + 6(k^2 - p^2)}{k^2 (k^2 - p^2) + M^2} \frac{2}{k} \frac{p \cdot (k^2 - p^2)}{k^2 (k^2 - p^2) + M^2} \quad (14b)$$

$$\Gamma_{ij}^{(2)} = \frac{C}{4} \Gamma_{ij} = 3 \frac{k_i k_j}{k^2} \int_0^1 d^4p \frac{p \cdot (k^2 - p^2)}{k^2 (k^2 - p^2) + M^2} \frac{p \cdot (k^2 - p^2)}{k^2 (k^2 - p^2) + M^2} + \frac{C}{8} \Gamma_{ij} + \frac{k_i k_j}{k^2} \int_0^1 d^4p \frac{p \cdot (k^2 - p^2)}{k^2 (k^2 - p^2) + M^2} \frac{p \cdot (k^2 - p^2)}{k^2 (k^2 - p^2) + M^2} + \frac{p \cdot (k^2 - p^2)}{k^2} \quad (14c)$$

$\Gamma_{ij}^{(1)}$ is the contribution of the standard Yang-Mills diagrams. $\Gamma_{ij}^{(2)}$ involves the new vertex (12c), by itself and mixed with (12a). The integrals can actually be evaluated in terms of elementary functions but we do not need the explicit form in what follows. The total contribution to Γ_{ij} can be written as

$$\Gamma_{ij} = \Gamma_{ij}^{(1)} \frac{k_i k_j}{k^2} \frac{M^2}{e^2} B + (K) \frac{(1)}{M^2} \quad (15a)$$

$$(K) = \frac{C e^2}{4 M} \int_0^1 d \frac{9p}{2} \frac{1}{K \left(\frac{p^2}{M^2} + 1 \right) + 1} + \frac{p}{2} \frac{1}{K \left(\frac{p^2}{M^2} + 2 \right)} + \frac{(K-1)}{4} \frac{p}{K \left(\frac{p^2}{M^2} + 1 \right) + 1} \frac{(5 \frac{p^2}{M^2} + 2)}{2} + \frac{p}{5} \frac{1}{1 + \frac{p^2}{M^2}} \quad (15a)$$

$$+ \frac{3}{4} \frac{p}{1 + \frac{p^2}{M^2}}$$

$$B = \frac{C e^2}{4 M} \int_0^1 d \frac{p}{5} \frac{1}{1 + \frac{p^2}{M^2}} + \frac{3}{4} \frac{p}{1 + \frac{p^2}{M^2}} \quad (15b)$$

$$\frac{C e^2}{4 M} \quad (2:384)$$

$K = (k^2 + M^2)/M^2$. Notice that Γ_{ij} is transverse as expected on grounds of gauge invariance. (K) vanishes at $K = 0$ and has no other zeros for positive K as can be checked graphically. (For negative K , (K) is complex and our calculation which uses time-ordered products is not applicable; see ref. [4].) The one-loop corrected inverse propagator has the form $\frac{M^2}{e^2} K + (K) + B \frac{(1)}{M^2}$: Since $(K) = 0$ at $K = 0$, we see that the pole of the propagator will not be shifted if we choose $B = \frac{(1)}{M^2}$ or $(1) \frac{C e^2 M}{4} \quad (2:384)$. The gap equation (7) then gives

$$M = (2:384) \frac{C e^2}{4} \quad (2:384) \frac{C g^2 T}{4} \quad (16)$$

With this choice of (1) , the inverse propagator has the form $\frac{M^2}{e^2} K + (K)$. The correction (K) is significant compared to K , for small K ; for large K , it approaches the result for massless Yang-Mills theory; this is a necessary check for any infrared cutoff.

Even though Γ_{ij} is transverse in agreement with gauge invariance, it depends on the gauge fixing used for the gluon propagator in the loop. The position of the pole and hence the gap equation do not depend on this. One can explicitly check this. The simplest way is as follows. In the effective action, one can have a gauge-dependent Γ_{ij} ; Generally, the higher point functions are gauge dependent as well. Physical results, such as scattering amplitudes, are independent of the gauge fixing used. Alternatively, one can define a new

two-point function, a new three-point function, etc., which are independent of the gauge-fixing and which lead to the same physical results. One shifts some of the contribution (to the scattering process) from the three-point vertex to the propagator; similar shifts are done for the higher point functions as well. The amount of shift is determined by requiring the physical results to be the same and leads to the pinching procedure [8]. For our gauge choice, the pinching terms arise from the diagrams shown in fig 2. (A quark scattering process suffices to identify these terms.)




fig1.- Gluon self-energy




fig2.- Pinching diagrams.

The total pinching contribution to the two-point function is

$$\tilde{\Gamma}(K) = \frac{C e^2 m}{16} K \int_0^1 d^d p \frac{5}{K^2 (p^2 + 1) + 2} - \frac{2}{K^2 (p^2 + 2)} \quad (17)$$

This is to be added to $\Gamma(K)$ to obtain the gauge-independent two-point vertex function.

From the explicit factor of K in Eq.(17), we see that the pole remains at $K = 0$.

We shall now briefly consider two-loop corrections. These must involve e^4 and since e^2 has the dimension of mass, the two-loop contribution to B has the form

$$B^{(2\text{-loop})} = \frac{(C e^2)^2}{M^2} (\alpha_1 + \alpha_2 \log(T=M)) \quad (18)$$

where $\alpha_1; \alpha_2$ are pure numbers. The gap equation now reads

$$x^2 = \frac{2.384}{4} x + \alpha_1 + \alpha_2 \log(T=M) \quad (19)$$

$M = C e^2 x$. The numerical values of $\alpha_1; \alpha_2$ determine whether the two-loop corrections are small. There are seventy-nine diagrams for α_{ij} at two-loop level as opposed to five at the one-loop level. Since the vertices given by S_m are fairly complicated we have not completed the two-loop calculation. A preliminary analysis of some of the diagrams suggests that the two-loop effects may be smaller by a factor of three or four. It should be noted that this issue does not impinge on the use of S_m as a gauge-invariant infrared cutoff using the above procedure. It only affects the numerical determination of M in a loop expansion.

References

- [1] R. Efraty and V.P. Nair, Phys.Rev.Lett. 68 (1992) 2891; Phys.Rev. D 47 (1993) 5601.
- [2] R. Pisarski, Physica A 158 (1989) 246; Phys.Rev.Lett. 63, (1989) 1129; E. Braaten and R. Pisarski, Phys.Rev. D 42 (1990) 2156; NuclPhys. B 337 (1990) 569; ibid. B 339, (1990) 310; Phys.Rev. D 45 (1992) 1827.
- [3] J. Frenkel and J.C. Taylor, NuclPhys. B 334 (1990) 199; J.C. Taylor and S.M.H. Wong, NuclPhys. B 346 (1990) 115.
- [4] R. Jackiw and V.P. Nair, Phys.Rev. D 48 (1993) 4991.
- [5] J.P. Blaizot and E. Iancu, Phys.Rev.Lett. 70 (1993) 3376; NuclPhys. B 417 (1994) 608.
- [6] V.P. Nair, Preprint CCNY-HEP 4/94.
- [7] A. Bilbire, G. Lazarides and Q. Shaha, Phys. Lett. 103 B (1981) 450; O.K. Kalashnikov, JETP Lett. 39 (1984) 405; T.S. Biro and B. Muller, NuclPhys. A 561

(1993) 477; O . Philipsen, in Proceedings of the NATO workshop, Sintra, Portugal, March 1994 (to be published); W . Buchmuller and O . Philipsen, Preprint DESY -94-202 (November 1994).

[8] JM . Cornwall, Phys.Rev. D 26 (1982) 1453; JM . Cornwall and J. Papavassiliou, Phys.Rev.D 40 (1989) 3474.