

CERN {TH /95-86
 ENSLAPP-A-514/95
 ISN 95{28
 ITKP 95{11

c
 AND OTHER CHARMED BARYONS
 REVISTED

A ndre M artin

CERN , Theory D ivision
 CH {1211 G eneve 23

and

Laboratoire de P hysique T heorique ENSLAPP[?]
 G roupe d'A nnecy, LAPP, B .P . 110
 F {74941 A nnecy-le-V ieux, France

Jean-M arc R ichard^y

Institut des Sciences N ucléaires{CNRS{IN 2P 3
 U niversité Joseph Fourier
 53, avenue des M artys, F-38026 G renoble, France

and

Institut für T heoretische K emphysik
 Rheinische Friedrich-W ilhelm s U niversität
 Nu allee 14{16, D {52115 Bonn, G erm any

A bstract

The mass of the Λ_c^0 baryon with quark content (ssc) is computed in a potential model whose parameters have been determined in 1981 by fitting the spectrum of heavy mesons. It is found in perfect agreement with a recent measurement at the CERN hyperon-beam experiment. The spectroscopy of other charmed baryons in potential models is briefly reviewed.

^y Supported by a A . von Humboldt French-G erm an Research Grant

[?] URA 14-36 du CNRS associée à l'Ecole Normale Supérieure de Lyon et à l'Université de Savoie

CERN {TH /95-86

March 25, 2024.

Many potential models have been successfully tuned to reproduce the spectrum and static properties of heavy quarkonia. A key property is flavour independence: the same potential holds for various quark-antiquark systems. Small corrections are however expected, in particular in the spin-dependent part. They arise when reducing any relativistic kernel into the Schrodinger framework.

One of these models with flavour independence built in is the simple power-law potential designed by one of us [1]

$$V = -8.064 + 6.8698 r^{0.1}; \quad (1)$$

where units are GeV for the potential V , and GeV⁻¹ for the interquark distance r . This is a variant of the logarithmic potential which produces the same spacings for all $Q\bar{Q}'$ spectra [2]. The central potential (1) is supplemented by a spin-spin term of contact type

$$V_{SS} = 1.112 \frac{1}{m_1 m_2} \vec{s}_1 \cdot \vec{s}_2 \delta^3(\vec{r}); \quad (2)$$

where the m_i are the constituent masses. V_{SS} is treated at first order, and is adjusted to reproduce the correct $J=1$ hyperfine splitting of charmonium. The quark masses are

$$m_s = 0.518; \quad m_c = 1.8; \quad m_b = 5.174 \text{ GeV}; \quad (3)$$

This model has been very successful in reproducing the masses of the cc , bb , ss and cs bound states and its predictions for the bs system have been checked experimentally [3].

The model of Eqs.(1-3) was applied in Ref. [4] to compute the mass of the $\psi(3100)$, using the semi-empirical rule

$$V_{Q\bar{Q}} = \frac{1}{2} V_{Q\bar{Q}'}; \quad (4)$$

which is discussed in [4], and references therein. The result, 1662 MeV (actually now 1666 MeV from an improved computation of the hyperfine contribution), comes very close to the experimental value $M(\psi(3100)) = 1672 \text{ MeV}$. New, and accurate determinations of the mass of the $\psi(3100)$ are expected, and preliminary results are already reported. A value

$$M(\psi(3100)) = 2706.8 \pm 1 \text{ MeV} \quad (5)$$

was presented at the Rencontres de Moriond, in march 1995, by the WA89 collaboration [5], which uses the CERN hyperon beam [6]. This value has to be compared with 2740 \pm 20 MeV by the WA62 CERN experiment [7], 2719 \pm 7 \pm 2.5 MeV by ARGUS [8], and 2705.9 \pm 3.3 \pm 2.0 MeV by the E687 experiment at Fermilab [9]. For a review on experimental and theoretical aspects of heavy baryons, see, e.g., [10].

It seems interesting to repeat the calculation of [4] for (ssc) configurations with spin $1/2$ (ψ_c), and $3/2$ (ψ_c'). We find

$$M(\psi_c) = 2708; \quad M(\psi_c') = 2760 \text{ MeV}; \quad (6)$$

Table 1: Comparison of predictions for the masses of Λ_c and Σ_c baryons (in MeV).

Authors	Ref.	Λ_c		Σ_c	
Roncaglia et al.	[12]	2710	30	2770	30
Samuele et al.	[13]	2717	25	2767	35
Izatt et al.	[14]	2610		2710	
Rho et al.	[15]	2786		2811	
de Rujula et al.	[16]	2680		2720	
Malmann et al.	[17]	2730		2790	
Chan	[18]	2773		2811	
Richard et al.	[19]	2664		2775	
Silvestre-Brac	[20]	2675		2749	
Present work		2708		2760	

using an hyperspherical expansion up to a "grand" orbital momentum $L = 8$ [11]. Hence, it is expected that Λ_c^0 will decay into Λ_c^- with emission of a photon of about 52 MeV.

Other predictions from the literature are listed in Table 1. The estimate of Roncaglia et al. [12] does not result from a specific model, but from a survey of the regularities of the hadron spectrum in flavour space. Not surprisingly, it comes very close to the (preliminary) experimental mass. Ref. [13] is a lattice calculation, [14] a bag model. The others are potential models. The closest to the present one is [19], where the same functional dependence as in Eqs. (1,2) is used, but, there, it is attempted to treat all baryons, even those with light quarks, and this results in a larger strength for the hyperfine correction.

On the same line as [12], one can derive inequalities which do not depend on the specific choice of the potential V , and would in fact hold for any flavour-independent model. Examples were already given for beautiful baryons [21] and for light flavour [11]. We assume that the interquark potential in baryons and quarkonia satisfy

$$V_{QQQ}(\mathbf{r}_1; \mathbf{r}_2; \mathbf{r}_3) \geq \frac{1}{2} \sum_{i < j} V_{Q\bar{Q}}(\mathbf{r}_i - \mathbf{r}_j); \quad (7)$$

as for instance with the prescription (4), or with the string-motivated model [22]

$$V_{Q\bar{Q}} = \kappa r; \quad V_{QQQ} = m_J \ln(d_1 + d_2 + d_3); \quad (8)$$

where d_i is the distance from the i^{th} quark to a junction J whose location is adjusted to minimize the potential.

From Eq. (7), one easily derives an inequality between spin-averaged ground-state masses [11]

$$M(\text{ssc}) \geq \frac{1}{2} M(\text{ss}) + M(\text{cs}); \quad (9)$$

This lower bound can be estimated near 2.5 GeV, and is rather crude. It can be improved in two ways. First one introduces hyperfine corrections, assuming a linear dependence of the Hamiltonian upon the spin operators $\sim \vec{1} \cdot \vec{\gamma}$. As shown in [21], one gets

$$M(\chi_c) = \frac{1}{2}M(\chi) + \frac{3}{4}M(D_s) + \frac{1}{4}M(D_s^*) : \quad (10)$$

One can also use the Hall-Post techniques [23] to remove the centre-of-mass energy left over in 2-body subsystems when deriving (9). Let us simplify the notations into $m_s = 1$ and $m_c = M$, with typically $M \sim 3-4$. The kinetic energy of the baryon

$$T = \frac{\vec{p}_1^2}{2} + \frac{\vec{p}_2^2}{2} + \frac{\vec{p}_3^2}{2M} \quad (11)$$

can be rewritten as [24]

$$T = \frac{(\vec{p}_1 + \vec{p}_2 + \vec{p}_3)^2}{2(1+x)} + \frac{(\vec{p}_1 + \vec{p}_2)^2}{2(1+x)} + \frac{\vec{p}_3^2}{2M} ; \quad (12)$$

where, for given x , the inverse masses are

$$a_{23} = a_{31} = \frac{1+x}{1+2x} \left(1 + \frac{2}{M} \right) ; \quad a_{12} = \frac{8}{(1+2x)^2} x(1+x) \frac{1}{2M} ; \quad (13)$$

This gives the following inequality on Hamiltonians

$$H_3 = T + V_{QQ} \geq \frac{1}{2} \sum_{i < j} H_2(a_{ij}; V_{QQ}) ; \quad (14)$$

where $H_2(a; V) = \frac{a^2}{2} + V$. This implies an inequality on ground-state energies

$$E_3(ssc) \geq \sum_{i < j} E_2[a_{ij}] ; \quad (15)$$

The l.h.s. can be optimized by varying x , leading to inverse masses a_{ij} which are larger than the inverse masses $b_{ij} = (m_i^{-1} + m_j^{-1})/2$ in actual ss or cs mesons. The corresponding change of binding energy can be bounded in terms of the orbital excitation energy $E = E_2(1P) - E_2(1S)$. The result is [25]

$$E_2[a] - E_2[b] \geq \frac{3}{4} \frac{a - b}{a} E ; \quad (16)$$

with mild restrictions on the $Q\bar{Q}$ potential, $V > 0$, and $V'' < 0$, i.e., a behaviour in between Coulomb and linear. If one estimates E from the experimental information

on positive-parity ss and cs mesons (with some uncertainty for estimating the spin-averaged masses), and varies the parameter x , one ends with a lower limit

$$M(\psi_c) > 2.65 \text{ GeV} : \quad (17)$$

The main conclusion of this study is that the $1/2$ rule works surprisingly well for relating mesons to baryons. The ψ and ψ_c masses are well reproduced. The predictions for the ground state with other flavour combinations are

$$M(\psi_{cc}) = 3.737; \quad M(\psi_{cc}^2) = 3.797; \quad M(\psi_{ccc}) = 4.787 \text{ GeV} : \quad (18)$$

There are reasonable expectations that these states, "the ultimate goal of baryon spectroscopy" [26, 27] would be seen at LHC.

Acknowledgments

One of us (A.M.) would like to repeat his thanks to Murray Gell-Mann for suggesting him to include strange quarks in potential models. He would also like to thank Tran Thanh Van for the hospitality at the Rencontres de M. Orind.

References

- [1] A. Martin, Phys. Lett. 100B (1981) 511.
- [2] C. Quigg and J.L. Rosner, Phys. Lett. 71B (1977) 153;
M. Machacek and Y. Tomozawa, Ann. Phys. (N.Y.) 110 (1978) 40.
- [3] A. Martin, preprint CERN-TH 7352/94, to appear in Symmetry and Simplicity in Physics, (A symposium on the occasion of Sergio Fubini's 65th birthday), Eds. W.M. Alberico and S. Sciuto, World Scientific (1994).
- [4] J.-M. Richard, Phys. Lett. 100B (1981) 515.
- [5] F. Droppmann (WA89 collaboration), Rencontres de M. Orind (1995), to be edited by J. Tran Thanh Van (Editions Frontieres).
- [6] Roland Werdinger et al., Production of Charmed Baryons in the CERN Hyperon Beam Experiment WA89, presented at 27th International Conference on High Energy Physics (ICHEP), Glasgow, Scotland, 20-27 Jul 1994.
- [7] S.F. Biagi et al. (WA62 collaboration), Z. Phys. C 28 (1985) 175.
- [8] H. Albrecht et al. (ARGUS collaboration), Phys. Lett. B 288 (1992) 367.
- [9] P.L. Frabetti et al. (E687 collaboration), Phys. Lett. B 300 (1993) 190.
- [10] J. Komer, M. Kramer, and D. Pirjol, Prog. Part. Nucl. Phys. 33 (1994) 787.

- [11] J.-M. Richard, Phys. Rep. 212 (1992) 1.
- [12] R. Roncaglia, D. B. Lichtenberg, and E. Predazzi, Predicting the masses of baryons containing one or two heavy quarks, preprint hep-ph 9502251.
- [13] S. Samuel and K. J. Moriarty, Phys. Lett. B 175 (1986) 197.
- [14] D. Izatt, C. DeTar and M. Stephenson, Nucl. Phys. B 199 (1982) 269.
- [15] M. Rho, D. O. Riska and N. N. Scoccola, Phys. Lett. B 251 (1990) 597.
- [16] A. de Rújula, H. Georgi and S. L. Glashow, Phys. Rev. D 12 (1975) 2060.
- [17] K. Maltman and N. Isgur, Phys. Rev. D 22 (1980) 1701.
- [18] L. Chan, Phys. Rev. D 15 (1977) 2478.
- [19] J.-M. Richard and P. Taxil, Phys. Lett. B 128 (1983) 453.
- [20] B. Silvestre-Brac, Spectrum and static properties of heavy baryons, Grenoble preprint ISN 94-60, to appear in Few Body Systems, and private communication.
- [21] A. Martin and J.-M. Richard, Phys. Lett. B 185 (1987) 426.
- [22] H. G. Dosch and V. F. Muller, Nucl. Phys. B 116 (1976) 470;
D. Gromes and I. O. Stamatescu, Z. Physik C 3 (1979) 4;
P. Hasenfratz et al., Phys. Lett. 94 (1980) 401;
J. Carlson, J. Kogut and V. R. Pandharipande, Phys. Rev. D 27 (1983) 233;
D 28 (1983) 2807.
- [23] R. L. Hall and H. R. Post, Proc. Phys. Soc. 90 (1967) 381.
- [24] J.-L. Basdevant, A. Martin, J.-M. Richard and T. T. Wu, Nucl. Phys. B 393 (1993) 111.
- [25] J.-L. Basdevant, A. Martin and J.-M. Richard, Nucl. Phys. B 343 (1990) 69.
- [26] B. J. Bjorken, in Proc. Int. Conf. on Hadron Spectroscopy, College Park, 1985, ed. S. Oneda (A.I.P., 1985).
- [27] M. J. Savage and M. B. Wise, Phys. Lett. B 248 (1990) 177;
M. J. Savage and R. P. Springer, Int. J. Modern Phys. A 6 (1991) 1701.