

Phenomenology of x_F Dependence of Quarkonium Production in Proton-Nucleus Interactions

W. Q. Chao^{1,2,3}, C. S. Gao^{1,3,4} and Y. B. He^{1,4}

1. China Center of Advanced Science and Technology (World Laboratory),

P. O. Box 8730, Beijing 100080, China

2. Institute of High Energy Physics, Academia Sinica, P. O. Box 918(4), Beijing 100039, China

3. Institute of Theoretical Physics, Academia Sinica, P. O. Box 2735, Beijing 100080, China

4. Physics Department, Peking University, Beijing 100871, China

Abstract

We present a phenomenological study of the x_F dependence of quarkonium production in high energy proton-nucleus collisions. The x_F dependence of comover contributions is introduced to account for the observed quarkonium suppression at low x_F . Combining comover contributions, nuclear shadowing effect, energy loss mechanism and nuclear absorption together we reproduce the overall x_F dependence of E 772/E 789 data.

Typeset using REVTeX

Partly supported by the National Science Foundation of China

I. INTRODUCTION

A strong suppression of charmonium ($J = \psi, \psi'$) and bottomonium ($1S, 2S+3S$) production has been observed in $p-A$ collisions on heavy relative to light nuclear targets [1,4]. Special attentions have been paid to the significant suppressions at low (small or negative) and high x_F regions. The suppression at low x_F is usually believed to come from nuclear absorption and comover contributions. The maximum contribution of nuclear absorption to the $J = \psi$ suppression was studied in ref. [5], and it was found that nuclear absorption alone cannot account for the $J = \psi$ suppression measured in either $p-A$ or $A-B$ collisions. Since at E 772 energy most of the physical bound states are formed well outside target nucleus due to the Lorentz dilation of the formation time, nuclear absorption can contribute little to the observed suppression at low x_F region. While comovers can continue to interact well outside of the nuclear volume, they may in principle play a more important role in proton-nucleus collisions, although the number of comovers in $p-A$ reactions is not so large as in $A-B$ collisions. In the literature comover contributions are generally estimated in the central rapidity region without x_F dependence. To our knowledge there is no direct explanation of the E 772/E 789 data at $x_F \neq 0$ and below [3,4]. In this work we use a simple model to extrapolate the rapidity distribution of comovers for $p-A$ reactions from $p-p$ data [6], introducing the x_F dependence of comover contributions, to explain the E 772/E 789 data.

On the other hand, several mechanisms have been proposed to account for the suppression at large x_F . Among these, nuclear shadowing effect alone is argued to be unable to explain the data [3]. The intrinsic charm model [7,8] was expected to explain this x_F dependence. However, E 789 data of $J = \psi$ production at very large x_F [9] show no evidence of intrinsic charm contribution. The energy loss mechanism [10] could account for the observed suppression at large x_F . Nevertheless, the amount of energy loss needed to explain the E 772 $J = \psi$ data appears to be significantly larger than the value determined from other considerations [11]. In our work, we combine energy loss mechanism together with nuclear shadowing effect, nuclear absorption and comover contributions, and reproduce the large

x_F data with a smaller amount of energy loss.

II. COM OVER CONTRIBUTIONS

In the case of quarkonium production, e.g., J/ψ production, in proton-nucleus collisions, comoving secondary particles can scatter with the cc pair, and contribute to the observed J/ψ suppression. These comover contributions have been studied in the literature [12,14,8]. However, since the shape of the rapidity density of the comovers in inclusive J/ψ production for p-A collisions is unknown, comover density is usually estimated in the central rapidity region, and taken as a constant independent of the comover rapidity. In this work we will use a simple model to extrapolate the rapidity distribution of comovers for p-A reactions from p-p data [6]. Assuming that comovers with rapidity close to the rapidity of a cc pair can cause the breakup of the cc pair, we translate the rapidity distribution of comovers into x_F distribution, and include the resulting x_F dependence of comover contributions to explain the E772/E789 low x_F data. We will follow the work of refs. [8,14], and improve the treatment of comover contributions.

Including only the comover contributions, the x_F dependent cross section of J/ψ production in p-A collisions can be expressed as

$$\frac{d_{pA}}{dx_F} = \frac{d_{pp}}{dx_F} \int_0^1 d^2b \int_{-1}^1 dz_A (b;z) \exp \left[- \int_0^z d\tau \sigma_{cc-com} v(\tau;b) \right]; \quad (1)$$

where ρ_A is the nuclear density profile, σ_{cc-com} is the (cc) -comover absorption cross section, τ_0 is the formation time of comovers, τ_f is the effective proper time over which the comovers can interact with cc pair, v is the relative velocity of cc with the comovers, and $n(\tau;b)$ is the density of comovers at the proper time τ and impact parameter b .

Assuming Bjorken's hydrodynamics [15], the comover density varies with the proper time according to,

$$n(\tau) = \frac{n(\tau_0)}{\tau}: \quad (2)$$

The comovers consist mostly of π^+ , π^- and K^0 mesons. We relate n , the density of comovers, to n_{ch} , the density of produced charged particles, by $n = f n_{ch}$. For example, $f = 1.5$ if comovers are all π^+ 's with equal numbers of π^- , π^0 and K^0 .

The charged particle density at y_0 can be related to rapidity distribution by

$$n_{ch}(y_0; b) = \frac{1}{\sigma_{in}^0} \frac{dN_{ch}^{pA}(b)}{dy}; \quad (3)$$

where $\sigma_{in}^0 = 30 \text{ mb}$ is the nucleon-nucleon inelastic scattering cross section.

Now we use the so called independent cluster model [16] to extrapolate the charged particle rapidity distribution for $p-A$ reactions from $p-p$ data [6].

We fit the data at $\sqrt{s} = 45.2 \text{ GeV}$, which is close to the E772 energy, in the form,

$$\frac{dN_{ch}^{pp}}{dy} = W \exp^{n(y+y_0)^2} + W \exp^{n(y-y_0)^2} \quad (4)$$

with $W = 1.75$, $y_0 = 1.5$ and $n = 2.9$.

In the framework of the independent cluster model, we have

$$\frac{dN_{ch}^{pA}(b)}{dy} = (b)W \exp^{n(y+y_0)^2} + W \exp^{n(y-y_0)^2}; \quad (5)$$

where $(b) = \int_{-R}^R dz_A (b; z)$ is the number of target participants at the impact parameter b .

In order to introduce the x_F dependence of comover contributions, we assume that those comovers with rapidity close to the rapidity of a $J=$ can cause the breakup of the $J=$, and we relate the rapidity of $J=$ in the center-of-mass system to x_F , which is defined as $x_F = P_L / (P_L)_{max}$, by

$$y = \frac{1}{2} \ln \frac{E + P_L}{E - P_L}; \quad (6)$$

where E and P_L are the energy and longitudinal momentum of $J=$ in the c.m.s..

In our approach to the x_F dependent comover density two main parameters are f and y_0 . In Fig.1 we show the x_F dependence of comover densities for various sets of f and y_0 . Note that the dotted curve with $f = 1.5$, $y_0 = 0.6$ and $\sigma_0 = 0.8 \text{ fm}$ is plotted in view of the

argument of ref. [12], which represents a com over density too low to explain the data. In our further calculations we will $x_F = 1.5$. The com over formation time has been estimated to be about $0.4 - 1.2 \text{ fm} = c$, the time scale of both quark and hadron formation in the early stages of the collision [17,18]. It can be seen from Fig.1 that the com over density is sensitive to the choice of ρ_0 . In the next section we will vary ρ_0 and ρ_∞ to attribute the observed suppression at low x_F region to com over contributions.

Since com overs can interact with the produced heavy quarks Q or $\bar{Q}$ before the quarkonium bound state can be formed, and the interaction is independent of whether or not the heavy quark pair is produced as a color octet or singlet, we expect that for charmonium production the (cc)-com over interaction cross section is the same for the $J=$ and 0 bound states. Furthermore, the com over contributions are expected to be independent of the final hadronic size. As a result, although we expect the occurrence of significant com over absorptions with strong nuclear dependence, the $J=$ and 0 production would have the same nuclear dependence under com over absorption, which is consistent with the earlier experimental observation [3].

III. COMBINATION OF MECHANISMS, NUMERICAL RESULTS AND DISCUSSIONS

In this section we combine com over contributions, nuclear absorption, nuclear shadowing effect and energy loss mechanism to reproduce the overall x_F dependence of E772/E789 data on $J=$ and 0 production.

We will calculate x_F dependence of σ for $J=$ and 0 production in the parametrization,

$$\frac{d \sigma^{pA}}{dx_F} = \frac{d \sigma^{pp}}{dx_F} A^{(\alpha_F)} : \quad (7)$$

For $J=$ production, we make use of the fact that about 60% of the final $J=$'s are produced directly, about 30% come from c intermediate states and the remaining 10% are produced through the decay of 0 . Analogously, about 41% of the final $1S$'s are produced directly,

about 35% and 18% come from the decay of ψ and ψ^0 , about 5% and 1% are produced through the decay of χ_{2S} and χ_{3S} [19].

The cross section of quarkonium production in p-A collisions is

$$\frac{d_{\text{ppA}}}{dx_F} = \frac{d_{\text{pp}}}{dx_F} z^{\frac{9}{8}} d^2 b_1 dz_A(b; z) \exp \left[-z^{\frac{1}{8}} dz_A^0(b; z^0)_{\text{abs}}(z^0, z); \right. \\ \left. \exp \int_0^Z d_{\infty} v n(\cdot; b) \right] \quad (8)$$

where several mechanisms have been included.

The averaged nucleon-nucleon cross section of pA reactions, σ_{NN}^{pA} in eq. (8), is the same as the bare nucleon-nucleon cross section except that in pp the distribution functions of the gluon, quark and antiquark within the target nucleus are nuclear modified. We follow the work of ref. [20] to incorporate the shadowing effect into σ_{NN}^{pA} , and use parameters determined in ref. [8].

The first exponential in eq. (8) represents the contribution of nuclear absorption. We take the values of parameters in the following way: the formation time of charmonia and bottomonia, $\tau = 0.89$ fm, $\tau = 0.76$ fm, etc., is taken from ref. [21], the absorption cross section between resonances and nucleon is chosen as

$$R_N = \sum_N (r_N - \bar{r})^2; \quad (9)$$

with $\sigma_N = 6 \text{ mb}$, and the radii of resonances r_R taken from ref. [21]. In particular, we want to point out that with the reason discussed in the first section, the nuclear absorption can contribute very little suppression even with a rather large absorption cross section $\sigma_N = 6 \text{ mb}$, as indicated by the fact that the nuclear dependence of J/ψ and ψ' production is the same within errors [3].

We follow the work of ref. [10] to include the contribution of energy loss mechanism in our calculations. In ref. [10] only energy loss mechanism is taken into account, and an upper bound of energy loss is obtained, $dE/dZ \sim 1.5 \text{ GeV/fm}$. Moreover, the energy loss contribution in ref. [10] has not been normalized, which implies an integrated suppression. In our calculations, we complete the normalization in a similar way as in ref. [22],

$$\frac{d^{pA}}{dx_F}(x_F)_{\text{normal}} = -\frac{1}{dx_F} \frac{d^{pA}}{dx_F}(x_F = 0); \quad (10)$$

where $\gamma = \frac{x_F}{x_F + \Delta x_F}$, and Δx_F is the shift of x_F due to the energy loss. Since nuclear shadowing effect can also contribute to the quarkonium production suppression at large x_F , in our calculation a value of $dE = dZ = 0.5 \text{ GeV/fm}$ is found to be able to fit the data quite well, which is consistent with the value obtained from other considerations [11].

The comover contributions are included in the second exponential in eq.(8). We take the effective proper time $\tau_f = r_0/c_s$, where r_0 is the projectile radius which we take to be 1.2 fm , and $c_s = \frac{1}{3} \frac{p}{m}$ [8].

The results of our calculations are compared to E772/E789 data for J/ψ production at 800 GeV/c in Figs 2 and 3, with different sets of σ_{∞} and ρ_0 . We find that $\sigma_{\infty} = 4.0 \text{ mb}$ and $\rho_0 = 0.8 \text{ fm}^{-3}$ is the best fit, which corresponds to a comover density with maximal value about 6 fm^{-3} . Fig 2 shows that a lower density (larger ρ_0) is not sufficient to explain the data. In Fig 3 we see that a comover absorption cross section σ_{∞} smaller than 4.0 mb is also insufficient to fit the data. Note that the dot-dashed curve in Fig 3 corresponds to the case without comover contributions. The curve illustrates a very small contribution of nuclear absorption at low x_F , as we expected. One can find in Fig 3 that although the number of comovers is not very large in $p-A$ reactions, comover interactions have a much more important effect than nuclear absorption.

A comparison of our calculations for J/ψ production with the E772 data is shown in Fig 4. The dashed curve with $\sigma_{\infty} = 4.0 \text{ mb}$ and $\rho_0 = 0.8 \text{ fm}^{-3}$ is somehow lower than the experimental points at $x_F < 0.3$. This may imply our over-estimate of comover contributions when we fit the data at low x_F region.

In the literature studies of nucleus-nucleus reactions indicated that the comover density at an early stage of the collision is very high, possibly from 1 to 5 fm^{-3} . In our work, a comover density with a maximum 6 fm^{-3} is used when we fit the E772/E789 data at low x_F . With such a high density one would rather think of these comovers as light quarks and antiquarks than as pions and low-mass resonances [14]. However, whether it is possible to

obtain such a high density in proton-nucleus collisions appears questionable. Presumably some new mechanism might also contribute at low x_F region.

According to ref. [23], absorption in confined hadronic matter is excluded as a possible cause of the J/ψ suppression observed in nucleus-nucleus collisions. Since at low x_F region neither energy loss nor nuclear shadowing effect is important, it seems difficult to understand the low x_F suppression in proton-nucleus collisions, in which no deconfined matter is expected. Our work shows that comover interactions play an important role at the low x_F region, and a comover absorption cross section as large as 4 mb is needed to confront the E772/E789 data. In addition, a recent work observed that nuclear antishadowing leads to enhanced J/ψ production at low x_F , which is in contradiction with the E772/E789 data [24]. These issues suggest that the low x_F suppression in $p-A$ reactions deserves careful studies, and possibly requires some new mechanism.

To summarize, we have reproduced the overall x_F dependence of quarkonium production in proton-nucleus collisions, by combining comover contributions, nuclear shadowing effect, nuclear absorption and energy loss mechanism together. In particular, by including the x_F dependence of comover distribution we show that comover interactions have a very important effect when one attempts to explain the observed quarkonium suppression, especially at low x_F region, in proton-nucleus collisions. However, our work indicates that a fairly high density of comovers is needed to explain the E772/E789 low x_F suppression, which might imply some new mechanism occurs at this region as well.

ACKNOWLEDGMENTS

The authors would like to thank Q.H. Zhang and H.M. Hu for helpful discussions.

REFERENCES

- [1] J. Badier et al., Z. Phys. C 20, 101 (1983).
- [2] S. Kartik et al., Phys. Rev. D 41, 1 (1990).
- [3] D. M. A. Ide et al., Phys. Rev. Lett. 66, 133 (1991).
- [4] M. J. Leitch et al., Nucl. Phys. A 544, 197c (1992).
- [5] S. Gavin, H. Satz, R. L. Thews and R. Vogt, Z. Phys. C 61, 351 (1994).
- [6] W. Thome et al., Nucl. Phys. B 129, 365 (1977).
- [7] S. J. Brodsky and P. Hoyer, Phys. Rev. Lett. 63, 1566 (1989).
- [8] R. Vogt, S. J. Brodsky and P. Hoyer, Nucl. Phys. B 360, 67 (1991).
- [9] M. S. Kowitt et al., Phys. Rev. Lett. 72, 1318 (1994).
- [10] S. Gavin and J. Milana, Phys. Rev. Lett. 68, 1834 (1992).
- [11] S. J. Brodsky and P. Hoyer, Phys. Lett. B 298, 165 (1993).
- [12] S. Gavin, M. Gyulassy and A. Jackson, Phys. Lett. B 207, 257 (1988).
- [13] R. Vogt, M. Prakash, P. Koch and T. H. Hansson, Phys. Lett. B 207, 263 (1988); J. Ftacnik, P. Lichard, and J. Pisut, Phys. Lett. B 207, 194 (1988).
- [14] S. Gavin and R. Vogt, Nucl. Phys. B 345, 104 (1990).
- [15] J. D. Bjorken, Phys. Rev. D 27, 140 (1983).
- [16] W. Q. Chao, M. K. Hegab and J. Hufner, Nucl. Phys. A 395, 482 (1983).
- [17] S. J. Brodsky and A. H. Mueller, Phys. Lett. B 206, 685, (1988).
- [18] C. Y. Wong, Introduction to High-Energy Heavy-Ion Collisions, Oak Ridge National Laboratory, 1994.

- [19] G . P iller, J.M utzbauer and W . W eise, Nucl.Phys.A 560, 437 (1993).
- [20] J. Q iu, Nucl.Phys.B 291, 746 (1987).
- [21] F.K arsch and R .Petronzio, Z.Phys.C 37, 627 (1988).
- [22] D .K harzeev and H .Satz, Z.Phys.C 60, 389 (1993).
- [23] D .K harzeev and H .Satz, Phys.Lett.B 334, 155 (1994).
- [24] S.L iuti and R .Vogt, LBL-35821, hep-ph/9407272.

Figure Captions

Fig.1. x_F dependence of com over densities for different sets of f and ρ_0 .

Fig.2. x_F dependence for $J=\psi$ production from E 772/E 789 [3,4]. The curves compare our calculations to the data, with a fixed $\sigma_{\infty} = 4.0$ mb and various values of ρ_0 .

Fig.3. x_F dependence for $J=\psi$ production from E 772/E 789 [3,4]. The curves compare our calculations to the data, with a fixed $\rho_0 = 0.8$ fm⁻³ and $\sigma_{\infty} = 4.0$ mb; 2.0 mb. The dot-dashed line corresponds to the case without com over contributions.

Fig.4. Our calculations of x_F dependence for bottomonium production with different sets of σ_{∞} and ρ_0 are compared to the E 772 data [3].

Fig. 1




Fig. 2




Fig. 3




Fig. 4

