

Multiloop Calculations in Heavy Quark Effective Theory

D.J.BROADHURST^y and A.G.GROZIN^z
 Physics Department, Open University
 Milton Keynes, MK7 6AA, UK

We review algorithmic methods for two-loop calculations in HQET, and the analogous methods for on-shell QCD, needed for matching HQET to QCD.

1. Introduction

In this talk, we review methods for two-loop calculations in heavy quark effective theory¹. In HQET, the heavy quark propagator is $S(k) = 1/(k \cdot v + i0)$, and the quark-gluon vertex is $igt^a v$, with a heavy quark velocity $v = (1; 0)$, in the rest frame. We shall discuss algorithmic methods, suitable for computer algebra implementation. All of them are based on dimensional regularization, with space-time dimension $d = 4 - 2\epsilon$, and integration by parts², a powerful calculational method based on the simple observation that integrals of full derivatives vanish. Tensor integrals are handled by decomposition into a suitable tensor basis, with coefficients that can be expressed in terms of scalar integrals, by solving linear systems of equations. In all cases which we shall discuss, numerators can be expressed in terms of factors of the denominator.

In Section 2, we recall the well-known integration-by-parts algorithm for calculation of massless two-point integrals². This example is used to introduce notations and to set up the pattern which will be repeated in other cases. In Section 3, we discuss the corresponding method in HQET³. This method was generalized⁴ to HQET three-point integrals, expanded in the velocity change $v^0 \rightarrow v$, to any finite order, as discussed in Section 4. In order to obtain QCD/HQET matching conditions⁵, it is necessary to equate QCD and HQET on-shell matrix elements. The latter are trivial (apart from massive quark loops). The former require methods for on-shell massive QCD calculations^{6;7;8;9}, discussed in Section 5.

2. Massless two-point integrals

We write the massless one-loop integral of Fig. 1a, with arbitrary indices, as

$$\int \frac{d^d k}{(2\pi)^d} \frac{1}{(k^2)^{\alpha_1} ((k+p)^2)^{\alpha_2}} = \frac{i(-p^2)^{d/2-\alpha_1-\alpha_2}}{(4\pi)^{d/2}} G(-\alpha_1; -\alpha_2): \quad (1)$$

¹ Talk presented by A.G.Grozin at the A'HE'NP'95 workshop, Pisa, April 1995

^y D.Broadhurst@open.ac.uk

^z A.Grozin@open.ac.uk;

on leave of absence from Budker Institute of Nuclear Physics, Novosibirsk, Russia

It can be calculated by using the Feynman parameterization

$$\frac{1}{a+b} = \frac{(1+x)}{(1-x)} \int_0^1 \frac{x^{1-1} (1-x)^{1-1} dx}{[ax + b(1-x)]^+} \quad (2)$$

or by using Fourier transformation to coordinate space and back. The result is

$$G(1; 2) = \frac{(1+2-d=2) (d=2-1) (d=2-2)}{(1) (2) (d-1-2)}: \quad (3)$$

All one-loop integrals with integer indices can therefore be expressed in terms of $G_0 = (1+2)^2 (1-2) = (1-2)$, with rational coefficients.




Figure 1: Massless two-point diagrams

Two-loop integrals of the form of Fig. 1b are evaluated by repeated use of (3). The integral of Fig. 1c is non-trivial. We write it as

$$\begin{aligned} & \int \frac{d^d k}{(2)^d} \frac{d^d l}{(2)^d} \frac{1}{(k^2)^1 (l^2)^2 (k+l)^3 (1+p)^4 ((k-l)^2)^5} \\ &= \frac{(p^2)^d}{(4)^d} F(1; 2; 3; 4; 5): \end{aligned} \quad (4)$$

Integration by parts gives the recurrence relation

$$(d-1-3-2-5)F = 1^+ (5-2) + 3^+ (5-4) F; \quad (5)$$

where, for example, 1^+ increases 1 , and 2^- decreases 2 . Applying this relation sufficiently many times, we can kill one of the lines 2, 4, or 5, in Fig. 1c. The resulting integrals, in Figs. 1d,e, are trivial:

$$\begin{aligned} F(1; 2; 3; 4; 0) &= G(1; 3)G(2; 4); \\ F(1; 2; 3; 0; 5) &= G(1+2+5-d=2; 3)G(2; 5): \end{aligned} \quad (6)$$

All two-loop integrals can be thus expressed in terms two structures: G_0^2 and $G_1 = (1+2)^3 (1-3) = (1-3)$, with rational coefficients. This algorithm² is implemented in REDUCE^{10,3}, and as part of the 3-loop package Mincer¹¹.

3. HQET two-point integrals

We write the HQET one-loop integral of Fig. 2a, with arbitrary indices, as

$$\int \frac{d^d k}{(2)^d} \frac{1}{(k^2)^1} \frac{1}{k_0 + i0} = \frac{i(2!)^{d-2-1}}{(4)^{d-2}} I(1; 2): \quad (7)$$

It can be calculated, using the parameterization

$$\frac{1}{a+b} = \frac{(\quad + \quad)}{(\quad)(\quad)} \int_0^1 \frac{y^{-1} dy}{(a+by)^+}; \quad (8)$$

where y has dimensions of energy. The result is

$$I(\quad; \quad) = \frac{(2\quad + \quad - d)(d-2\quad - 1)}{(\quad)(\quad)}; \quad (9)$$

All one-loop integrals with integer indices can be expressed in terms of $I_0 = (1 + 2)^{-(1-\epsilon)}$ with rational coefficients.




Figure 2: Two-point diagrams in HQET or on-shell QCD

The two-loop integrals of Figs. 2b,c are trivial. Multiplying the integrand of Fig. 2d by

$$1 = \frac{1}{!} [(k_0 + !) + (l_0 + !) - (k_0 + l_0 + !)]; \quad (10)$$

we can kill one of the heavy lines, reducing this integral to trivial ones, like those of Figs. 2f,g. Only the integral of Fig. 2e is non-trivial. We write it as

$$\begin{aligned} & \int \frac{d^d k}{(2\quad)^d} \frac{d^d l}{(2\quad)^d} \frac{1}{(k^2 - \quad)(l^2 - \quad)((k-l)^2 - \quad)} \frac{!}{k_0 + !} \frac{!}{l_0 + !} \\ &= \frac{(\quad!)^{2(d-\quad-\quad-\quad)}}{(4\quad)^d} I(\quad; \quad; \quad; \quad; \quad); \end{aligned} \quad (11)$$

The recurrence relation

$$\begin{aligned} & (d - \quad - \quad - \quad - 2\quad - 1) I \\ &= \quad! + (5 - 2\quad) + (2(d - \quad - \quad - \quad) - \quad - 4 + 1) 4\quad I \end{aligned} \quad (12)$$

allows us to kill one of the lines 2, 4, or 5, in Fig. 2e. The resulting integrals, in Figs. 2f,g,h, are trivial:

$$\begin{aligned} I(\quad; \quad; \quad; \quad; 0) &= I(\quad; \quad) I(\quad; \quad); \\ I(\quad; 0; \quad; \quad; \quad) &= I(\quad; \quad + \quad + 2\quad - d) I(\quad; \quad); \\ I(\quad; \quad; \quad; 0; \quad) &= I(\quad + \quad + \quad - d; \quad) G(\quad; \quad); \end{aligned} \quad (13)$$

All two-loop integrals can thus be expressed in terms of I_0^2 and $I_1 = (1 + 4\epsilon)^{-2} (1 - \epsilon)$, with rational coefficients.

This algorithm was proposed in³, where it was implemented in REDUCE. It was used for calculation of: the heavy quark field anomalous dimension³ (previously considered in Wilson¹² and on-shell⁷ QCD calculations); the heavy-light current anomalous dimension³ (also obtained using Feynman parameterization¹³ and Gegenbauer polynomial expansion, with a more difficult momentum routing¹⁴); and the correlator of two heavy-light currents^{15,16}. The coefficient of ϵ in this correlator is of order 10, making sum rules for heavy-light mesons unreliable, in our opinion¹⁵.

4. HQET three-point integrals expanded in v^0/v

Here we consider three-point diagrams, containing HQET propagators with two velocities, v and v^0 , and depending on two residual energies, ω and ω^0 . Their expansions in v^0/v to any finite order can be evaluated algebraically. The heavy quark propagators are expanded in v^0/v , and the resulting tensor integrals are reduced to scalar integrals. The one-loop integral of Fig. 2i, with arbitrary indices specifying the dependence on ω and ω^0 , is

$$\int \frac{d^d k}{(2\pi)^d} \frac{1}{(k^2 - \omega^2)} \frac{\omega^0}{k_0 + \omega^0} = \frac{i(2\omega^0)^{d-2}}{(4\pi)^{d-2}} I_{-1;2;3;\frac{\omega^0}{\omega}};$$

$$I(-1;2;3;x) = I(-1;2+3)x^{-2} {}_2F_1(2+2+3-d;2+3;1-x):$$

We write the two-loop integral of Fig. 2e as

$$\int \frac{d^d k}{(2\pi)^d} \frac{d^d l}{(2\pi)^d} \frac{1}{(k^2 - \omega^2)(l^2 - (\omega - l)^2)} \frac{\omega^0}{k_0 + \omega^0} \frac{l^0}{l_0 + \omega^0}$$

$$= \frac{(2\omega^0)^{2(d-1-2-5)}}{(4\pi)^d} I_{-1;2;3;4;5;\frac{\omega^0}{\omega}}: \quad (14)$$

In the Feynman gauge, all diagrams generate scalar integrals with $1 \leq 5$. Moreover, the cases with $3,4 > 1$ can be obtained from $I(-1;2;1;1;5;x)$ by differentiation with respect to ω and ω^0 . Then, either some of the lines are already absent, or one of the lines 1, 2, 4, 5 can be killed by applying the recurrence relation⁴

$$5 + 1 + \frac{\omega^0}{\omega} (2 + 5 + 3 - d) I$$

$$= 5 + 1 + \frac{\omega^0}{\omega} 5 + (1 - 2) + 1 + (2 - 5) + \frac{\omega^0}{\omega} 3 + 4 - I: \quad (15)$$

This produces the trivial integrals of Figs. 2f,h, and the integral of Fig. 2g, given by $I(-5;4)I(-1;3;4+2-5-d;\frac{\omega^0}{\omega})$.

This algorithm was proposed in⁴; we are not aware of a package that implements it. The method was applied for checking¹⁷, at small velocity transfer, the general result for the heavy-heavy current anomalous dimension¹⁸, and for obtaining⁴ the

correlator of two heavy{light currents and one heavy{heavy current, at small $v^0 \sim v$ (subsequently obtained, for all velocity transfers¹⁹, using methods based on differential equations). Again there are large radiative corrections. However, these are largely cancelled in the ratio of the three{point and two{point sum rules, making predictions for the Isgur{Wise function^{4,19} relatively safe.

5. On-shell massive two{point integrals

Expansions of massive two{point diagrams to any finite order in p^2/m^2 can be evaluated algebraically. Such on-shell diagrams are necessary for QCD/HQET matching. We write the one{loop integral of Fig. 2a, with arbitrary indices, as

$$\int \frac{d^d k}{(2\pi)^d} \frac{1}{(k^2)^{-1} ((k+m v)^2 + m^2)^{-2}} = \frac{m^{d-2(1+2)}}{(4\pi)^{d-2}} M(1; 2): \quad (16)$$

It can be evaluated by Feynman parameterization. Alternatively, we can use the inversion $k \rightarrow k/k^2$, after transforming to a dimensionless euclidean loop momentum. This transforms the on-shell massive denominator into an HQET denominator, and we obtain

$$M(1; 2) = I(d-1, 2; 2) = \frac{(1+2-d-2)(d-2-1-2)}{(2)(d-1-2)}: \quad (17)$$

All one{loop integrals with integer indices can be expressed in terms of $M_0 = (1+)$ with rational coefficients.

There are two distinct non-trivial topologies for two{loop integrals. We shall start with the simpler type M, of Fig. 2e,

$$\int \frac{d^d k}{(2\pi)^d} \frac{d^d l}{(2\pi)^d} \frac{1}{(k^2)^{-1} (l^2)^{-2} ((k+l)^2)^{-5} ((k+m v)^2 + m^2)^{-3}} \\ \frac{1}{((l+m v)^2 + m^2)^{-4}} = \frac{m^{2(d-1)}}{(4\pi)^d} M(1; 2; 3; 4; 5): \quad (18)$$

Using inversion, we can relate it to an HQET two{loop integral

$$M(1; 2; 3; 4; 5) = I(d-1, 3, 5; d-2, 4, 5; 3; 4; 5): \quad (19)$$

However, this HQET integral contains two non-integer indices, and is hence more difficult to calculate than before. The recurrence relation for M has a form identical to (5), allowing us to kill one of the lines 2, 4, or 5 of Fig. 2e. The integrals of Figs. 2f,h are trivial, as in (13), but that of Fig. 2g is not. Special combinations of recurrence relations allow us to kill one more line. As a result, all two{loop integrals of the type M can be expressed in terms of M_0^2 and $M_1 = (1+)^2(1-)(1+2^2)(1-4^2) = (1-2^2)(1-3^2)$ with rational coefficients.

We write the most difficult integrals, of the type N of Fig. 2d, as

$$\int \frac{d^d k}{(2\pi)^d} \frac{d^d l}{(2\pi)^d} \frac{1}{(k^2)^{-1} (l^2)^{-2} ((k+m v)^2 + m^2)^{-3} ((l+m v)^2 + m^2)^{-4}} \\ \frac{1}{((k+l+m v)^2 + m^2)^{-5}} = \frac{m^{2(d-1)}}{(4\pi)^d} N(1; 2; 3; 4; 5): \quad (20)$$

The recurrence relation

$$(d-2-2-4-5)N = [4^+2 + 5^+(2-3)]N \quad (21)$$

allows us to kill one of the lines 2 or 3 of Fig. 2d. This gives integrals shown in Fig. 2g, belonging to the class M , and integrals shown in Fig. 2j, which are still non-trivial. Special combinations of recurrence relations allow us to kill one more line. As a result, all two-loop integrals of the type N can be expressed in terms of M_0^2 , M_1 , and $N(0;0;1;1;1)$ with rational coefficients. Instead of using $N(0;0;1;1;1)$ as a basis integral, it is more convenient to use the convergent integral^{20,8}

$$N(1;1;1;1;1) = \frac{1}{2} \log^2 2 - \frac{3}{2} (3) + O(\epsilon^2); \quad (22)$$

whose $O(\epsilon^2)$ and $O(\epsilon^4)$ terms have been found by algebraic methods⁸, but are not necessary for two-loop calculations.

This algorithm is implemented in the REDUCE package RECURSOR⁸ and the FORM package SHELL2⁹. It was used for calculating heavy-quark on-shell mass⁶ and wavefunction⁷ renormalizations, and the QCD/HQET matching of heavy-light currents⁵. Coefficients of $(\epsilon_s)^2$ in the matching coefficients are typically of order 10, leading to substantial two-loop corrections in ratios such as $f_B = f_B$.

Acknowledgements

We are grateful to the Royal Society and PPARC, for grants which made our collaboration possible, to HUCAM, for travel funds, and to P. Gosdzinsky, for useful comments about the three-point case⁴.

References

1. M. Neubert, Phys. Reports 245 (1994) 260.
2. K. G. Chetyrkin, F. V. Tkachov, Nucl. Phys. B 192 (1981) 159.
3. D. J. Broadhurst, A. G. G. Rozin, Phys. Lett. B 267 (1991) 105.
4. E. Bagan, P. Ball, P. Gosdzinsky, Phys. Lett. B 301 (1993) 249.
5. D. J. Broadhurst, A. G. G. Rozin, preprint OUT-4102-52, hep-ph/9410240 (1994).
6. N. Gray, D. J. Broadhurst, W. Grafe, K. Schilcher, Z. Phys. C 48 (1990) 673.
7. D. J. Broadhurst, N. Gray, K. Schilcher, Z. Phys. C 52 (1991) 111.
8. D. J. Broadhurst, Z. Phys. C 54 (1992) 599.
9. J. Fleischer, O. V. Tarasov, CPC 71 (1992) 193; Phys. Lett. B 283 (1992) 129.
10. L. R. Surguladze, F. V. Tkachov, CPC 55 (1989) 205.
11. S. G. Gorishny, S. A. Larin, L. R. Surguladze, F. V. Tkachov, CPC 55 (1989) 381.
12. D. Knauss, K. Schamhorst, Annalen der Physik 41 (1984) 331.
13. X. Ji, M. J. Musolf, Phys. Lett. B 257 (1991) 409.
14. V. Gimenez, Nucl. Phys. B 375 (1992) 582; B 401 (1993) 116.
15. D. J. Broadhurst, A. G. G. Rozin, Phys. Lett. B 274 (1992) 421.
16. E. Bagan, P. Ball, V. M. Braun, H. G. Dosch, Phys. Lett. B 278 (1992) 457.
17. E. Bagan, P. Gosdzinsky, Phys. Lett. B 305 (1993) 157.
18. G. P. Korchemsky, A. V. Radyushkin, Nucl. Phys. B 283 (1987) 342.
19. M. Neubert, Phys. Rev. D 47 (1993) 4063.
20. D. J. Broadhurst, Z. Phys. C 47 (1990) 115.