

Sem i-Leptonic Decays of Λ_c and Λ_b Baryons involving Heavy to Light Transitions and the determination of $|V_{ub}|$

A lakabha D atta

Physics Department, University of Hawaii at Manoa, 2505 Correa Road, Honolulu, HI 96822, USA .

Abstract

We calculate the sem i-leptonic decays of Λ_c and Λ_b baryons which involve transitions of a heavy quark to a light quark . In a previous work we used data on the Cabibbo favoured non-leptonic decays of charmed baryons $\Lambda_c \rightarrow \Sigma^+ \pi^-$ and $\Lambda_c \rightarrow \Sigma^+ \pi^0$ to obtain information on the form factors in the $\Lambda_c \rightarrow \Sigma$ transition. This form factor information is used to study the sem i-leptonic decay $\Lambda_c \rightarrow \Sigma l \bar{\nu}_l$. Using SU (3) symmetry and HQET it is shown how a knowledge of form factors in $\Lambda_c \rightarrow \Sigma$ can be applied to $\Lambda_b \rightarrow p l \bar{\nu}_l$ decay and used to measure $|V_{ub}| = |V_{cs}| f$.

1 Introduction

Recently there has been a measurement of the form factors in the sem i-leptonic decay $\Lambda_c \rightarrow \Sigma^+ e^- \bar{\nu}_e$ [1], where a fit to the data was done using the model of Komer and Kramer [2]. The model, constructed within the framework of HQET, treats both the charm quark and the strange quark as heavy and considers $1/m_s$ expansion keeping $m_c \rightarrow 0$. Even though the

model has only one input parameter, use of $1=m_s$ expansion is questionable and the omission of $1=m_c$ corrections is also not justified. For a heavy to light transition, for example of the type $c \rightarrow l$, the use of HQET in the limit $m_Q \rightarrow \infty$ allows one to express all the form factors in terms of only two form factors [3]. Semileptonic decays of c have been studied in this limit [4, 2]. In Ref. [5, 6] we constructed a model for the form factors which takes into account $1=m_Q$ corrections for heavy-to-light transitions of the type baryon (light degrees in spin 0 state). In this work we use the form factors calculated in Ref. [6] to study the decays $c \rightarrow l$ and $b \rightarrow pl$.

The paper is organized in the following way. In the next section we present the formulae for the calculation of asymmetries and decay rates for the semileptonic decays $c \rightarrow l$ and $b \rightarrow pl$ while in the third section we present our results.

2 Formalism

Here we present the formalism for the semileptonic decays of $c \rightarrow l$ and $b \rightarrow pl$. The hadronic part of the amplitude is the matrix elements of the weak quark current between baryonic states that is parametrized in terms of form factors. We define the six vector and axial vector form factors through the following equations

$$\begin{aligned} \langle B^0(p^0; s^0) | j_Q^\mu | B_Q(p; s) \rangle &= u_{B^0}(p^0; s^0) \left[f_1 \frac{q^\mu}{m_{B_Q}} + \frac{f_2}{m_{B_Q}} q^\mu + \frac{f_3}{m_{B_Q}} q^\mu \right] u_{B_Q}(p; s) \\ \langle B^0(p^0; s^0) | j_Q^{\mu 5} | B_Q(p; s) \rangle &= u_{B^0}(p^0; s^0) \left[g_1 \frac{q^\mu}{m_{B_Q}} + \frac{g_2}{m_{B_Q}} q^\mu + \frac{g_3}{m_{B_Q}} q^\mu \right] u_{B_Q}(p; s) \end{aligned} \quad (1)$$

where $q = p - p^0$ is the four momentum transfer, B_Q is the baryon with a heavy quark and B^0 is the light baryon. We have shown in Ref. [5, 6] that we can write the form factors in the above equation in terms of two form factors F_1 and F_2 provided some assumptions are made about the $1=m_c$ corrections. For both c and b semileptonic decays we will work in the approximation that the lepton mass is negligible. The decay $c \rightarrow s l_1$ proceeds via $c \rightarrow s W_{\text{off shell}}$ followed by $W_{\text{off shell}} \rightarrow l_1$. Following [2] we define the helicity amplitudes which are given by

$$\begin{aligned} H_{2; W}^V &= H_{2; W}^{V^+} + H_{2; W}^{A^+} \\ H_{2; W}^{V(A)^+} &= \frac{1}{\sqrt{2}} (H_{2; W}^{V^+} \mp H_{2; W}^{A^+}) \end{aligned} \quad (2)$$

where λ_2 ; λ_W are the polarizations of the daughter baryon and the W^- -boson respectively. In terms of the form factors the helicity amplitudes are given by

$$\begin{aligned} H_{1=2;0}^V &= a_- (M_1 + M_2) f_1 + \frac{q^2}{m_{B_c}} f_2 V(q^2) \\ H_{1=2;0}^A &= a_+ (M_1 - M_2) g_1 + \frac{q^2}{m_{B_c}} g_2 A(q^2) \\ H_{1=2;1}^V &= \frac{q}{2Q} f_1 - \frac{(M_1 + M_2)}{m_{B_c}} f_2 V(q^2) \\ H_{1=2;1}^A &= \frac{q}{2Q} g_1 - \frac{(M_1 - M_2)}{m_{B_c}} g_2 A(q^2) \end{aligned} \quad (3)$$

where

$$\begin{aligned} Q &= (M_1 - M_2)^2 - q^2; \quad a_{\pm} = \frac{s_{\pm}}{q^2} \\ V(q^2) &= \frac{(1 - \frac{q_{max}^2}{m_{FF}^2})^2}{(1 - \frac{q^2}{m_{FF}^2})^2}; \quad A(q^2) = \frac{(1 - \frac{q_{max}^2}{m_{FF}^2})^2}{(1 - \frac{q^2}{m_{FF}^2})^2} \end{aligned} \quad (4)$$

where M_1, M_2 are the parent and daughter baryon masses and $m_{FF}^{V,A}$ are the appropriate pole masses. The decay $c \rightarrow s l_1$ is analysed by looking at the two sided cascade decay $c \rightarrow s [l_1 p] + W^- [l_1 l_2]$. The normalized four-fold joint angular distribution for the decay $1 \rightarrow 2 [l_1 a(\frac{1}{2}^+)] + b(0^-) + W^- [l_1 l_2]$ is given, following the notation in Ref. [2], as

$$\frac{d}{dq^2 d\cos\theta d\cos\theta_1 d\cos\theta_2} = B (c \rightarrow s [l_1 a] + b) \frac{1}{2} \frac{G^2}{(2)^4} \frac{q^2 p}{24M_1} |V_{Q_1 Q_2}|^2 [T_1 + T_2 + T_3] \quad (5)$$

where

$$\begin{aligned} T_1 &= \frac{3}{8} (1 - \cos\theta)^2 |H_{1=2;1}|^2 (1 + \cos\theta_1) + \frac{3}{8} (1 - \cos\theta)^2 |H_{1=2;-1}|^2 (1 - \cos\theta_1) \\ T_2 &= \frac{3}{4} \sin^2\theta |H_{1=2;0}|^2 (1 + \cos\theta_1) + |H_{1=2;0}|^2 (1 - \cos\theta_1) \\ T_3 &= \frac{3}{2} \cos\theta \sin\theta \sin\theta_1 (1 - \cos\theta) \text{Re}(H_{1=2;0} H_{1=2;1}) + (1 - \cos\theta) \text{Re}(H_{1=2;0} H_{1=2;-1}) \end{aligned}$$

where $p = \frac{p}{Q + Q} = 2M_1$, the upper and lower signs in the above equation hold for the l_1 and l_1^+ leptonic final states respectively and $V_{Q_1 Q_2}$ is the CKM element for the $Q_1 \rightarrow Q_2$ transition. The polar angles are defined in Ref. [2]. By integrating over two of the angles one can look at the following distributions

$$\frac{d}{dq^2 d\cos\theta} / (1 + \cos\theta)$$

$$\begin{aligned} \frac{d}{dq^2 d\cos\theta} &= \frac{1}{2} \cos^2\theta + \frac{1}{2} \cos\theta \\ \frac{d}{dq^2 d\phi} &= \frac{1}{32} \frac{3}{2} \cos\phi \end{aligned} \quad (6)$$

The first distribution in the above equation gives the polar angle distribution for the cascade decay $\chi_s \rightarrow p \gamma$. The second distribution is the polar angle distribution for the decay $W \rightarrow l_1 \nu$, while the third distribution is the azimuthal angle distribution. The asymmetries are given by the expressions below [2] and will depend only on the ratio $F_2^0(l=1)/F_1^0(l=1)$ in our model for the form factors

$$\begin{aligned} &= \frac{\mathcal{H}_{1=2,1}^J \mathcal{H}_{1=2,1}^J + \mathcal{H}_{1=2,0}^J \mathcal{H}_{1=2,0}^J}{\mathcal{H}_{1=2,1}^J + \mathcal{H}_{1=2,1}^J + \mathcal{H}_{1=2,0}^J + \mathcal{H}_{1=2,0}^J} \\ 0 &= \frac{\mathcal{H}_{1=2,1}^J \mathcal{H}_{1=2,1}^J}{\mathcal{H}_{1=2,1}^J + \mathcal{H}_{1=2,1}^J + 2(\mathcal{H}_{1=2,0}^J + \mathcal{H}_{1=2,0}^J)} \\ 0 &= \frac{\mathcal{H}_{1=2,1}^J + \mathcal{H}_{1=2,1}^J - 2(\mathcal{H}_{1=2,0}^J + \mathcal{H}_{1=2,0}^J)}{\mathcal{H}_{1=2,1}^J + \mathcal{H}_{1=2,1}^J + 2(\mathcal{H}_{1=2,0}^J + \mathcal{H}_{1=2,0}^J)} \\ &= \frac{2 \operatorname{Re}(\mathcal{H}_{1=2,0} \mathcal{H}_{1=2,1}^* + \mathcal{H}_{1=2,0} \mathcal{H}_{1=2,1}^*)}{\mathcal{H}_{1=2,1}^J + \mathcal{H}_{1=2,1}^J + \mathcal{H}_{1=2,0}^J + \mathcal{H}_{1=2,0}^J} \end{aligned} \quad (7)$$

for unpolarized χ_c . For polarized χ_c one has similar decay distributions and we refer the reader to Ref. [2] for the relevant details. We give here the expression for the asymmetries

$$\begin{aligned} P &= \frac{\mathcal{H}_{1=2,1}^J \mathcal{H}_{1=2,1}^J - \mathcal{H}_{1=2,0}^J \mathcal{H}_{1=2,0}^J}{\mathcal{H}_{1=2,1}^J + \mathcal{H}_{1=2,1}^J + \mathcal{H}_{1=2,0}^J + \mathcal{H}_{1=2,0}^J} \\ P &= \frac{2 \operatorname{Re}(\mathcal{H}_{1=2,0} \mathcal{H}_{1=2,0}^*)}{\mathcal{H}_{1=2,1}^J + \mathcal{H}_{1=2,1}^J + \mathcal{H}_{1=2,0}^J + \mathcal{H}_{1=2,0}^J} \end{aligned} \quad (8)$$

All the asymmetries have limiting values as $q^2 \rightarrow q_{\text{max}}^2$. All the polar asymmetries vanish at this limit while the azimuthal asymmetries $\rightarrow 2/3$ and $P \rightarrow 1/3$. At the $q^2 = 0$ point however the limiting value of the asymmetries A_0 , P and P depend on the dynamics of the semi-leptonic decay and we will comment on this issue further in the next section. The other asymmetries A_0 and A_0 tends to zero while A_0 tend to -1 in the $q^2 = 0$ point. These asymmetries were studied for different representative values of the ratio $r_{KK} = f_2/f_1$ [2] which is related to the ratio $r = F_2^0(l=1)/F_1^0(l=1)$ via $r_{KK} = r/(2+r)$. The fit, performed by Ref. [1], to the semi-leptonic decay data on $\chi_c \rightarrow s e^+ e^-$ uses the Komer Kramer (KK) model to extract r_{KK} and the asymmetry [1]. Since, as already mentioned in the introduction, the KK model does not include $1/m_c$ corrections and the use of $1/m_s$ expansion is questionable, a more correct approach would be therefore to do a fit to the

semi-leptonic data including $1=m_c$ corrections to extract the ratio $r(r_{KK})$ and hence the asymmetries. The formula for the calculation of the absolute decay rate for the semi-leptonic decay of baryons is given in Ref. [7]

3 Results

In our analysis we have studied the semi-leptonic decays for $j=c$ and $0 < F_1^0(!=1) < 1$. In Ref. [6] we found the $1=m_c$ expansion to be valid for $j=c$ and from the study of non-leptonic two body charmed baryon decays $F_1^0(!=1) < 1$ was obtained (we assumed $F_1^0(!=1) > 0$). In our analysis we found that the $1=m_q$ corrections depend on the value of r . As far as the $1=m_q$ corrections to the form factors are concerned for the c decay the $1=m_c$ corrections to the form factors are $< 50\%$ for $0.62 < r < 1$. For b decays the $1=m_b$ corrections are small and are less than 30% for $0.48 < r < 1$. Fig.1 shows $\langle q^2 \rangle$ versus q^2 for different r values. We note that for $0.5 < r < 1$ the asymmetry is not very sensitive to r . This feature is common to most of the calculated integrated asymmetries. In Fig.2 we show $\langle q^2 = 0 \rangle$ versus r . In the KK model or in general with $m_c \neq 1$, $\langle q^2 = 0 \rangle = 1$ for all r . Fig.2 shows the effect of including $1=m_c$ corrections. Fig.3 and Fig.4 show the predictions for the integrated asymmetries for different r for unpolarised and polarised c and except for c we see that the integrated asymmetries are insensitive to r in the range $0.5 < r < 1$. The $1=m_c$ corrections to the integrated asymmetries again depend on the value of r and can be as large as 40% .

The decay rates depend on both r and $F_1^0(!=1)$. So instead of calculating the individual decay rates for $c \rightarrow 1$ and $b \rightarrow pl$, to reduce the uncertainties from these sources we calculate the ratio of the decay rates which will depend only on r as the factor $F_1^0(!=1)$ cancels. Fig.5 shows this ratio as a function of r with and without $1=m_q$ corrections with a monopole and dipole form for the form factors. We see that the $1=m_q$ corrections are small for the dipole form factors. The ratio above will receive corrections of the order $1=m_q^2$ and higher where m_q is the c or b quark. An estimate of these corrections will depend on the value of r which can be extracted by performing a fit to the $c \rightarrow e^+ e^-$ data. The final result can be written as

$$\frac{\langle b \rightarrow pl \rangle}{\langle c \rightarrow 1 \rangle} = R_0(r) [1 + \alpha_1(r) + \alpha_2(r) + \dots] \frac{V_{ub}}{V_{cs}} \quad (9)$$

where R_0 is the ratio with $m_q \neq 1$ and $\alpha_j(r)$ represents the corrections to the ratio due to $1=m_q^j$ corrections to the form factors. Since we do not know r we can make an estimate of

the ratio above by using the value of r extracted from non-leptonic charm ed baryon decays with dipole form factors [6] i.e $r = r_0 = 0.47$ ($r_{KK} = 0.31$). We write

$$R_0(r) = R_0(r_0) + r R_0'(r_0) + \frac{r^2}{2} R_0''(r_0) + \dots \quad (10)$$

where $R_0'(r_0)$ is the derivative of $R_0(r)$ at $r = r_0 = 0.47$, and where r may also include experimental uncertainties in extracting r . So an estimate of the ratio defined in eqn.(9) can be made through the following approximation

$$\frac{(b \rightarrow c)_{\text{pl}}}{(b \rightarrow c)_{\text{sl}}} = R_0(r_0) [1 + r R_0'(r_0) + \frac{r^2}{2} R_0''(r_0) + \dots] \frac{V_{ub}}{V_{cs}} \quad (11)$$

For monopole and dipole form factors we then have

$$\frac{(b \rightarrow c)_{\text{pl}}}{(b \rightarrow c)_{\text{sl}}} = 196.65 [1 - 0.78 r + 0.02 r^2 + \frac{r^3}{6} R_0'''(r_0) + \dots] \frac{V_{ub}}{V_{cs}} \quad (12)$$

and

$$\frac{(b \rightarrow c)_{\text{pl}}}{(b \rightarrow c)_{\text{sl}}} = 83.41 [1 - 0.94 r + 0.15 r^2 + \frac{r^3}{6} R_0'''(r_0) + \dots] \frac{V_{ub}}{V_{cs}} \quad (13)$$

Naively we can set $r_2(r_0) = r_1(r_0) = 2m_Q$ and so we see that $r_2(r_0) > 0.05$ for the dipole form factors and $r_2(r_0)$ is negligible for the monopole form factors. Measurement of the ratio R can be used to extract V_{ub}/V_{cs} . Clearly the largest uncertainty in R comes from the assumed q^2 dependence of the form factors and to reliably extract V_{ub}/V_{cs} using R the q^2 dependence of the form factors must be measured. Our model for the form factors at $q^2 = 1$ can be tested by studying the decay distribution in $(b \rightarrow c)_{\text{pl}}$. With enough statistical power such a study could determine the q^2 dependence of the form factors. So if our model for the form factors at $q^2 = 1$ is tested successfully by doing a fit to the $(b \rightarrow c)_{\text{pl}}$ decay distribution, and if r and information on the q^2 dependence of form factors extracted, we can use the ratio in eqn.(9) to determine V_{ub}/V_{cs} with a small error.

It is interesting to compare this method of extracting V_{ub}/V_{cs} and hence V_{ub} with other methods of extracting V_{ub} . One method is to use semi-leptonic inclusive B decays and study the decay distribution near the end point of the spectrum in the narrow window $2.3 < E_1 < 2.6$ GeV [11]. However in this region the $b \rightarrow u$ transition is subject to large QCD radiative corrections as well as large non-perturbative corrections which are entangled with each other and theoretical calculations are model dependent. Another way to extract V_{ub} is through the study of exclusive B decays [12] and this approach is similar to our approach

with the difference that we are working with baryons instead of mesons. The calculations of form factors in exclusive decays of B mesons are done in specific models, like the quark model, or with chiral Lagrangian with heavy meson. In Ref. [13] it is suggested that $|V_{ub}|$ may be extracted from the ratio of the lepton distribution in $B \rightarrow l$ and $D \rightarrow l$ with an theoretical uncertainty of 10-20 %. The relevant form factors are calculated up to order $1/m_Q$ in the soft pion limit with the additional assumption that only intermediate states degenerate with the ground state meson contribute and so the calculations of the form factors are model dependent. In Ref. [14] the ratio of the lepton distribution in $B \rightarrow l$ and $D \rightarrow l$ at maximum q^2 is used for an extraction of $|V_{ub}|$. The $1/m_Q$ corrections are calculated in quark model and the theoretical uncertainty in the measurement of $|V_{ub}|$ is small. However at maximum q^2 the decay rates vanish, actual measurements are performed away from maximum q^2 where additional form factors contribute adding to the theoretical uncertainty in the measurement of $|V_{ub}|$. Another approach to measuring $|V_{ub}|$ which is the mesonic counterpart of our method, is given in Ref. [15] where experimental data on $D \rightarrow K$ semi-leptonic decay are used to calculate the $B \rightarrow$ semi-leptonic decay. The q^2 dependence of the form factors is assumed to be of the monopole type, no $1/m_Q$ corrections are considered and the extraction of $|V_{ub}|$ is found to depend on the value of f_{D_s} and f_B whose values are not known very accurately. Since most approaches to extracting $|V_{ub}|$ from B meson decays are model dependent and can involve significant theoretical uncertainties, extraction of $|V_{ub}|$ using heavy baryon decays is interesting as it is possible that theoretical uncertainties in heavy baryon decays might be small.

The predictions for the decay rates for $(\Lambda_c \rightarrow l)$ and $(\Lambda_b \rightarrow pl)$ are subject to significant $1/m_Q$ corrections and may not be reliable. Nevertheless we give the predictions for the decay rates with $r = 0.47$ and $F^0(l = 1) = 0.46$ [6]

$$(\Lambda_c \rightarrow l) = 5.36 \cdot 10^{10} s^{-1} \quad (14)$$

$$(\Lambda_b \rightarrow pl) = |V_{ub}|^2 6.48 \cdot 10^{12} s^{-1} \quad (15)$$

The Particle Data Group gives $(\Lambda_c \rightarrow e^+ X) = (7.0 \pm 2.5) \cdot 10^{10} s^{-1}$. Measurement of $(\Lambda_b \rightarrow pl)$ is likely to be made at LEP in the future. The numbers for the individual decay rates are calculated using $|V_{cs}| = 0.9745$ [9], the pole masses $m_V = 5.37$ GeV and $m_A = 5.80$ GeV [10] and the quark masses $m_b = 4.74$ GeV and $m_u = 0.005$ GeV [8]

In conclusion we have studied the semi-leptonic decays of charmed and bottom baryons involving transition of a heavy to light quark based on a model for form factors that includes $1/m_Q$ corrections. We also suggest a method for determining $|V_{ub}| = |V_{cs}|$.

Acknowledgement : I would like to thank Professor Sandip Pakvasa and Professor Tom Browder for useful discussions. This work was supported in part by US D.O.E grant # DE-FG 03-94ER40833.

References

- [1] G.Crawford et al, CLNS 94/1306, CLEO 94-34.
- [2] J.G.Komer and M.Kramer, Phys.Lett.B 275, 495 (1992).
- [3] T.Mannel, W.Roberts and Z.Ryzak, Nucl.Phys.B 355, 38 (1991); F.Hussain, J.G.Komer, M.Kramer and G.Thompson, Z.Phys.C 51, 321 (1991).
- [4] T.Mannel, W.Roberts and Z.Ryzak, Phys.Lett.B 255, 593 (1991).
- [5] A.lakabha Datta, UH-511-809-94, hep/ph 9411432 (To appear in Phys.Lett.B).
- [6] A.lakabha Datta, UH-511-824-95.
- [7] H. Pietschmann, Weak Interactions - Formulae, Results, and Derivations (Springer-Verlag Wien New York 1983).
- [8] Particle Data Group, Phys. Rev.D 50, 1435 (1994).
- [9] Particle Data Group, Phys. Rev.D 50, 1225 (1994).
- [10] S.Godfrey, N.Isgur, Phys. Rev.D 32, 189 (1985).
- [11] B.Blok, T.Mannel, Phys. Rev.D 51, 2208 (1995).
- [12] C.S.Kim et al, Phys. Rev.D 50, 5762 (1994).
- [13] G.Burdmann et al, Phys. Rev.D 49, 2331 (1994).
- [14] C.O.Dib, F.Vera, Phys. Rev.D 47, 3938 (1993).
- [15] D.Du, C.Liu, Phys. Rev.D 50, 4558 (1994).

3.1 Figure Captions

Fig. 1 The asymmetry $\langle q^2 \rangle$ versus q^2 for some representative values of the form factor ratio r .

Fig. 2 The asymmetry $\langle q^2 = 0 \rangle$ versus r . The $m_c \rightarrow 1$ prediction for $\langle q^2 = 0 \rangle$ is 1 for all r . This figure therefore shows the effect of $1=m_c$ corrections.

Fig. 3 The average asymmetries as functions of r when the initial heavy baryon is unpolarised.

Fig. 4 The average asymmetries as functions of r when the initial heavy baryon is polarised.

Fig. 5 The ratio of the decay rates of $(\Lambda_b \rightarrow p l) = (\Lambda_c \rightarrow l)$ versus r . R_0 and R represents the ratio without and with $1=m_c$ corrections. Monopole and dipole form factors are used for the q^2 dependence of the form factors.

This figure "fig1-1.png" is available in "png" format from:

<http://arxiv.org/ps/hep-ph/9504429v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arxiv.org/ps/hep-ph/9504429v1>

This figure "fig1-3.png" is available in "png" format from:

<http://arxiv.org/ps/hep-ph/9504429v1>

This figure "fig1-4.png" is available in "png" format from:

<http://arxiv.org/ps/hep-ph/9504429v1>

This figure "fig1-5.png" is available in "png" format from:

<http://arxiv.org/ps/hep-ph/9504429v1>