



University of Wisconsin – Madison

MADPH-95-885

May 1995

# Flux Tubes in Effective Field Theory

M. G. Olsson and Sinisa Veseli

Department of Physics, University of Wisconsin, Madison, WI 53706

Ken Williams

Continuous Electron Beam Accelerator Facility

Newport News, VA 23606, USA

and

Physics Department, Hampton University, Hampton, VA 23668

## Abstract

Quark-antiquark bound states are examined in the long-range strong-coupling limit with the minimal area law of lattice gauge theory assumed as input. Matrix element relations are established which in the effective theory obtain dynamical equations equivalent to a formulation of the flux-tube model.

# 1 Introduction

The success of QCD as the correct theory of strong interactions has rested primarily upon perturbative calculations. The prediction of hadronic properties is inherently non-perturbative and has been much slower to develop. Most of lattice progress has turned out to be largely irrelevant to hadron dynamics. An exception has been the low quark velocity Wilson loop expansion [1]. The aim of this paper is to extend this method to arbitrary quark velocity. Our principal result is that under natural assumptions QCD provides a framework for meson dynamics that is identical with the relativistic flux tube (RFT) model [2, 3].

The RFT model assumes that a chromoelectric flux tube stretches in a straight line between the quark and the antiquark in a meson. The tube rest frame energy per unit length also contributes to rotational energy, momentum, and angular momentum in the meson rest frame. Also, as pointed out by Buchmüller [4], a pure chromoelectric field in the tube rest frame implies no long range spin-spin correlation. In the last few years the RFT model has been demonstrated to be numerically calculable for any quark mass case [5]. The RFT model provides intuitive physical pictures for relativistic corrections in agreement with QCD and becomes a Nambu-Goto string for high rotational states [2].

The QCD Lagrangian density is a fundamental object in the theory of strongly interacting particles. For a meson it is given by the minimally transformed free Lagrangian

$$p = p + gA; \quad (1)$$

$$L_{free} = L_{QCD} = \sum_j q_j (\not{D} - m_j) q_j - \frac{1}{4} F^a F_a; \quad (2)$$

from which follow the Euler-Lagrange equations

$$(\not{D} - m_j) q_j = 0; \quad (3)$$

$$D F = g \sum_j q_j T q_j; \quad (4)$$

These are non-linear and amenable only as a perturbation expansion in small coupling  $g$ . Wilson's observation [6] over two decades ago has led to some of the most fruitful work in the non-perturbative regime. The simple picture offered in his area law for static quarks is compelling: Contributions to the  $q\bar{q}$  propagator fall off exponentially in the area swept out by lines of gauge invariant chromoelectric flux joining the constituents' world lines; in this way, widely separated paths are suppressed. Subsequent lattice simulations verify the same qualitative behavior for dynamical quarks.

Progress in obtaining a consistent description of confinement for realistic  $0+$ -lattice calculations has been largely uncertain. The variety of semi-relativised confinement models based on the above simple idea is striking and is partly due to the limited theoretical input available beyond the static limit. Notable exceptions are in works of Eichten and Feinberg, Gromes and later those of Brambilla and Prosperetti [1] where analytic expressions are derived from the minimal area law to first order in the inverse heavy quark mass. More recently, a relativistic QCD-string Lagrangian for spinless quarks [7] has been deduced.

Here we derive an effective Hamiltonian for a quark-antiquark system in the confinement region with the minimal area (MA) law of lattice gauge theory as input. In this arrangement, the gauge field's non-Abelian character is manifest through the minimal area law; complete non-Abelian expressions are therefore given only where useful, e.g., to clarify a point. Our method consists in expressing the usual canonical matrix elements from (2) in terms of Wilson loop expectation values derived from the MA law. In the effective theory these become first-quantized operators acting on state vectors. The results are fully relativistic.

## 2 Wilson loop expectation values

We begin with the Lagrangian density assuming that the MA analysis has been carried out; hence fermion and gauge fields are interrelated at the outset, effectively reducing the degrees of freedom in (2). The idea is simple: gauge fields specified by MA relations are taken to be "external" to the Lagrangian; Euler-Lagrange nonlinearities (3-4) are thereby hidden within undetermined matrix elements.

From the energy-momentum tensor of (2) the conserved quantities are the usual

$$H = \int d^3x q_j^y \left[ \frac{1}{2} (\vec{r} \cdot \vec{g}A) + gA_0 + m_j \right] q_j^y ; \quad (5)$$

$$P = \int d^3x q_j^y (\vec{r}) q_j^y ; \quad (6)$$

$$J = \int d^3x q_j^y \left[ \vec{r} \cdot (\vec{r}) + \frac{1}{2} \right] q_j^y ; \quad (7)$$

which with the MA law,

$$\langle \text{Tr} W(C) \rangle = a S_{\text{min}} ; \quad (8)$$

define our bound state problem. Eventually, we take independent variations of (8) along the paths C.

We consider the Wilson loop of a straight slice out of the qqm in an all world sheet from time  $t_0$ ,

$$\begin{aligned} \langle \text{Tr} W(C) \rangle &= \frac{1}{3} \text{Tr} P \exp \left( i g \oint_C dx^\mu A_\mu(x) \right) \\ &= \frac{1}{3} \text{Tr} P \exp \left( i g \int_{t_0}^{t_f} dt \left[ A_0(z_1) - A_0(z_2) \right] U(z_2^0; z_1^0) \right. \\ &\quad \left. P \exp \left( i g \int_{z_1^0}^{z_2^0} dz^\mu A_\mu(z) \right) U(z_{1f}; z_{2f}) \right) ; \end{aligned}$$

The U's here are straight-line path ordered exponentials, and the average is taken over gauge fields,

$$\langle \# \rangle = \frac{\int D A \exp \left( i S[A] \right) \# [A]}{\int D A \exp \left( i S[A] \right)} ; \quad (9)$$

where the pure gauge action  $S[A]$  includes a gauge fixing term.

The c-number path variables for a given gauge field  $(z_i$  and  $\bar{z}_i)$  are related to coordinate and mechanical momentum matrix elements,

$$z_i = \langle \psi_i | z | \psi_i \rangle = \langle \psi_i | \hat{z} | \psi_i \rangle; \quad (10)$$

$$m_{ij} = \langle \psi_i | \hat{p} | \psi_j \rangle = \langle \psi_i | \hat{p} | \psi_j \rangle; \quad (11)$$

$$\langle \psi_i | \hat{p} | \psi_j \rangle = \frac{1}{1 + z_i^2}; \quad (12)$$

where  $\psi_i$  is either a particle or antiparticle Euler-Lagrange solution (3-4) with a classical gauge field. These valence or quenched solutions prevent the possibility of time-backtracking [7] not present in the M A picture. If  $\psi$  is the meson wavefunction in a product space of  $\psi_i$  and  $\bar{\psi}_i$ , a reasonable ansatz is the auxiliary condition

$$\psi = \sum_i \psi_i^{(1)} \psi_i^{(2)}; \quad (13)$$

where  $\psi_i^{(i)}$  are positive or negative energy projection operators. In this approximation, the Casimir operators reappear in the Hamiltonian (5).

On the contrary, we point out that the quark degrees of freedom have not yet entered the dynamics except as a means to specify the physical gauge field configurations. The above identifications are made for later use.

The enclosed area we parameterize in the Nambu-Goto form,

$$S = \int_0^1 \int_0^1 d\tau d\sigma \left[ \frac{1}{2} \dot{x}^2 + \frac{1}{2} x'^2 \right] \quad (14)$$

with  $x = x(\tau, \sigma)$ ;  $x_0 = \frac{e x}{e}$ , and  $x^0 = \frac{e x}{e}$ . At the boundary,  $x(\tau, 1) = h_1(\tau)$  and  $x(\tau, 0) = h_2(\tau)$ . In the usual straight-line and equal times approximations, which we assume here, the minimum is given by

$$x_{min}(\tau) = \frac{1}{2} (h_1(\tau) + h_2(\tau)); \quad (15)$$

A general path variation of (8) for fixed endpoints and fixed time,

$$\begin{aligned} \frac{\langle \ln W(C) \rangle}{\langle W(C) \rangle} &= \frac{\{ \int_0^Z \frac{1}{3} \text{TrP exp} (\{ g_c dx A(x) \})}{\langle W(C) \rangle} \\ &\quad g \left( z_1 \frac{\partial}{\partial z_1} + z_1 \frac{\partial}{\partial z_1} \right) A_0(z_1) - z_1 A(z) \\ &\quad \left( z_2 \frac{\partial}{\partial z_2} + z_2 \frac{\partial}{\partial z_2} \right) A_0(z_2) - z_2 A(z) \\ &= \int_0^Z \frac{1}{3} \frac{\partial S}{\partial x^0} dx^0 + \frac{\partial S}{\partial x} x_{\text{min}} ; \end{aligned} \quad (16)$$

specified to  $z_1 = z_2$ , obtains in the center of momentum (see Appendix for details)

$$\begin{aligned} \langle \ln gA(z) \rangle &= \frac{1}{\langle W(C) \rangle} \frac{1}{3} \text{TrP exp} (\{ g_c dx A(x) \}) gA(z) \\ &= \int_0^Z dx \cdot x \cdot p_t ; \end{aligned} \quad (17)$$

which are the desired Wilson loop expectation values with reference points chosen at the origin,  $A(0) = 0$ , and "?" defined relative to the straight line,  $v \cdot v$  ( $v \cdot \hat{t}$ ) $\hat{t}$ . It should be clear that the gauge fields quantized in this average carry spatial dependence in the radial coordinate only. Evidently, this procedure selects the physically realizable transverse polarizations. The time component is obtained by simple differentiation of (8) with respect to  $\tau$ ,

$$\begin{aligned} \frac{d}{d\tau} \{ \ln W(C) \} &= \frac{\{ \frac{1}{3} \text{TrP exp} (\{ g_c dx A(x) \})}{\langle W(C) \rangle} \\ &\quad g f A_0(z_1) - A_0(z_2) - [z_1 A(z) - z_2 A(z)] g \\ &= \int_0^Z \frac{1}{3} \frac{\partial S}{\partial x_{\text{min}}} dx ; \end{aligned} \quad (18)$$

yielding (using the procedure described in the Appendix)

$$\langle \ln gA_0(z) \rangle = \int_0^Z dx \cdot x \cdot H_t ; \quad (19)$$

and also

$$\langle \ln z \cdot gA(z) \rangle = \int_0^Z dx (x \cdot x_{\text{min}}) \cdot L_t ; \quad (20)$$

### 3 Relation to Flux Tube Model

With the Wilson loop expectation values we re-express conserved quantities (5-7) in terms of coordinate and velocity matrix elements. In the effective theory these are promoted to noncommuting quantum operators, observables requiring symmetrization. We give the naive expressions

$$H = \sum_{i=1}^N \frac{1}{2} m_i \dot{x}_i^2 + \sum_{i=1}^N \frac{1}{2} g^2 A_0^2(x_i) \quad (21)$$

$$P = \sum_{i=1}^N m_i \dot{x}_i = 0 \quad (22)$$

$$J = \sum_{i=1}^N \left[ x_i \times m_i \dot{x}_i + \frac{1}{2} g^2 x_i \times \nabla A_0^2(x_i) \right] \quad (23)$$

This compares well with other results from the Wilson loop. Both spin and spin-independent Hamiltonians of [1], for example, are reproduced on semi-relativistic reduction of (22). Equations (21-23) in fact define the relativistic flux tube model [8] as formulated by the present authors in [2, 3]. There, dynamical relations are derived from a heuristic Lagrangian in which the flux tube is fashioned as a simple constant energy density. The "tube operators" are equivalent to the Wilson loop expectation values above. This straightforward derivation serves to clarify the model's close relation to the underlying QCD.

### 4 Conclusion

We have demonstrated that under the same assumptions as in previous work [1, 7] that the Wilson loop area law and the QCD Lagrangian yields a relativistically valid picture of dynamical confinement. We emphasize that we have not "solved" QCD, but that our ignorance can be distilled into three loop expectation values each of which is equal to that of the mechanical RFT model. We end that

$$\langle \text{Area} \rangle = H_t \quad (24)$$

$$\langle H_{\text{gA}} \rangle = p_t ; \quad (25)$$

$$\langle H_{\text{r-gA}} \rangle = L_t ; \quad (26)$$

for the tube energy, momentum, and angular momentum respectively. In addition we have shown that the natural equation of motion is the Salpeter equation in which the "covariant tube substitution" [3] has been made.

The verification of the RFT model structure promises more than to legitimize a physically reasonable model. The close relation of the RFT model to lattice QCD should shed light on both subjects. In addition, a systematic program of improving the RFT model to include field fluctuations [8] can now be envisioned.

## A Appendix

Equation (17) follows from the physical gauge field's spatial dependence in the straight line and equal times approximations

$$A(r) = \sum_k \exp(-ik \cdot r) \hat{A}(k) \exp(ik \cdot r) ; \quad (27)$$

Then small angular variations of a given path at fixed time leave  $A$  unchanged. It will suffice to consider the transverse part of (16) with

$$z_i(\tau; A) = z_i(\tau; A^0) + \delta z_i ; \quad (28)$$

for all gauge fields at a given  $\tau$ . Naturally,

$$\langle \delta z_i \rangle = 0 ; \quad (29)$$

We write the Wilson loop in discrete form,

$$W(C) = \frac{1}{3} \text{Tr} P \prod_{n=0}^N \exp(\{g t_n [A_0(z_{1n}) - z_{1n} \cdot A(z_n)] U(z_2^0; z_1^0) \exp(-g t_n [A_0(z_{2n}) - z_{2n} \cdot A(z_n)] U(z_{1f}; z_{2f}) ; \quad (30)$$



and take the path variations of (8) according to

$$\begin{aligned} \langle \{ \ln W(C) \} \rangle &= \frac{\{ \ln W(C) \}}{\langle W(C) \rangle} \langle W(C) \rangle \\ &= \frac{\{ \ln W(C) \}}{\langle W(C) \rangle} \left( \sum_{n=0}^N \left( z_{1n} \frac{\partial}{\partial z_{1n}} + z_{2n} \frac{\partial}{\partial z_{2n}} \right) W(C) \right) ; \end{aligned} \quad (31)$$

giving equation (16),

$$\begin{aligned} \frac{1}{\langle W(C) \rangle} \langle \ln W(C) \rangle &= \frac{1}{\langle W(C) \rangle} \int_0^1 dx \exp \left( \int_0^x A(x) dx \right) \left( z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} \right) f(z_1; z_2) \\ &= \frac{1}{\langle W(C) \rangle} \int_0^1 dx \frac{\partial S}{\partial x} \frac{\partial}{\partial x} \left( z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} \right) f(z_1; z_2) ; \end{aligned} \quad (32)$$

where

$$f(z_1; z_2) = \langle (1 - \langle z_1 \rangle) g_A(z_1) - g_A(0(z_1)) \rangle ; \quad (33)$$

The temporal derivative is transferred from the variations by partial integration, so that

$$\begin{aligned} \frac{1}{\langle W(C) \rangle} \langle \ln W(C) \rangle &= \frac{1}{\langle W(C) \rangle} \int_0^1 dx \exp \left( \int_0^x A(x) dx \right) \left( \frac{d}{dx} \frac{\partial}{\partial z_1} + \frac{\partial}{\partial z_1} \right) f(z_1; z_2) \\ &= \frac{1}{\langle W(C) \rangle} \int_0^1 dx \frac{\partial S}{\partial x} \frac{\partial}{\partial x} \left( z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} \right) f(z_1; z_2) ; \end{aligned} \quad (34)$$

yielding in the  $z_1 = z_2 (1 - z_2)$  case of interest

$$\langle \ln \frac{\partial}{\partial z_1} f(z_1; z_2) \rangle = \frac{1}{\langle W(C) \rangle} \int_0^1 dx \frac{\partial S}{\partial x} \frac{\partial}{\partial x} \left( z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} \right) f(z_1; z_2) ; \quad (35)$$

or

$$\langle \ln g_A(z_1) - g_A(z_2) \rangle = \frac{1}{\langle W(C) \rangle} \int_0^1 dx \frac{h_{z_1?} + (1 - h_{z_2?})}{[1 - (h_{z_1?} + (1 - h_{z_2?}))^2]^{1/2}} ; \quad (36)$$

This obtains (17) when reference points are chosen at the origin of coordinates.

Also, with the  $A_0$  reference point chosen at the origin,  $A_0(0) = 0$ , and (18) can be written

$$\frac{1}{h\omega} \frac{1}{(C)i3} \frac{\text{Tr P} \int_0^R dA_t g[A_t(z_{A_t})] \int_0^R dA_{\epsilon t} \exp(i g_c \int_0^R dx A_t) \exp(iS[A])}{D A \exp(iS[A])} \\ = ahz_t i_0^{R_1} d [1 - 2 hz_t i_2^2]^{1=2} ; \quad (37)$$

with

$$hz_t i = \frac{\int_0^R dA_t z_{A_t} \int_0^R dA_{\epsilon t} \exp(iS[A])}{D A \exp(iS[A])} : \quad (38)$$

Then, the functional derivative of (37) with respect to  $z_{A_t}^0$  is

$$\frac{1}{h\omega} \frac{1}{(C)i3} \frac{\text{Tr P} \int_0^R (g A_t^0) \int_0^R dA_{\epsilon t} \exp(i g_c \int_0^R dx A_t) \exp(iS[A])}{D A \exp(iS[A])} \\ = ahz_t i_0^{R_1} d (1 - 2 hz_t i_2^2) [1 - 2 hz_t i_2^2]^{1=2} \frac{\int_0^R dA_{\epsilon t} \exp(iS[A])}{D A \exp(iS[A])} : \quad (39)$$

Taking the scalar product of the above with  $z_{A_t}^0$ , and integrating over  $A_t^0$ , gives

$$hz_t g A_{ii} = a \int_0^Z hz_t i dx x_2^2 ; \quad (40)$$

which is used in obtaining (19). Taking the vector product of (39) with  $z_{A_t}^0$  yields (20) after integration over  $A_t^0$ ,

$$hz_t g A(z)_{ii} = a \int_0^Z hz_t i dx (x_1 \otimes x_2) : \quad (41)$$

#### ACKNOWLEDGMENTS

We would like to thank N. Brambilla and G. Proserpi for reading the manuscript and making many helpful comments. This work was supported in part by the U.S. Department of Energy under Contract Nos. DE-FG 02-95ER 40896 and DE-AC 05-84ER 40150, the National Science Foundation under Grant No. HRD 9154080, and in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation.

## References

- [1] E. Eichten and F. Feinberg, *Phys. Rev. D* 23, 2724 (1981); D. Gromes, *Z. Phys. C* 22, 265 (1984); N. Brambilla, P. Consoli, and G. M. Prosperi, *Phys. Rev. D* 50, 5878 (1994); N. Brambilla and G. M. Prosperi, Wilson loops,  $q\bar{q}$  and  $3q$  potentials, Bethe-Salpeter equation, Proceedings of the Conference "Quark Confinement and the Hadron Spectrum", World Scientific Singapore, 1995.
- [2] D. LaCourse and M. G. Olsson, *Phys. Rev. D* 39, 2751 (1989); C. Olson, M. G. Olsson, and K. Williams, *Phys. Rev. D* 45, 4307 (1992).
- [3] M. G. Olsson and Ken Williams, *Phys. Rev. D* 48, 417 (1993); M. G. Olsson, Sinisa Veseli, and Ken Williams, Fermion Confinement by a Relativistic Flux Tube, UW-Madison preprint MADPH-95-868 (hep-ph/9501323).
- [4] W. Buchmüller, *Phys. Lett. B* 112, 479 (1982).
- [5] M. G. Olsson and Sinisa Veseli, *Phys. Rev. D* 51, 3578 (1995).
- [6] K. Wilson, *Phys. Rev. D* 10, 2445 (1974).
- [7] A. Yu. Dubin, A. B. Kaidalov and Yu. A. Simonov, *Phys. Lett. B* 323, 41 (1994).
- [8] N. Isgur and J. Paton, *Phys. Lett. B* 124, 247 (1983).