

hep-ph/9505219
RAL-95-036
3 May 1995

Evidence for a Scalar Glueball

Claude Amisler

Physik-Institut, Universitat Zurich, CH-8057 Zurich, Switzerland

Frank E. Close^z

Particle Theory, Rutherford-Appleton Laboratory, Chilton, Didcot OX11 0QX, UK

May 1995

Abstract

We show that the newly discovered scalar meson $f_0(1500)$ at LEAR has properties compatible with the lightest scalar glueball predicted by lattice QCD and incompatible with a $Q\bar{Q}$ state. We suggest that decays of glueballs are into pairs of glueballs (including $\pi^0\pi^0$ or $(\pi\pi)_S$) or by mixing with nearby $Q\bar{Q}$ states. The partial widths of $f_0(1500)$ are in accord with this hypothesis, tests of which include characteristic radiative decays to $\pi^0\gamma$; $\eta\gamma$; and the prediction of a further scalar state, $f_0'(1500-1800)$ which couples strongly to $K\bar{K}$, $\eta\eta$ and $\pi^0\pi^0$.

Submitted to Physics Letters

E-mail : amisler@cernvm.cern.ch

^yE-mail : fec@v2.rl.ac.uk

^zWork supported in part by the European Community Human Capital Mobility program "Eurodafne", Contract CHRX-CT 92-0026

Glueballs are a major missing component of the standard model. For two decades, experimental searches for bound states of gluons have produced only controversial signals. Whereas the gluonic degrees of freedom expressed in L_{QCD} have been established beyond doubt in high momentum data, their dynamics in the strongly interacting limit epitomised by hadron spectroscopy remain obscure.

In this letter we show that if intuition based on lattice QCD is a reliable guide, the emerging properties of the scalar meson $f_0(1500)$ discovered at LEAR are consistent with those of a glueball mixed with nearby members of the quark model scalar nonet. This hypothesis may be tested in forthcoming experiments.

There are now clear signals for the $f_0(1500)$ in a range of production processes that are traditionally believed to favour glueballs, namely [1]

1. Radiative J/ψ decay: $J/\psi \rightarrow \gamma + G$ [2]
2. Collisions in the central region away from quark beams and target: $pp \rightarrow p_f(G)p_s$ [3, 4].
3. Proton-antiproton annihilation where the destruction of quarks creates opportunity for gluons to be manifested [5]–[10]. The $f_0(1500)$ was observed to decay into $\pi^0\pi^0$, and π^0 in $\bar{p}p \rightarrow 3\pi^0$ [5, 6], $\pi^0\pi^0$ [7, 8] and $\pi^0\pi^0\pi^0$ [9].

By contrast, there are no significant sightings of $f_0(1500)$ in processes where glueballs are not expected to be enhanced.

This *prima facie* evidence for a scalar glueball is particularly interesting now that studies of lattice-QCD appear to be coming to a consensus that the lightest glueball is indeed a scalar in this region of mass [11, 12]. In these circumstances it is natural to speculate that the $f_0(1500)$ and the "primitive" (i.e. quenched approximation) scalar glueball of lattice-QCD are intimately related. The purpose of this letter is to evaluate and to propose further tests of this hypothesis. A detailed discussion may be found in ref. [13].

Lattice QCD is not yet able to make detailed quantitative statements about the partial decay widths of glueballs against which we could evaluate those of $f_0(1500)$. However, qualitative features may be abstracted from flux-tube models which are probably the nearest we can presently get to simulating the strong coupling lattice formulation of QCD. In this simulation [14] mesons consist of a quark and antiquark connected by a tube of coloured flux whereas glueballs G_0 consist of a loop of flux. These eigenstates are perturbed by two types of interaction [15]

1. V_1 which creates a Q and a \bar{Q} at neighbouring lattice sites, together with an elementary flux-tube connecting them. When V_1 acts on conventional $Q\bar{Q}$ mesons it causes their decays (g. 1a), and is thereby known to be significant. Fig. 1b shows that when V_1 acts on a glueball G_0 , in leading order it causes glueball- $Q\bar{Q}$ mixing.
2. V_2 which creates or destroys a unit of flux around any plaquette, where a plaquette is an elementary square with links on its edges: in leading order this causes glueballs to decay to pairs of glueballs (g. 1c).

Note that in this approach G_0 does not decay to $\pi\pi$ or $K\bar{K}$ in leading order since gluons are isoscalar; decays to $\pi\pi$ and η are allowed to the extent that these have nonzero overlap with glue.

The mixing with $Q\bar{Q}$ is likely to play a significant role in glueball phenomenology if a quarkonium nonet of the same J^{PC} is nearby. As a result of extensive data from the Crystal Barrel collaboration at LEAR it is now clear that there is a scalar nonet in the 1.5 GeV region, as well as the $f_0(1500)$.

(i) The discovery of the $I = 1$ state $a_0(1450)$ [16] sets the natural scale of masses and widths for the other members of the scalar nonet. In quark models such as ref. [15, 17] the widths of the scalar $Q\bar{Q}$ are qualitatively ordered as $(\bar{n}n) > (s\bar{s}) > (a_0) > (K_0)$. Empirically $(a_0) = 270 \pm 40$ MeV, $(K_0) = 287 \pm 23$ MeV which are consistent with quark model expectations and lead one to expect for their partners $(\bar{n}n) \sim 400\text{--}700$ MeV and $(s\bar{s}) \sim 300\text{--}500$ MeV [13, 15, 17, 18].

(ii) The Crystal Barrel data show clear signals for an independent scalar meson $f_0(1370)$ in $\pi\pi$ and $\eta\eta$ whose mass is consistent with it being the nn partner of the $a_0(1450)$ and whose ratio of partial widths to $\pi\pi$ and $\eta\eta$ also is consistent with it being the nn $(u\bar{u} + d\bar{d}) = \frac{2}{3}$ state of a nonet [5, 6, 8, 16]. The total width of $f_0(1370)$ is not yet well determined, 200–700 MeV being possible [5, 8, 6] depending on the theoretical model used in the analysis and in accord with the nn hypothesis. The width in this region is also consistent with it containing the nn state [19, 20]. By contrast, the $f_0(1500)$ width is 116 ± 17 MeV [6, 9, 8] and is clearly out of line with the scalar nonet, being even smaller than the K_0 and a_0 widths.

The properties of the $f_0(1500)$ – $f_0(1370)$ system are incompatible with them both belonging to a $Q\bar{Q}$ nonet. The $f_0(1370)$ appears to be dominantly nn and on mass grounds we expect that the ss would lie some 200–300 MeV higher than this and the $a_0(1450)$. The strong coupling of $f_0(1500)$ to pions implies that it is not primarily an ss state; the decoupling from $K\bar{K}$ further distances it from the nonet. Specifically [13] after correcting for phase space and minor effects of form factors, the Crystal Barrel data imply for the ratios of partial widths

$$R_1 = \frac{\Gamma(f_0(1500) \rightarrow \pi\pi)}{\Gamma(f_0(1500) \rightarrow \eta\eta)} = 0.27 \pm 0.11; \quad (1)$$

$$R_2 = \frac{\Gamma(f_0(1500) \rightarrow \eta\eta)}{\Gamma(f_0(1500) \rightarrow \eta\eta')} = 0.19 \pm 0.08; \quad (2)$$

while a bubble chamber experiment leads to the (95% C.L.) upper limit [21]

$$R_3 = \frac{\Gamma(f_0(1500) \rightarrow K\bar{K})}{\Gamma(f_0(1500) \rightarrow \pi\pi)} < 0.1; \quad (3)$$

In the $Q\bar{Q}$ hypothesis it is possible to fit two ratios but not all three for the $f_0(1500)$. This is shown in fig. 2 [22]: The roughly equal couplings to $\pi\pi$ and $\eta\eta$ together with the dominance of $\pi\pi$ (eqn. 1 and 2) imply that $f_0(1500)$, if $Q\bar{Q}$, is nearly $\bar{n}n$. However, for an $\bar{n}n$ state, $\Gamma(K\bar{K}) = 1/3 \Gamma(\pi\pi)$, in contradiction with data (eqn. 3). Furthermore, the

strong affinity of $f_0(1370)$ for η and its branching ratios and widths suggest that this state is strongly nn which creates further problems for a $Q\bar{Q}$ interpretation of $f_0(1500)$. This remains true for any reasonable breaking of $SU(3)_f$ symmetry [13]. These results are stable against form factor choice [13, 18] and contrast the experience with other known nonets [23].

However, the observed decay branching ratios and in particular the suppression of $K\bar{K}$ for the $f_0(1500)$ are natural for a scalar glueball which is in the vicinity of the $Q\bar{Q}$ nonet and mixes with the two nearby $Q\bar{Q}$ isoscalars, one (nn) with mass below 1.5 GeV and the other (ss) above. The $f_0(1370)$ is a natural candidate for the nn ; this hypothesis requires that a (mainly) $s\bar{s}$ state lies in the 1600 MeV region.

We demonstrate this more quantitatively by considering the effect of V_1 on a primitive glueball G_0 . For this first look we assume flavour blindness at the fundamental level such that $\langle \bar{s}s | V_1 | G_0 \rangle = \langle \bar{d}d | V_1 | G_0 \rangle$. The quarkonium mixing into the glueball state is then

$$N_G |G\rangle = |G_0\rangle + \frac{P_-}{f^2(2 + \lambda^2)} \lambda \text{sig} |G_0\rangle + \frac{P_-}{2} |\bar{Q}Q\rangle \quad (4)$$

where N_G is the normalisation $\frac{1}{1 + \frac{1}{2}(2 + \lambda^2)}$, λ is the dimensionless mixing parameter

$$\lambda = \frac{\langle \bar{d}d | V_1 | G_0 \rangle}{E_{G_0} - E_{nn}}; \quad (5)$$

and

$$\lambda = \frac{E_{G_0} - E_{nn}}{E_{G_0} - E_{ss}} \quad (6)$$

is the ratio of the mass gap of G_0 and the nn and ss intermediate states in old fashioned perturbation theory (eq. 3).

Only in the particular case $\lambda = 1$, where $E_{nn} = E_{ss}$, does the underlying flavour blindness naturally survive at hadron level as G_0 mixes into the flavour singlet

$$|\bar{Q}Q\rangle = \frac{1}{\sqrt{3}} (|\bar{u}u\rangle + |\bar{d}d\rangle + |\bar{s}s\rangle) \quad (7)$$

However, when $\lambda \neq 1$, as will tend to be the case when G_0 is in the vicinity of a $Q\bar{Q}$ nonet with the same J^{PC} quantum numbers (as in the 0^{++} case of interest here) significant distortion from naive flavour singlet can arise: the mass breaking, $m_s - m_d$, which is usually regarded as a small perturbation in hadron dynamics, is here magnified.

There are three data that suggest a self consistency with this hypothesis.

1. The suppression of $K\bar{K}$ in the $f_0(1500)$ decays suggests a destructive interference between nn and ss such that $\lambda \approx 1$. This arises naturally if the primitive glueball mass is equidistant between those of nn and the primitive ss - a situation not inconsistent with lattice QCD. As the mass of $G_0 \rightarrow m_{nn}$ or m_{ss} , the $K\bar{K}$ remains suppressed though non-zero. Specially from eqn. 4 and $SU(3)_f$ we obtain for $G = f_0(1500)$

$$\langle K\bar{K} | V_1 | G \rangle = \frac{1 + \lambda}{2} \langle V_1 | G \rangle \quad (8)$$

per unit charge combination for $K \bar{K}$ and $\pi\pi$. The upper limit R_3 (eqn. 3) leads to the constraint

$$0.27 < \frac{\langle K \bar{K} | j_1 j_2 | \rangle}{\langle \pi\pi | j_1 j_2 | \rangle} < 0.27 \quad (9)$$

which then gives the range

$$1.5 < \alpha < 0.5 \quad (10)$$

Thus eventual quantification of the $K \bar{K}$ signal may be used to constrain m_{G_0} .

2. Lattice QCD suggests that the primitive scalar glueball G_0 lies at or above 1500 MeV, hence above the $I = 1$ QQ state $a_0(1450)$ and the (presumed) associated $\pi\pi$ $f_0(1370)$. Hence $E_{G_0} - E_{\pi\pi} > 0$ in the numerator of α , eqn. 6.
3. The allowed range for α (eqn. 10) enables predictions of the mass of the $\pi\pi$ state. Using eqn. 6 and $m(G) = 1509 \pm 10$ MeV, $m(\pi\pi) = 1360 \pm 40$ MeV, which dominates the error, we find [13]

$$1580 [m(\pi\pi) = 1400] < m(\pi\pi) < 1890 [m(\pi\pi) = 1320] \text{ MeV} \quad (11)$$

This is consistent with naive mass estimates whereby the $m = m_{ss} - m_{\pi\pi}$
200 \pm 300 MeV

If this were the whole story, the mixing eqn. 4 would suppress not just $K \bar{K}$ but also $\pi\pi$. However, this is where the effect of the perturbation V_2 comes into prominence: At $O(V_2)$ the glueball decays directly into pairs of glueballs or (gg) continuum and thereby into mesons whose Fock states have strong overlap with gg . To the extent that there is significant gg coupling to π^0 or to $(\pi\pi)_s$ (e.g. $\pi^0 \rightarrow gg$ and $(\pi\pi)_s$ have large intrinsic couplings notwithstanding the fact that they are supercially OZI violating) one may anticipate π^0 and $\pi\pi$ in the two-body decays of scalar glueballs. Note that the perturbation V_2 triggering $G_0 \rightarrow G_0 G_0, G_0(gg)$ or $(gg)(gg)$ will not directly feed exclusive two-body favoured states such as $\pi\pi$ nor $K \bar{K}$ since gluons are isoscalar. However, from the above phenomenology, one may anticipate $(gg)(gg) \rightarrow (\pi\pi)_s(\pi\pi)_s$ in the decay of $G(0^{++})$ and analogously for 0^+ glueballs one may anticipate π^0 or $\pi\pi$ decays.

The manifestation of this mechanism in $\pi\pi$ states involving the π^0 mesons depends on the unknown overlaps such as $\langle gg | j_1 j_2 | \pi\pi \rangle$ in the pseudoscalars. Chiral symmetry suggests that the coupling of glue to π^0 or $\pi\pi$ vanishes in the limit $m_q \rightarrow 0$ and hence occurs dominantly through their ss component, thereby favouring the π^0 :

$$\frac{\langle gg | j_1 j_2 | \pi^0 \rangle}{\langle gg | j_1 j_2 | \pi \rangle} = \frac{\langle ss | j_1 j_2 | \pi^0 \rangle}{\langle ss | j_1 j_2 | \pi \rangle} = \frac{4}{3} \quad (12)$$

This appears to be consistent with the decays of $f_0(1500)$ as we now show. Combining the amplitudes from both the QQ and G_0 components we have, in the approximation

that χ and χ^0 are approximately 50 : 50 mixtures of nn and ss ¹

$$\frac{h \rightarrow j \bar{j} G_i}{h \rightarrow j \bar{j} G_i} = \frac{h \rightarrow j \bar{j} G_{0i}}{N_G h \rightarrow j \bar{j} G_i} + \left(\frac{1 + \alpha}{2} \right) = (0.90 \pm 0.20) \quad (13)$$

from R_1 (eqn. 1) and

$$\frac{h \rightarrow j \bar{j} G_i}{h \rightarrow j \bar{j} G_i} = \frac{h \rightarrow j \bar{j} G_{0i}}{N_G h \rightarrow j \bar{j} G_i} + \left(\frac{1 - \alpha}{2} \right) = (0.53 \pm 0.11) \quad (14)$$

from R_2 (eqn. 2). Hence using the range eqn. 10 for α one finds with the + sign in eqn. 13 and the - sign in eqn. 14

$$1.8 < \frac{h \rightarrow j \bar{j} G_{0i}}{h \rightarrow j \bar{j} G_i} < 1.5 \quad (15)$$

A small breaking of chiral symmetry is consistent with this.

An upper limit for α^2 can be obtained by comparing the ratio of amplitudes (eqn. 13) for $f_0(1500)$ decay to that for $f_0(1370)$. We find [13]: $\alpha^2 < 0.2$. The magnitudes of the partial widths of $1500=1370$ also fit with the $G \rightarrow Q \bar{Q}$ mixing scenario. The above analysis shows that the decay amplitudes from the $Q \bar{Q}$ component of are all at $O(\alpha)$ and so we expect

$$\frac{\Gamma(f_0(1500) \rightarrow j \bar{j})}{\Gamma(f_0(1370) \rightarrow j \bar{j})} \sim O(\alpha^2) \quad (16)$$

Empirically this ratio is smaller than 0.4 and hence we again find $\alpha^2 < 0.2$.

This hypothesis implies that there are small G_0 admixtures in nn, ss :

$$\begin{aligned} q \frac{1 + \alpha^2}{1 + 2\alpha^2} j \bar{s} i &= j \bar{s} i + \alpha \frac{1}{2} G_{0i} \\ q \frac{1 + \alpha^2}{1 + 2\alpha^2} j \bar{n} i &= j \bar{n} i + \alpha \frac{1}{2} G_{0i} \end{aligned} \quad (17)$$

which in turn implies that the sum of the partial widths of the two states is

$$\Gamma(\chi_n) + \Gamma(\chi^0) \approx \Gamma(nn); \quad (18)$$

in accord with quark model estimates [17, 18].

The smallness of α then also implies that χ_s decays essentially like an ss state with the branching ratios to $K \bar{K}$ and χ^0 dominating over χ_n and with χ_n much suppressed. If 3P_0 quark pair creation is important in the decay dynamics, the values of the branching ratios and total width may be strongly mass dependent [13, 15, 18]

The quantitative predictions of our analysis depend on the apparent suppression of $f_0(1500)$ decay to $K \bar{K}$. Thus detailed study of $pp \rightarrow f_0(1500) K \bar{K}$ can be seminal (i) in confirming

¹The actual numbers vary slightly when a mixing angle of -17.3° [24] is used, see ref. [13]

the $K\bar{K}$ suppression, (ii) in confirming the $K_0(1430) \rightarrow K\bar{K}$ and $a_0(1450) \rightarrow \pi\pi$ and KK , (iii) in quantifying the signal for $f_0(1370)$ and $f_0(1500)$ and (iv) in isolating the predicted $s\bar{s}$ member of the nonet. Furthermore, study of radiative decays $f_0(1500) \rightarrow \pi^0\pi^0(\gamma)$ may probe the flavour content in the QQ mixing. To the extent that $\pi\pi$ and $s\bar{s}$, the amplitude ratios will be

$$f_0(1500) \rightarrow \pi\pi : \pi^0\pi^0 : \pi^+\pi^- = 1 : 1 : 3 \quad (19)$$

It seems clear that new dynamics, beyond simple QQ , is operating at the $f_0(1500)$. The simplest explanation within strong QCD is that a glueball excitation is seeding the phenomena, in particular the clarity of the signals in channels that are believed favourable to glue. It is also tantalising that the signal appears particularly sharp in pp ! [25] which might be consistent with direct production of a 0^+ glueball above 2 GeV (in line with lattice predictions) decaying via the V_2 process into G_0G_0 with consequent affinity for $f_0(1500) \rightarrow \pi\pi$. An excitation curve for this process could be interesting.

We thank T. Barnes, M. Benayoun, K. Bowler, D. Bugg, Y. Dokshitzer, G. Gounaris, A. Grigorian, R. Kenway, E. Klempt, H. Lipkin, J. Paton, S. Spanier, M. Teper, D. Wyler and B. Zou for helpful discussions.

References

- [1] F.E. Close, Rep. Prog. Phys. 51 (1988) 833
- [2] D.V. Bugg, in Proc. of Int. Symp. on Medium Energy Physics, Beijing, Aug. 1994 (unpublished); D.V. Bugg et al., "Further amplitude analysis of $\pi^+ \pi^- \rightarrow 4\pi$ ", Phys. Lett. B (in press)
- [3] T.A. Armstrong et al., Phys. Lett. B 228 (1989) 536; S. Abatzis et al., Phys. Lett. B 324 (1994) 509
- [4] D. Alder et al., Phys. Lett. B 201 (1988) 160
- [5] V.V. Anisovich et al., Phys. Lett. B 323 (1994) 233
- [6] C. Amisler et al., Phys. Lett. B 342 (1995) 433;
- [7] C. Amisler et al., Phys. Lett. B 291 (1992) 347
- [8] C. Amisler et al., Phys. Lett. B, to appear
- [9] C. Amisler et al., Phys. Lett. B 340 (1994) 259
- [10] T.A. Armstrong et al., Phys. Lett. B 307 (1993) 399; B 307 (1993) 394
- [11] G. Bali et al. (UKQCD), Phys. Lett. B 309 378 (1993)
- [12] D. Weingarten, Nucl. Phys. B (Proc. Suppl.) 34 (1994) 29
- [13] C. Amisler and F.E. Close, "Evidence for Glueballs" Rutherford Appleton Lab report CCL-TR-95-003 (to be submitted to Phys. Rev. D)
- [14] N. Isgur and J. Paton, Phys. Rev. D 31 (1985) 2910
- [15] R. Kokoski and N. Isgur Phys. Rev. D 35 (1987) 907
- [16] C. Amisler et al., Phys. Lett. B 333 (1994) 277
- [17] S. Godfrey and N. Isgur, Phys. Rev. D 32 (1985) 189
- [18] E. Aklis, T. Barnes and E. Swanson, unpublished (private communication from T. Barnes)
- [19] D. Morgan and M. Pennington, Z. Phys. C 48 (1990) 623; F.E. Close, Proc. 26th Int. Conf. HEP, Dallas, 1992, p. 562
- [20] Z.P. Li, F.E. Close and T. Barnes, Phys. Rev. D 43 (1991) 2161; F.E. Close and Z.P. Li, Z. Phys. C 54 (1992) 147
- [21] L. Gray et al., Phys. Rev. D 27 (1983) 307

- [22] C. Am sler, in P roc. of 27th Int. Conf. on H igh Energy Physics, G lasgow , 1994, p. 199
- [23] Y .D okshitzer, private com m unication
- [24] C .Am sler et al., Phys. Lett. B 294 (1992) 451
- [25] J. Ludem ann (C rystal Barrel Collaboration), P roc. Low Energy $\bar{p}p$ Conf. LEAP '94, B led, Slovenia, 1994

Figure Captions

Figure 1: The effect of perturbation V_1 causes $Q\bar{Q}$ decay (a) or G_0 mixing with $Q\bar{Q}$ (b). The effect of V_2 on G_0 causes decay to two glueballs (c).

Figure 2: Invariant couplings 2 to two pseudoscalars for an isoscalar $0^{++} Q\bar{Q}$ meson as a function of nonet mixing angle for a pseudoscalar mixing angle of 17.3° (up to a common arbitrary normalization constant). Full curve: π^0 ; dashed curve: η ; dotted curve: η' ; dashed-dotted curve: $K\bar{K}$. Ideal mixing occurs at 35.3° ($s\bar{s}$) or at 125.3° ($\bar{u}u$).

Figure 3: Gluonium- $Q\bar{Q}$ mixing involving the energy denominator $E_G - E_{Q\bar{Q}}$

This figure "fig1-1.png" is available in "png" format from:

<http://arxiv.org/ps/hep-ph/9505219v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arxiv.org/ps/hep-ph/9505219v1>

This figure "fig1-3.png" is available in "png" format from:

<http://arxiv.org/ps/hep-ph/9505219v1>