

P T H A 95/12

hep-ph/9505354

M ay 1995

Angular Distribution and Polarization of Photons in the Inclusive Decay $b \rightarrow s \gamma$

M .G remm , F .K ruger^y and L .M .S ehgal^z

III. Physikalisches Institut (A), RW TH Aachen

D -52074 Aachen, G erm any

A B S T R A C T

We study the angular distribution of photons produced in the inclusive decay of a polarized b , $b \rightarrow s \gamma$, using the technique of heavy quark effective theory. Finite non-perturbative corrections are obtained relative to the free quark decay $b \rightarrow s \gamma$. These corrections affect significantly the intensity and polarization of photons emitted at small angles relative to the b spin direction.

E lectronic address: gremm@physik.rwth-aachen.de

^yE lectronic address: krueger@physik.rwth-aachen.de

^zE lectronic address: sehgal@physik.rwth-aachen.de

I. INTRODUCTION

The decay $b \rightarrow s \gamma$, calculated on the level of free quarks, has three interesting features:

- (i) The photon is monochromatic, its energy spectrum being

$$\frac{1}{dy} \frac{d\Gamma}{dy}_{\text{FQM}} = (y - y_0) ; \quad y = \frac{2E}{m_b} ; \quad y_0 = 1 - \frac{m_s^2}{m_b^2} ; \quad (1.1)$$

- (ii) The photon is emitted preferentially backwards relative to the spin of the b-quark, the angular distribution being

$$\frac{1}{d\cos\theta} \frac{d\Gamma}{d\cos\theta}_{\text{FQM}} = \frac{1}{2} \left(1 - \frac{m_b^2 - m_s^2}{m_b^2 + m_s^2} \cos\theta \right) ; \quad (1.2)$$

- (iii) The photon is predominantly left-handed, the angular distribution for helicities $\lambda = \pm 1$ being

$$\begin{aligned} \frac{1}{d\cos\theta} \frac{d\Gamma_{+}}{d\cos\theta}_{\text{FQM}} &= \frac{1}{2} \frac{m_s^2}{m_b^2} (1 + \cos\theta) \left(1 + \frac{m_s^2}{m_b^2} \right) ; \\ \frac{1}{d\cos\theta} \frac{d\Gamma_{-}}{d\cos\theta}_{\text{FQM}} &= \frac{1}{2} (1 - \cos\theta) \left(1 + \frac{m_s^2}{m_b^2} \right) ; \\ P_{++} &= \frac{m_b^2 - m_s^2}{m_b^2 + m_s^2} ; \end{aligned} \quad (1.3)$$

Whereas feature (i) is purely kinematical (reflecting the two-body nature of the decay), features (ii) and (iii) are specific consequences of the standard model, in which the effective Hamiltonian governing $b \rightarrow s \gamma$ has the structure

$$\begin{aligned} H_{\text{eff}} &= -\frac{4G_F}{2} \frac{e}{16\pi^2} V_{tb}V_{ts}^* C_7(m_b) s \left(m_b P_R + m_s P_L \right) b\gamma^\mu \\ &\quad -\frac{4G_F}{2} \frac{e}{16\pi^2} V_{tb}V_{ts}^* C_7(m_b) s \quad b\gamma^\mu ; \end{aligned} \quad (1.4)$$

leading to a decay width

$$\Gamma_{\text{FQM}}(b \rightarrow s \gamma) = \frac{G_F^2 m_b^5}{32\pi^4} |V_{tb}V_{ts}^*|^2 |C_7(m_b)|^2 \left(1 + \frac{m_s^2}{m_b^2} \right) \left(1 - \frac{m_s^2}{m_b^2} \right)^3 ; \quad (1.5)$$

The purpose of this paper is to study how the characteristics of $b \rightarrow s$, summarized in Eqs. (1.1)-(1.3), are altered when the b -quark is embedded in a polarized b baryon, and when the final s -quark is a part of a hadronic system that is summed over. These changes have their origin in the "Fermi motion" of the b -quark within the hadron, as well as its spin-dependent interaction with the environment, and can be parametrized using the method of heavy quark effective theory (HQET) [1,2]. In Sec. II, we outline the method of calculation, and discuss our results in Sec. III.

II. HQET AND THE DECAY $b \rightarrow s X_s$

The differential decay rate for the inclusive decay $b(p) \rightarrow X_s(p_X) + (p)$ in the standard model can be written as

$$d\Gamma_{X_s, \text{pol}}^X = \frac{1}{(2\pi)^4} (p \cdot p_X - p^2) \frac{d^3 p_X}{(2\pi)^3 2E_X} \langle h_b | j_e^\gamma(0) | X_s \rangle \langle X_s | j_e^\gamma(0) | h_b \rangle \frac{d^3 p}{(2\pi)^3 2E} ; \quad (2.1)$$

where the effective Hamiltonian H_e is given in (1.4). Using the optical theorem, and writing the phase space element as

$$\frac{d^3 p}{(2\pi)^3 2E} = \frac{m_b^2}{32\pi^2} y dy d\cos\theta ; \quad (2.2)$$

Eq. (2.1) can be rewritten in the form

$$\frac{d\Gamma}{dy d\cos\theta} = \frac{G_F^2 m_b^2}{2^7 \pi^5} |V_{tb} V_{ts}|^2 |j_\gamma(m_b)|^2 y \text{Im} T(y; \cos\theta) ; \quad (2.3)$$

where T is given by

$$T = E - T ;$$

with

$$E = \int d^4x \, \bar{\psi} \gamma^\mu \psi \, A_\mu(x) \quad (2.4)$$

and

$$T = i \int d^4x \, e^{ip \cdot x} \bar{h}_b \gamma^\mu T^N b(x) \gamma^\nu s(x); s(0) = b(0) \gamma^\mu \gamma^\nu \quad (2.5)$$

The time-ordered product appearing in T can be expanded in powers of $1/m_b$, using methods described in [28]. To order $1/m_b^2$, we obtain

$$T(y; \cos \theta) = 2y^2 m_b^3 \frac{1}{(y - y_0 - i)} \left(1 + K \frac{5}{3} - \frac{7}{3} \frac{y}{(y - y_0 - i)} + \frac{2}{3} \frac{y^2}{(y - y_0 - i)^2} \right) \left(1 + \frac{m_s^2}{m_b^2} \right) \cos \theta + (1 + \bar{b} + K) \frac{5}{3} - \frac{7}{3} \frac{y}{(y - y_0 - i)} + \frac{2}{3} \frac{y^2}{(y - y_0 - i)^2} \left(1 - \frac{m_s^2}{m_b^2} \right) : \quad (2.6)$$

The imaginary part $\text{Im} \, T(y; \cos \theta)$ is then obtained by the formal replacement

$$\begin{aligned} \frac{1}{y - y_0 - i} &\rightarrow \delta(y - y_0) ; \\ \frac{1}{(y - y_0 - i)^2} &\rightarrow -\delta'(y - y_0) ; \\ \frac{1}{(y - y_0 - i)^3} &\rightarrow -\frac{1}{2} \delta''(y - y_0) : \end{aligned} \quad (2.7)$$

The distributions in Eq. (2.7) yield physically meaningful results only if one integrates over the photon energy (see e.g. Ref. [8]). The leading term proportional to

$(y - y_0)$ reproduces the result of the free quark model (FQM). The corrections to the free quark result, involving $\delta'(y - y_0)$ and $\delta''(y - y_0)$, are parametrized by two phenomenological constants K and $\bar{b}$, defined as

$$K = \bar{h}_b \gamma^\mu \gamma^\nu \frac{(iD)^2}{2m_b^2} h_v \gamma_\mu \gamma_\nu \quad (2.8)$$

$$(1 + \bar{b})s = \bar{h}_b \gamma^\mu \gamma^\nu \gamma^5 b \gamma_\mu \gamma_\nu \quad (2.9)$$

(Refs. [4,8] use a parameter κ_1 related to K by $K = \kappa_1 = 2m_b^2$). Integration of Eq. (2.3) over y then yields the angular distribution

$$\frac{d(\Gamma_b \rightarrow X_s \gamma)}{d\cos\theta} = \frac{1}{2} \frac{1}{F_{QM}} [(1 - K) + (1 + \kappa_b - K) \frac{m_b^2 - m_s^2}{m_b^2 + m_s^2} \cos\theta] : \quad (2.10)$$

If the decay $\Gamma_b \rightarrow X_s \gamma$ is calculated for a fixed photon helicity ($\lambda = \pm 1$), the angular distributions are

$$\frac{d(\Gamma_b \rightarrow X_s \gamma)}{d\cos\theta} = \frac{1}{2} \frac{1}{F_{QM}} \frac{m_s^2}{m_b^2} [(1 - K) + (1 + \kappa_b - K) \cos\theta] \left[1 + \frac{m_s^2}{m_b^2} \right]^{-1} ; \quad (2.11)$$

$$\frac{d(\Gamma_b \rightarrow X_s \gamma)}{d\cos\theta} = \frac{1}{2} \frac{1}{F_{QM}} [(1 - K) + (1 + \kappa_b - K) \cos\theta] \left[1 + \frac{m_s^2}{m_b^2} \right]^{-1} : \quad (2.12)$$

Eqs. (2.10)-(2.12) are the HQET analogs of the free quark result (1.2)-(1.3).

III. DISCUSSION OF RESULTS

To evaluate the HQET corrections, we need an estimate of the parameters K and κ_b . Following Ref. [5] we use $K \approx 0.01$, although this parameter has a large uncertainty. From the fact that $d\Gamma/d\cos\theta \geq 0$ for all scattering angles (independent of the value of m_s or K) we must have $\kappa_b < 0$ (see Eq. (2.10)). We will use the value $\kappa_b = -\frac{2}{3}K$ suggested in Ref. [3].¹ This is in agreement with the bound derived in Ref. [9].

From the angular distribution in Eq. (2.10), we infer that the effect of non-perturbative corrections is to reduce slightly the forward-backward asymmetry in the photon emission. In Fig. 1, we plot the fraction of decays producing a photon

¹This value is obtained if one neglects the contributions coming from the double insertion of the chromomagnetic operator [3].

in the forward cone $\cos \theta_0 < \cos \theta < 1$:

$$I(\cos \theta_0) = \frac{\int_{\cos \theta_0}^1 d\cos \theta \frac{d}{d\cos \theta}}{\int_0^1 d\cos \theta \frac{d}{d\cos \theta}} : \quad (3.1)$$

The result is compared with that in the free quark model.

From the results in Eqs. (2.11) and (2.12), we obtain the polarization of the photon as a function of its direction:

$$P(\cos \theta) = \frac{d_{++} = d\cos \theta}{d_{++} = d\cos \theta + d_{--} = d\cos \theta} : \quad (3.2)$$

Once again, as shown in Fig. 2, corrections to the free quark model are significant for photons emitted in the near-forward direction.

To summarize, the angular distribution of inclusively produced photons in the decay $\Lambda_b \rightarrow X_s \gamma$ of a polarized Λ_b baryon can be calculated unambiguously in HQET. This distribution tests the structure of the underlying Hamiltonian H_e in a way that is not possible by studying the mesonic decay $B \rightarrow X_s \gamma$. The angular distribution is a well defined observable which involves no divergences of the type that appear in the calculation of the energy spectrum (delta functions and derivatives thereof). Our calculation may be viewed as an illustration of the utility (and limitation) of the HQET approach in determining the effects of hadronic binding on the decay of a heavy quark. The deviations from free quark decay in the reaction $\Lambda_b \rightarrow X_s \gamma$ are found to be globally small, but are significant for photons emitted in the forward direction.

ACKNOWLEDGEMENTS

We acknowledge helpful discussions with Gabi Köpp, and collaboration on a related paper [10]. This work has been supported in part by the Deutsche Forschungsgemeinschaft (DFG) under Grant No. Se 502/4-1, which we gratefully acknowledge.

REFERENCES

- [1] M .Neubert, Phys. Rep. 245 (1994) 259, and references therein; N .Isgur and M .B .W ise, in Heavy Flavours, edited by A .J. Buras and M .Lindner (W orld Scienti c, Singapore, 1992).
- [2] J.Chay, H .Georgi and B .G rinstein, Phys. Lett. B 247 (1990) 399.
- [3] A .F .Falk and M .Neubert, Phys. Rev. D 47 (1993) 2982.
- [4] A .F .Falk, M .Luke and M .J. Savage, Phys. Rev. D 49 (1994) 3367.
- [5] A .V .M anohar and M .B .W ise, Phys. Rev. D 49 (1994) 1310.
- [6] I.I.B igi, M .Shifm an, N .G .U raltsev and A .I.Vainshtein, Int. J. M od. Phys. A 9 (1994) 2467.
- [7] T .M annel, Nucl. Phys. B 413 (1994) 396.
- [8] M .Neubert, Phys. Rev. D 49 (1994) 4623.
- [9] J.G .K omer and D .P irjol, Phys. Lett. B 334 (1994) 399.
- [10] M .G rem m , G .K opp and L.M .Sehgal, preprint P ITHA 95/2, to appear in Physical Review D .

FIGURE CAPTIONS

Figure 1 The fractional intensity I of photons in the forward cone $\cos \theta_0 = \cos \theta = 1$.

Figure 2 The photon polarization P in the inclusive b decay as a function of the photon direction with $K = 0.01$ and $\theta_b = \frac{2}{3}K$.




FIG .1:




FIG .2: