

CHIRAL PREDICTION FOR THE N S(WAVE) SCATTERING LENGTH a TO ORDER $O(M^{-4})$

V. Bernard^{a#1#2}, N. Kaiser^{b#3}, Ulf-G. Meißner^{c#4}

^aCentre de Recherches Nucleaires, Physique Theorique
BP 28, F-67037 Strasbourg Cedex 2, France

^bTechnische Universität München, Physik Department T 39
James-Frank-Straße, D-85747 Garching, Germany

^cUniversität Bonn, Institut für Theoretische Kernphysik
Nussallee 14-16, D-53115 Bonn, Germany

Abstract

We evaluate the S-wave pion-nucleon scattering length a in the framework of heavy baryon chiral perturbation theory up to and including terms of order M^{-4} . We show that the order M^{-4} piece of the isovector amplitude at threshold, T_{thr} , vanishes exactly. We predict for the isovector scattering length, $0.088 M^{-1} \leq a \leq 0.096 M^{-1}$.

March 25, 2024

^{#1}also at Laboratoire de Physique Theorique, Institut de Physique, 3-5 rue de l'Université, F-67084 Strasbourg Cedex, France.

^{#2}electronic address: bernard@cmhp4.in2p3.fr

^{#3}electronic address: nkaiser@physik.tu-muenchen.de

^{#4}electronic address: meissner@thp.uni-bonn.de

Chiral symmetry severely constrains the interactions of the strongly interacting particles at low energies, first formulated in terms of current algebra (CA). One of the most splendid successes of CA was Weinberg's prediction for the S-wave pion-nucleon scattering lengths [1],

$$a_{\text{CA}}^+ = 0 ; \quad a_{\text{CA}} = \frac{M}{8 F^2} \frac{1}{1 + M/m_p} = 0.079 M^{-1} ; \quad (1)$$

with $M = 139.57 \text{ MeV}$ the charged pion mass, $m_p = 938.27 \text{ MeV}$ the proton mass, $F = 92.5 \text{ MeV}$ the pion decay constant and the superscripts $+$ refer to the isoscalar and isovector N amplitude, respectively. The Karlsruhe(Helsinki) phase shift analysis of N scattering [2] leads to $a = 0.092 \pm 0.002 M^{-1}$ and $a^+ = -0.008 \pm 0.004 M^{-1}$, impressively close to the CA prediction, eq.(1). The Karlsruhe(Helsinki) analysis is based on dispersion relations thus requiring a solid data base as input. However, over the last few years there has been some controversy about the low-energy N data which has not yet been settled. Consequently, the uncertainties in a are presumably larger and even the sign of a^+ could be positive. A more direct way to get a handle at these zero momentum (i.e. threshold) quantities is the measurement of the strong interaction shift (δ_{1S}) and the decay width (Γ_{1S}) in pionic atoms. The PSI-ETH group has recently presented first results of their impressive measurements in pionic deuterium and pionic hydrogen [3] [4] [5] (for earlier work in the field see e.g. [6], [7] [8]). The S-wave scattering length are related to δ_{1S} and Γ_{1S} via the Deser-type formulae [9],

$$\delta_{1S} = C (a^+ + a) (1 + \epsilon) ; \quad \Gamma_{1S} = 4 q C (1 + 1/P) [a (1 + \epsilon)]^2 ; \quad (2)$$

with q the momentum of the π^0 , P the Panofsky ratio and the constant $C = 4 E_{1S} r_B$ is proportional to the point-Coulomb binding energy, E_{1S} , of the system in the 1S state divided by the pion Bohr radius, r_B . ϵ and δ describe electromagnetic corrections to the scattering lengths, see [10]. The consequent analysis of the data leads to [5] $a = 0.096 \pm 0.007 M^{-1}$ and $a^+ = -0.0077 \pm 0.0071 M^{-1}$. If one combines the pionic hydrogen shift measurement with the one from the pionic deuterium, one has $a = 0.086 \pm 0.002 M^{-1}$ and $a^+ = 0.002 \pm 0.001 M^{-1}$. The largest uncertainty comes from the width measurement of pionic hydrogen. Both determinations are consistent within one standard deviation. We conclude that a is larger than the CA value and that a^+ is consistent with zero.

Heavy baryon chiral perturbation [11] allows to systematically calculate the corrections to the CA predictions, eq.(1), for the S-wave scattering lengths. This topic was already addressed in ref.[12]. In this note, we want to further sharpen the prediction for the isovector scattering length. In particular, we will calculate all contributions which are of order M^{-4} and furthermore give a realistic estimate of the uncertainties for the one loop result (which was not done in [12]). The starting point is the effective pion-nucleon Lagrangian,

$$L_N = L_N^{(1)} + L_N^{(2)} + L_N^{(3)} + L_N^{(4)} \quad (3)$$

where the superscript (i) refers to the number of derivatives or quark mass insertions. The structure of L_N is discussed in detail in the review [13] and a pedagogical introduction can be found in [14]. The S-wave scattering lengths can be calculated from the on-shell forward N scattering amplitude $T^{ba} = T^+(\!)^{ba} + i^{bac} c T^-(\!)$ with $\! = p - q = m_p$, the pion lab energy. Evaluation at threshold $\! = M$ gives them as $a = T_{\text{thr}} = (4(1 + M/m_p))$. The one-loop result for T_{thr} to order M^4 reads

$$T_{\text{thr}} = \frac{M}{2F^2} + \frac{g_N^2 M^3}{8m_p^4} + \frac{M^3}{16F^4} \left[1 - 2 \ln \frac{M}{\Lambda} \right] + B^r(\Lambda) M^3 + T_{\text{thr}}^{i4} + O(M^5); \quad (4)$$

where the chiral corrections at order M^3 come from the expansion of the nucleon pseudovector Born term, the loop corrections and the counterterms. T_{thr}^{i4} denotes the order M^4 contribution to the isovector forward N amplitude at threshold to be discussed below. We note that the contribution from the Born term is just 1% of a_{CA} . To determine the counterterm contribution, one needs to pin down the value of $B^r(\Lambda)$ which is a particular combination of some low-energy constants from $L_N^{(3)}$ [12] [13]. In fact, these low-energy constants are not known and have been estimated in [12] via resonance exchange,

$$B^r(\Lambda) = B^{\text{res}} = \frac{g_N^2}{2m_p^2 m^2} \left[Z - \frac{1}{2} \right] + \frac{R g_N^2}{8m_p^2 (m_p + m_N)^2}; \quad (5)$$

In eq.(5), we have explicitated the corresponding (1232) and the $N(1440)$ contributions. The range of the various resonance parameters like R and Z is discussed in [12]. This procedure induces a spurious dependence on the scale of dimensional renormalization (Λ) which enters through the divergences from the one loop contribution. In [12], Λ was fixed at the mass of the (1232) since this resonance is most important in the estimation of the contact term $B^r(\Lambda)$. However, in the framework of resonance saturation this scale should be varied between $M = 550 \text{ MeV}$ and $m_N = 1.44 \text{ GeV}$. Would we be able to fix $B^r(\Lambda)$ from some data, the prediction for a would, of course, be Λ -independent.

We now turn to the calculation of the order M^4 corrections. First, contact terms from $L_N^{(4)}$ can not contribute to T_{thr} at $O(M^4)$ because of the crossing property of the isovector forward N amplitude,

$$T^-(\!) = -T^+(\!); \quad (6)$$

with $\!$ the pion lab energy. Thus we are left with one-loop graphs with exactly one insertion from $L_N^{(2)}$. Many of these diagrams cancel with their crossed partners due to the crossing property, eq.(6), and specific properties of the loop functions entering, as exemplified in Fig.1. In particular, the terms with one insertion proportional to $c_{1,2;3,4}$ from $L_N^{(2)}$ vanish because

$$J_0^N(\!) + J_0^N(-\!) = \frac{1}{4} \frac{q^2}{M^2 - \!^2} = 0; \quad (7)$$

at threshold, $\ell = M$ (the pertinent loop functions are given in appendix B of [13]). However, this does not hold for the thirteen diagrams shown in Fig 2. These remaining graphs can be grouped into five classes as indicated in Fig 2. Straightforward calculation gives

$$T_{\text{thr}}^{\ell^4} = \frac{g_A^2 M^4}{128 m_p F^4} (9 + 8 \ell^2 + 5 \ell^4 - 20 \ell^6) = 0; \quad (8)$$

where we have exhibited the contributions from the various classes separately. This result is renorm invariant of the renormalization of the isovector charge of the nucleon. Consequently, the next chiral corrections appear at order M^5 , from two-loop graphs, one-loop graphs with one insertion from $L_N^{(3)}$, counterterms and so on. They are suppressed by four powers of $M = (4 F)$ compared to the leading order term, eq.(1) and are thus expected to be small, even though the exact numerical coefficients can be of order ten (as indicated from the $O(M^3)$ correction to the CA result).

Varying the various resonance parameters which enter the estimate of B^{Res} and the scale within their bounds, we find as a conservative estimate for the one-loop prediction of a^- ,

$$0.088 M^{-1} \leq a^- \leq 0.096 M^{-1}; \quad (9)$$

which is consistent with the various empirical values discussed before and 10:::20% larger than the CA prediction [1]. As already stressed in [12], it is the chiral loop correction at order M^3 which closes the gap between the lowest order (CA) prediction and the empirical value. An indication of the size of the next corrections can be obtained by writing the one-loop result as $a^- = a_{\text{CA}} (1 + \delta_1) \approx a_{\text{CA}} \exp(\delta_1)$. The next correction follows to be $\delta_1^2 = 2$ which is of the order of 1:::2% of a_{CA} . We stress that this should only be considered indicative. The uncertainty in the prediction eq.(9) could be further decreased if one could determine $B^{\pi}(\ell)$ from some other process.

To summarize, we have sharpened the chiral perturbation theory prediction for the isovector N S-wave scattering length a^- by showing that the term of order M^4 in T_{thr} is identical to zero. Varying all uncertainties entering the one-loop expression for a^- [12], we arrived at the band, eq.(9), which is consistent with the existing data. Concerning the isoscalar scattering length a^+ , no firm prediction can be given at present as discussed in [12] since there are large cancellations which make the result for a^+ very sensitive to some not very accurately known counterterms.

Acknowledgements

We are grateful to Jurg Leisi and Daniel Sigg for triggering our interest and communication of results before publication.

References

- [1] S.W einberg, Phys. Rev. Lett. 17, 616 (1966)
- [2] R .K och, Nucl. Phys. A 448, 707 (1986)
- [3] A .Badertscher et al., Nucl. Instr. & M ethods A 335, 470 (1993)
- [4] D .Chatellard et al., "D etermination of the S-wave scattering length in pionic deuterium with a high resolution crystal spectrom eter", Jan. 1995, subm itted to Phys. Rev. Lett.
- [5] D .Sigg, Ph.D .Thesis ETH No. 11049, Zurich, 1995 (unpublished)
- [6] A .Forster et al., Phys. Rev C 28, 2374 (1983)
- [7] D .Bovet et al, Phys. Lett. B 153, 231 (1985)
- [8] W .Beer et al, Phys. Lett. B 261, 16 (1991)
- [9] S. D eser, M .L. G oldberger, K .Baum ann and W .Thirring, Phys. Rev 96, 774 (1954)
- [10] D .Sigg et al., "E lectrom agnetic Corrections to the S-W ave Scattering Lengths in P ionic Hydrogen", subm . for publication
- [11] E .Jenkins and A .V .M anohar, Phys. Lett. B 255, 558 (1991)
- [12] V .Bernard, N .K aiser and U lf-G .M ei ner, Phys. Lett. B 309, 421 (1993)
- [13] V .Bernard, N .K aiser and U lf-G .M ei ner, "Chiral D ynam ics in Nucleons and Nuclei", preprint CRN 95/3 and TK 95 1, Int. J M od. Phys. E (1995), in print
- [14] U lf(G .M ei ner, C zech. J. Phys. 45, 153 (1995)

Figure C aptions

Fig.1 Cancellation of diagram s at order M^4 due to the crossing property, eq.(6) . Solid and dashed lines denote nucleons and pions, respectively. The heavy dot denotes an insertion from $L_N^{(2)}$.

Fig.2 C lasses of non{vanishing diagram s at order M^4 . For notations, see Fig.1.






Fig. 1



I



II

III

IV

V

Fig. 2