

CERN-TH/95-100
 archive/9506208

WWGENPV A Monte Carlo event generator for four-fermion production in $e^+e^- \rightarrow W^+W^- \rightarrow 4f$

Guido MONTAGNA
 Istituto Nazionale di Fisica Nucleare, Sezione di Pavia
 Via A. Bassini, 6 - 27100 PAVIA - ITALY

Oreste NICROSINI¹
 CERN, Theory Division
 CH-1211 - Geneva 23 - SWITZERLAND

Fulvio PICCINI
 Istituto Nazionale di Fisica Nucleare, Sezione di Pavia
 Via A. Bassini, 6 - 27100 PAVIA - ITALY

Program classification: 11.1

The Monte Carlo program WWGENPV, designed for computing distributions and generating events for the four-fermion process $e^+e^- \rightarrow W^+W^- \rightarrow 4f$, is described. It is based on the calculation of the exact tree-level matrix element of the four-fermion reaction and includes initial-state radiation in the leading-log approximation within the structure function approach. The program can be used in a two-fold way: as a Monte Carlo integrator for weighted events, providing predictions for the total cross section, the W invariant-mass distribution, the radiative energy and invariant-mass loss; as a true event generator of unweighted events, useful for simulation purposes.

Submitted to Computer Physics Communications

CERN-TH/95-100
 April 1995

¹On leave from INFN, Sezione di Pavia, Italy

PROGRAM SUMMARY

Title of program : WWGENPV

Computer: DEC VAX, HP/APOLLO, IBM/RS6000; Installation: INFN,
Sezione di Pavia, via A. Bassi 6, 27100 Pavia, Italy

Operating system : VMS, UNIX

Programming language used: FORTRAN 77

Memory required to execute with typical data: 500 KByte

No. of bits in a word: 32

The code has not been vectorized

Subprograms used: random number generator RANLUX [1]; routines from the
CERN Program Library are also used.

No. of lines in distributed program, including test data, etc.: 2135.

Correspondence to:

MONTAGNA@PV.INFN.IT;

NICROSINI@VXCERN.CERN.CH, NICROSINI@PV.INFN.IT;

PICCINI@PV.INFN.IT

Keywords: high energy electron-positron collisions, W -pair on-shell production, four-fermion final state, LEP 2, QED corrections, electron structure functions, experimental cuts, Monte Carlo integration, event generation, importance sampling.

Nature of physical problem

The precise measurement of the W -boson mass M_W constitutes a primary task of the forthcoming experiments at the high energy electron-positron collider LEP 2 ($2M_W \leq \sqrt{s} \leq 210 \text{ GeV}$). This experimental goal will be possible through the study of the four-fermion reaction $e^+e^- \rightarrow W^+W^- \rightarrow 4f$. A meaningful comparison between theory and experiment requires an accurate description of the fully exclusive process $e^+e^- \rightarrow W^+W^- \rightarrow 4f$, including the main effects of radiative corrections, with the final goal of providing predictions for the realistic distributions measured by the experiments [2].

Method of solution

In order to efficiently perform the high-dimensional integration (due to the four-fermion final state) in the presence of realistic cuts, the total cross section, the energy and invariant-mass loss and the W -invariant-mass distributions are

computed by means of a Monte Carlo integration for weighted events. For simulation purposes, the program can also be used as an event generator that provides a sample of unweighted events, defined as the components of the four final-state fermion momenta, plus the radiative variables of the incoming electron and positron, plus $p_{\bar{s}}$, stored into proper n-tuples. The importance-sampling technique [3] is employed to take care of the peaking behaviour of the integrand, both in the integration and in the generation modes.

Restrictions on the complexity of the problem

Electroweak processes originating the same four-fermion final state of the double-resonating diagrams $e^+e^- \rightarrow W^+W^- \rightarrow 4f$ (background processes) are not included in the present version of the program. The treatment of the initial-state radiation is limited to the gauge-invariant subset of the leading-logarithmic corrections which are the most important contribution from the phenomenological point of view. Furthermore, the kinematics of the event in the presence of the initial-state radiation is limited to the collinear approximation, i.e. $p_T \approx p_L$ effects are neglected for the time being.

Typical running time

As integrator, the code needs about 5 h of HP 9000/735 for generating 10^8 weighted events; the relative errors on the cross section and on the energy and invariant-mass losses are about $2 \cdot 10^{-4}$ and $6 \cdot 10^{-4}$, respectively. The generation of a sample of 10^4 unweighted events requires about 3 min on the same system.

Unusual features of the program : none

References

- [1] F. James, Comput. Phys. Commun. 79 (1994) 111.
- [2] G. Montagna, O. Nicrosini, G. Passarino and F. Piccinini, Phys. Lett. B 348 (1995) 178, and references therein.
- [3] F. James, Rep. Prog. Phys. 34 (1980) 1145.

LONG WRITE { U P

1 Introduction

The precise measurement of the W -boson mass M_W , and more generally the study of its physical properties, constitutes a primary task of the forthcoming experiments at the high energy electron-positron collider LEP 2 ($\sqrt{s} = 2M_W \approx 210$ GeV).

A large amount of theoretical work has been performed up to now, following essentially two main streams.

On the one hand, one can find semi-analytical works, whose leitmotiv is computing analytically as much phase-space integrations as possible [1]. Typically, these works give (QED-corrected) distributions for the invariant masses of the W , integrating a priori all the other independent phase-space variables, from which total cross sections can be easily computed. After such an analytical effort, the semi-analytical codes developed along this line are very fast, precise and efficient. They are hence typically well suited for fitting purposes, their main drawback being the lack of a full reconstruction of the true event phase-space configuration.

On the other hand, one can find Monte Carlo codes. In principle they can guarantee the full event reconstruction, since they handle the four-momenta of the final-state particles. Of course, the accuracy with which this goal is achieved depends on the dynamics built-in in the code; in particular, the exclusive control of all the features of the full four-fermion event, including all the possible correlations between final-state particles, requires the knowledge of the exact four-fermion matrix element, at least in the Born approximation (for instance, no isotropic decays of the intermediate bosons can be allowed).

Common to both the above-mentioned approaches, is the problem of treating QED corrections, which, as is well known, are large and strongly dependent on the experimental cuts applied. In particular, it happens that the energy lost in the beam pipe (which at the leading-logarithmic level is well described by initial-state radiation in the collinear approximation) represents an important part of the systematic error on M_W . Therefore one needs also a reliable model for, at least, initial-state radiation.

Some Monte Carlo codes are already available for the four-fermion processes of interest at LEP 2 [2]. Nevertheless, the critical comparison between independent approaches to a problem has always been a useful and powerful tool for understanding the theoretical accuracy of the results. The aim of this work is to present and describe in some detail WWGENPV, a Monte Carlo event generator for $e^+e^- \rightarrow W^+W^- + 4f$, including initial-state QED corrections

in the structure functions formalism.

2 Theoretical Background and General Features

The theoretical formulation implemented in the program has been described in some detail in [3]. To calculate the cross section of the $e^+e^- \rightarrow W^+W^- + 4\gamma$ process in the presence of the initial-state radiation, the basic formula reads:

$$\sigma(s) = \int dx_1 dx_2 D(x_1; s) D(x_2; s) d\mathbb{P}[S] \frac{d\sigma_0}{d\mathbb{P}[S]}; \quad (1)$$

In this equation, $d\mathbb{P}[S]$ denotes the volume element in the 7-dimensional phase space connected to the independent variables of the hard scattering reaction in the centre-of-mass frame; $d\sigma_0 = d\mathbb{P}[S]$ is the exact differential kernel cross section in the Born approximation, limited at present to the W -signal diagrams. The inclusion of the complete set of background diagrams is in progress. At any rate their effect can be cut-off by proper selection criteria on the final-state invariant masses. Finally $D(x; Q^2)$ is the electron structure function describing initial-state radiation [4]. For its explicit expression the following form is chosen:

$$D(x; s) = \frac{\exp\left\{-\frac{1}{2} \ln^2 \frac{1}{x} - \frac{3}{4} \ln \frac{1}{x} - \frac{\gamma_E}{2}\right\}}{1 + \frac{1}{2} \ln^2 \frac{1}{x}} \left[\frac{1}{2} (1-x)^{-\frac{1}{2}} - \frac{1}{4} (1+x) + \frac{1}{32} \left(4(1+x) \ln(1-x) + 3(1+x) \ln x - 4 \frac{\ln x}{1-x} \right) \right]; \quad (2)$$

with

$$L = 2 - (L - 1); \quad (3)$$

where $L = \ln(s/m^2)$ is the collinear logarithm, γ_E is the Euler constant and $\Gamma(z)$ is the gamma function. The first exponentiated term of Gribov-Lipatov form describes multiphoton soft radiation, the second and third ones hard collinear bremsstrahlung. This choice of the structure function is a consistent solution of the Renormalization Group evolution equation at the leading-logarithmic level, and guarantees that, after an expansion down to $O(\alpha_s)$, the well-known factorized and hence universal bremsstrahlung spectrum is exactly recovered.

With the choice of the independent variables adopted in [3], the differential kernel cross section is proportional to the tree-level squared matrix element as follows:

$$\frac{d\sigma}{d\mathbf{P}S} \propto \frac{M^2}{(a^2/3 + b^2/3 + c)(a^2/2 + b^2/2 + c^0)} : \quad (4)$$

Therefore, in the kernel cross section two kinds of "singularities" can be recognized, namely the one related to the vector boson propagators (on $1/2$), the second arising from the Jacobian appearing in (4) (on $2/3$). To efficiently perform the Monte Carlo integration in the presence of the above peaking structure, the importance-sampling technique [5] has been employed. For instance, for the W -boson propagator singularities, an integration must be performed, which has the form

$$I = \int_{x_m^2}^{x_M^2} dx^2 \frac{f(x^2)}{(x^2 - a^2)^2 + b^2}; \quad (5)$$

where $x^2 = \frac{2}{s}$, $a^2 = M_W^2/s$, $b^2 = 2M_W^2/s^2$ and the function $f(x^2)$ is regular. Multiplying and dividing the integrand by the weight function

$$p(x^2) = \frac{N^{-1}(x_m^2; x_M^2)}{(x^2 - a^2)^2 + b^2}; \quad (6)$$

with

$$\begin{aligned} N(x_m^2; x_M^2) &= \int_{x_m^2}^{x_M^2} dx^2 \frac{1}{(x^2 - a^2)^2 + b^2} \\ &= \frac{1}{b} \operatorname{Arctg} \frac{x_M^2 - a^2}{b} - \operatorname{Arctg} \frac{x_m^2 - a^2}{b}; \end{aligned} \quad (7)$$

one obtains

$$I = \int_{x_m^2}^{x_M^2} dx^2 p(x^2) \frac{p^{-1}(x^2) f(x^2)}{(x^2 - a^2)^2 + b^2}; \quad (8)$$

where the quantity in square brackets is no longer singular at $x^2 = a^2$. The integral I can be rewritten as

$$I = \int_0^1 dc \frac{p^{-1}(x^2(c)) f(x^2(c))}{(x^2(c) - a^2)^2 + b^2}; \quad (9)$$

introducing the cumulative variable c as

$$c(x^2) = \int_{x_m^2}^{x^2} dz^2 p(z^2); \quad (10)$$

Therefore, by generating y at between 0 and 1 an accurate estimate of I can be obtained.

Concerning the singularities arising from the Jacobian factor, the integrations to be performed are of the following kind:

$$I = \int_{y_m}^{y_M} dy \frac{g(y)}{(y - y_1)(y_2 - y)} : \quad (11)$$

In this case the importance sampling can be applied by introducing the weight function

$$p(y) = N^{-1}(y_m; y_M) (y_2 - y)^{-\frac{1}{2}} (y - y_1)^{-\frac{1}{2}} ; \quad (12)$$

with

$$\begin{aligned} N(y_m; y_M) &= \int_{y_m}^{y_M} dy (y - y_m)^{-\frac{1}{2}} (y_M - y)^{-\frac{1}{2}} \\ &= 2 \operatorname{Arcsin} \frac{y_M - y_1}{y_2 - y_1} \operatorname{Arcsin} \frac{y_m - y_1}{y_2 - y_1} : \end{aligned} \quad (13)$$

When including initial-state QED corrections, in order to perform the importance sampling on the generation of the radiative variable x in the soft region $x \rightarrow 1$, the weight function is chosen as the all-order soft part of the structure function, i.e.

$$p(x) = N^{-1}(x_m; x_M) \frac{1}{2} (1 - x)^{-\frac{1}{2}} ; \quad (14)$$

with the normalization factor $N(x_m; x_M)$ given by:

$$\begin{aligned} N(x_m; x_M) &= \int_{x_m}^{x_M} dx \frac{1}{2} (1 - x)^{-\frac{1}{2}} \\ &= (1 - x_m)^{-\frac{1}{2}} - (1 - x_M)^{-\frac{1}{2}} : \end{aligned} \quad (15)$$

With respect to the original formulation of QED corrections given in [3], besides a different choice of the structure function, the Coulomb singularity, which is important in the threshold region, is included in the code according to the very recent formulae given in [6]. For details and links with other approaches to the problem, the reader is referred to the original literature.

3 Program Structure

After the initialization of the Standard Model parameters and of the electromagnetic quantities, the calculation within the program proceeds as described

in the following. As a first step, the random number generator RANLUX is called to extract a vector of twelve components which are converted from REAL*4 to REAL*8 precision in order to be consistent with the calculational environment, which is in REAL*8 precision. The first two random numbers are used to generate the radiative variables $x_{1,2}$ according to the distribution given by the soft exponentiated part of the structure function; eight random numbers are employed to fully reconstruct the kinematics of the four-fermion hard scattering process in the centre-of-mass frame: seven of them are needed to generate the independent variables previously introduced for the kernel reaction, one to choose between the two possible solutions of the non-linear constraint. Given the independent variables, all the invariants related to the $2 \rightarrow 4$ kernel reaction are computed. The variables $s_{1,2,3}^2$ and s_2 are, in particular, generated by means of the importance-sampling procedure to take care of the peaking behaviour of the integrand, as described in the previous section. The other variables $(z_{1,3}; \epsilon_1)$ are generated according to the uniform distribution in the physical range relative to an extrapolated set-up. When initial-state radiation is switched on, the four-momenta of the outgoing fermions are reconstructed in the laboratory frame, taking into account the Lorentz boost due to the emission of unbalanced electron and positron radiation and given by the γ_L Lorentz factor $\gamma_L = (x_1 - x_2)/(x_1 + x_2)$. At this point, one further random number is generated to perform a random rotation around the beam axis to account for a trivial overall azimuthal angle. The last random number is required in the generation branch for hit-or-miss. After that, a call to SUBROUTINE CUTUSER is performed in order to implement the rejection algorithm on the final-state particles. If the event is accepted, the squared matrix element in Born approximation is returned by a call to SUBROUTINE BORN. In conclusion, at the end of the loop over the events, the averages and squared averages are cumulated to provide the "weighted events" Monte Carlo results for the required observables and their numerical errors.

When the program runs as a generator of unweighted events, after the hit-or-miss procedure, the components of the four final-state fermions momenta, plus the radiative variables of the incoming electron and positron, plus $p_{\overline{s}}$, are stored into a proper n-tuple. Each event stored in the n-tuple has the structure $(x_1; x_2; E_b; q_1; q_2; q_3; q_4)$, where E_b is the beam energy and q_i are the three-momenta of the outgoing particles, assumed to be ultrarelativistic. In this case, the cross section for unweighted events is given as output.

4 Input

Some input parameters and flags are introduced in the program in order to allow the user to select the output of interest. A short explanation of their meaning is addressed in the following and is also printed when running the code interactively.

OGEN (CHARACTER*1)

It controls the use of the program as a Monte Carlo integrator for weighted events (OGEN = I) or as a Monte Carlo event generator of unweighted events (OGEN = G). If the first choice is requested, the code returns in a LOGFILE the value of the cross section for weighted events, together with the average energy loss and invariant-mass loss, defined as $\langle h(2 \sqrt{s_1} \sqrt{s_2}) \rangle$ and $\langle h(1 \sqrt{s_1} \sqrt{s_2}) \rangle$, respectively. If the second choice is requested, it prints in a LOGFILE the value of the cross section for unweighted events and fills an n-tuple, EXAMPLE.DAT, containing the generated (unweighted) events. The LOGFILE contains also the value of NBIAS/NMAX. If it is > 0 , the generation procedure is biased and it is necessary to run the code again, after having properly increased the value of FMAX (default = 1000).

RS (REAL*8)

The centre-of-mass energy (in GeV).

NHITWMAX (INTEGER)

Required by the integration branch. It gives the maximum number of calls for the Monte Carlo loop.

NHITMAX (INTEGER)

Required by the event-generation branch. It is the maximum number of hits for the hit-or-miss procedure.

IQED (INTEGER)

This flag allows the user to switch on/off the contribution of the initial-state radiation. If IQED = 0 the distributions are computed in lowest-order approximation, while for IQED = 1 the initial-state QED corrections are included in the calculation. This choice can be performed both for the integration and for the generation mode.

ODIS (CHARACTER*1)

Required by the integration branch. It selects the kind of experimental distribution. For ODIS = T the program computes the total cross section (in pb) of the process, eventually including the cuts supplied by the user in SUBROUTINE CUTUSER (see below); for ODIS = W (only possible with OGEN = I) the W invariant-mass distribution is returned (in pb/GeV). In this case one can choose between the two opposite charged W -bosons, namely W^- (IWCH = 1) and W^+ (IWCH = 2), and the invariant-mass of interest must be entered by the user in GeV.

OWIDTH (CHARACTER*1)

It allows a different choice of the value of the W -width. OWIDTH = Y means that the tree-level Standard Model formula for the W -width is used; OWIDTH = N requires that the W -width is supplied by the user in GeV. This freedom in the choice of the W -width has been introduced so as to simplify the comparisons of the results of our code with those of other programs.

NSCH (INTEGER)

The value of NSCH allows the user to choose the calculational scheme for the weak mixing angle and the gauge coupling. Three choices are available. If NSCH=1, the input parameters used are $G_F; M_W; M_Z$ and the calculation is performed at tree level. If NSCH = 2 or 3, the input parameters used are $(Q^2); G_F; M_W$ or $(Q^2); G_F; M_Z$, respectively, and the calculation is performed using the QED coupling constant at a proper scale Q^2 , which is requested as further input.

OCOUL (CHARACTER*1)

This flag allows the user to switch on/off the contribution of the Coulomb correction. If OCOUL = Y (N) the distributions are computed with (without) the above electromagnetic effect.

5 Test Run Output

The typical calculations that can be performed with WWGENPV are illustrated by the following examples of test run output.

Sam ple 1 Calculation of total cross section, average energy loss and average invariant-m ass loss for 10^8 weighted events, including initial-state QED corrections.

Sam ple 2 Calculation of total cross section for 10^4 unweighted events and lling the n-tuple EXAMPLE.DAT, including initial-state QED corrections.

The output of sam ple 1 can be obtained w ith the follow ing input:

OGEN = 'I'

The integration branch is chosen.

RS = 176.D0

The centre-of-m ass energy is xed at 176 GeV .

NHITWMAX = 100000000

The m axim um number of calls is xed.

IQED = 1

Initial-state QED corrections are included.

ODIS = 'T'

The total cross section is com puted (together w ith the corresponding average energy and invariant-m ass loss).

OWIDTH = 'Y'

The W -boson w idth is com puted at the tree level in the Standard M odel.

NSCH = 2

(Q^2) , G_F and M_W are used as input param eters.

1/ALPHAI = 128.07D0

The value of $1/(Q^2)$ is entered.

OCOUL = 'N'

The Coulomb correction is not included.

The output of sample 2 can be obtained with the following modifications:

OGEN = 'G'

The generation branch is chosen.

NHITMAX = 10000

The number of generated events is fixed.

Moreover the value of ODIS is not required.

Both for the integration and generation branch, the output returns the values of the "Feynman rules" GVE, GAE, GWF and GWWZ linked to the vertices occurring in the diagrams of $e^+e^- \rightarrow W^+W^- \rightarrow 4f$ as follows:

$$\begin{aligned} Z_{ff} \text{ vertex} &= (GVE + GAE_{\gamma}) \\ W_{ud;l_1} \text{ vertex} &= GWF (1 + \gamma_5) \\ W W Z \text{ vertex} &= GWWZ \quad \text{Lorentz structure:} \end{aligned}$$

6 Program Implementation

In this section a short description of each routine implemented in the program is given. For random-number generation the routine RANLUX by F. James is used and inserted in the code. For more details about RANLUX the reader is referred to the proper literature [7].

```
SUBROUTINE CUTUSER(EVENT,CUTFLAG)
REAL*8 CUTFLAG
DIMENSION EVENT(15)
```

To be supplied by the user. The subroutine allows the user to implement any kind of experimental cut on the four-momenta of the final-state fermions. Globally, the event is specified by 15 ingredients: the radiative variables $x_{1,2}$ (EVENT(1), EVENT(2)), the beam energy (EVENT(3)) and the components of the three-momenta of the four outgoing fermions (EVENT(I), with $I = 4;15$). The energies are simply derived by the mass-shell condition for ultrarelativistic fermions. The reference frame is assumed to be with the z-axis along the direction of the incoming positron.

The routine returns a flag (CUTFLAG = 0, 1), which controls if the event is accepted (CUTFLAG = 1) or rejected (CUTFLAG = 0) by the selection criteria.

```

SUBROUTINE BORN(S,X13,X14,X34,X35,X15,X16,X23,X24,
               X25,X26,X36,X45,X46,X56,SQM,IOVFL,IOEFL,
               IOES,JOEFL,JOES)
IMPLICIT REAL*8 (A-H,O-Z)

```

It returns in SQM the squared tree-level matrix element associated to the s- and t-channels Feynman diagrams of the kernel process $e^+e^- \rightarrow W^+W^- \rightarrow 4f$. The squared matrix element is computed numerically, squaring the amplitude obtained by means of SCHOONSCHIP [8] in the framework of the helicity-amplitude formalism [9] for massless fermions. If the effect of the Coulomb correction is required, the value returned for SQM is given by:

$$SQM = SQM_{fl} + \gamma_c v; \quad (16)$$

where γ_c represents the Coulomb correction factor and v the relative velocity of the two W -bosons (see [6] for more details and explicit formulae).

```

FUNCTION DOP(X)
IMPLICIT REAL*8 (A-H,O-Z)
COMMON/EM/RL,ETA,BETA,SDELTA

```

The QED structure function, adopted to account for the radiation emitted by the incoming leptons, is computed according to the analytical expression reported above. It is given by the sum of an exponentiated term of the Gribov-Lipatov form, associated to soft multiphoton emission, with up to $O(\alpha_s^2)$ contributions describing hard bremsstrahlung in collinear approximation. Actually, in order to perform importance sampling on the generation of the radiative variable x in the soft region $x \rightarrow 1$, the following formula is implemented in FUNCTION DOP:

$$\overline{D}(x) = D(x) = p(x); \quad (17)$$

where $p(x)$ is the weight function given by (14).

Acknowledgements

The authors would like to thank the INFN, Sezione di Pavia, for having provided computer resources. They are also grateful to Giorgio Fumagalli and Valerio Vercesi for helpful assistance in numerical and graphical problems. Useful discussions and exchanges of information with D. Bardin and D. Charlton are gratefully acknowledged.

References

- [1] M . Cacciari, A . Deandrea, G . Montagna and O . Nicrosini, Z . Phys. C 52 (1991) 421;
D . Bardin, M . Bilenky, A . O lhevsky and T . R iem ann, Phys. Lett. B 308 (1993) 403;
V . Fadin, V A . Khoze and A D . Martin, Phys. Lett. B 311 (1993) 311.
- [2] F . A . Berends, R . K leiss and R . P ittau, Nucl. Phys. B 424 (1994) 308 and B 426 (1994) 344; Comput. Phys. Comm un. 85 (1995) 437;
H . Anlauf et al., Comput. Phys. Comm un. 79 (1994) 487;
G . J. van Oldenborgh, P . J. Franzini and A . Borrelli, Comput. Phys. Comm un. 83 (1994) 14;
T . Sjstrand, Comput. Phys. Comm un. 82 (1994) 74; CERN -TH .7112/93.
- [3] G . Montagna, O . Nicrosini, G . Passarino and F . Piccinini, Phys. Lett. B 348 (1995) 178.
- [4] E . A . Kuraev and V . S. Fadin, Sov. J. Nucl. Phys. 41 (1985) 466;
G . Altarelli and G . Martinelli, in Physics at LEP , CERN Report 86-02, J. Ellis and R . Peccei eds. (CERN , Geneva, 1986);
O . Nicrosini and L . Trentadue, Phys. Lett. B 196 (1987) 551; Z . Phys. C 39 (1988) 479.
- [5] F . James, Rep. Prog. Phys. 34 (1980) 1145.
- [6] V . S. Fadin, V A . Khoze, A D . Martin and A . Chapovsky, Coulomb Effects in W^+W^- Production, Preprint DTP/94/116 and references therein.
- [7] F . James, Comput. Phys. Comm un. 79 (1994) 111.
- [8] SCHOONSCHIP, a program for symbol handling by M . Veltman, see H . Strubbe, Comput. Phys. Comm un. 8 (1974) 1.
- [9] G . Passarino, Nucl. Phys. B 237 (1984) 249.

TEST RUN OUTPUT

Sample 1

SQRT(S) = 176.0 GEV
M_Z = 91.1888 GEV
G_Z = 2.4974 GEV
M_W = 80.23 GEV
G_W = 2.03367033063195 GEV
STH2 = 0.2310309124510679

GVE = -1.409737267299689E-02
GAE = -0.185794027211796
GWF = 0.230409927395451
GWWZ = -0.571479454308384

RANLUX INITIALIZED BY RLUXGO FROM SEEDS 2 0 0

NCALLS = 100000000

XSECT FOR WEIGHTED EVENTS

EFF = 0.999998000004
XSECT = 13.52642598410257 +- 2.200641628422248E-03 (PB)

ENERGY LOSS
MEAN EG = 1.18064898022852 +- 6.041171111535717E-04 (GEV)

INVARIANT-MASS LOSS
MEAN IML = 1.17777558571609 +- 6.023399436481718E-04 (GEV)

Sam ple 2

SQRT(S) = 176.0 GEV
M_Z = 91.1888 GEV
G_Z = 2.4974 GEV
M_W = 80.23 GEV
G_W = 2.03367033063195 GEV
STH2 = 0.2310309124510679

GVE = -1.409737267299689E-02
GAE = -0.185794027211796
GWF = 0.230409927395451
GWWZ = -0.571479454308384

RANLUX INITIALIZED BY RLUXGO FROM SEEDS 2 0 0

NHIT = 10000

XSECT FOR UNWEIGHTED EVENTS

EFF = 1.341451799624662E-02
XSECT = 13.4145179962466 +- 0.133242395670936 (PB)
NBIAS/NMAX = 0.0