# THE DOUBLET-TRIPLET SPLITTING PROBLEM AND HIGGSES AS PSEUDOGOLDSTONE BOSONS

LISA RANDALL<sup>1</sup> and CSABA CSAK I<sup>2</sup>

Center for Theoretical Physics, Laboratory for Nuclear Science and Department of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139{4307 1: E-mail: lisa@ctptopmit.edu 2: E-mail: csaki@mithsmit.edu

The doublet-triplet splitting problem is probably the most signi cant challenge to supersymmetric GUT theories. In this talk, we review potential solutions and their problem atic aspects. We also present a complete consistent realization of our preferred solution, higgses as pseudogoldstone bosons, and discuss some distinctive aspects of its phenom enology.

W eak scale supersymmetry might be the solution to the hierarchy problem. Recently in light of the very accurate unication of gauge couplings, supersymmetric GUT theories have received a good deal of attention. However, the status of supersymmetric GUT theories is perhaps not so rosy as it appears. The essential problem is to nd a consistent picture of the Higgs sector which requires very light (weak scale) doublets but very heavy (GUT scale) triplets. The purpose of this talk is to emphasize the importance of solving the doublet-triplet splitting problem in establishing the credibility of supersymmetric GUT theories, and to suggest a possible solution.

The outline is as follows. We rst review the basic problem and the status of solutions. We will see that most solutions have some problem atic feature, and in general are quite complicated. We then present our favored solution, namely Higgses as pseudo-Goldstone bosons. We show that we can build a complete consistent model of a supersymmetric grand united theory in which the doublet Higgses are light and the triplet Higgses are heavy. We implement a potential which protects the Higgs mass is protected by an accidental global symmetry which is accurate at su ciently high order in Planck mass suppressed terms to keep the Higgs boson as light as required on phenomenological grounds.

The existence of a complete consistent theory is very important, both from the vantage point of the viability of SUSY GUTs, and as we will discuss, from the view -

This work is supported in part by funds provided by the U S.D epartment of Energy (D O E .) under contract # DE-FC 02-94ER 40818. Talk presented at SUSY '95 conference and at the Joint M eeting of the Johns H opkins W orkshop and PASCOS.M II -CTP-2460. hep-ph/9508208

point of phenom enology. We will show that the predictions of our speci cm odeldier signicantly from what would be predicted for a \generic" GUT theory, particularly with respect to avor changing neutral currents.

 Introduction: D oublet-Triplet Splitting, The Q uagm ire of Supersymmetric G U T s

The major phenom enological motivation for weak scale supersymmetry is the resolution of the hierarchy problem . W hile the Higgs doublet mass parameters should be in the 100 GeV range, higgsino mediated proton decay argues (in the simplest models) that the Higgs triplet mass parameters are in the  $10^{16}$  GeV range. In the minimal SU (5) model, a parameter must be chosen with accuracy of order  $10^{13}$ . A though this might be technically natural, it is not very compelling as a solution to the hierarchy problem.

In the absence of simple solutions, we are faced with the question of whether it makes sense to believe a model with a parameter as small as  $10^{13}$ . C learly, a better solution is warranted and it is important to explicitly realize it. For those of you who are still not convinced this is a problem, consider the analogous situation for technicolor. Technicolor is not ruled out by precision electroweak measurements.<sup>a</sup> It is the di culty and complication of making a model which can generate masses without excessive avor changing neutral currents which puts technicolor models in disfavor.

Now consider the status of supersymmetric GUTs. In some ways, the situation is not so dierent. There are serious fundamental problems from the standpoint of model building. As I have emphasized, the most serious problem is doublet-triplet splitting, but there are other aspects one would like to see simply addressed in a model, such as generating mass, suppressing dangerous avor changing neutral currents, and suppressing dangerous CP violation. Without a model, it is di cult to believe in the existence of the theory. Without an economical solution to the avor problem in technicolor theories, they were put in question. But at least the \Higgs" sector works. In supersymmetric GUTs, the theory is stymied at an even earlier stage. W hat should be emphasized here is that the doublet-triplet splitting problem is not a peripheral issue, but an essential part of the supersymmetric GUT theory. Furthermore, any aspect of the phenomenology which probes the GUT structure can depend in an essential way on this part of the theory. O nce we explicitly construct a model, we will exploit it to test the sensitivity of certain predictions, and compare to what might

<sup>&</sup>lt;sup>a</sup>For example, the models of Refs. [1,2] contain only a minimal technicolor sector and are therefore acceptable.

have been derived from a \generic" GUT.

## 2. Review of Proposed Solutions

There have been a number of solutions proposed to avoid ne tuning and naturally distinguish the doublet and triplet H iggs masses. They generally have clever group theory structure, which prevents a doublet H iggs mass in the renormalizable Lagrangian. However, as we will see, most models su er from some unsatisfactory feature. I will review a few of the more promising suggestions here and evaluate their satisfactory and unsatisfactory features.

Before proceeding, let us present some criteria for a good model. First, there should be no unduly small numbers, that is no tuning of parameters. This is a somewhat loosely de ned thing (one person's tuned parameter might be perfectly acceptable to someone else. We suggest the following subjective, but probably adequate, de nition. Any number which you would write only in scientic notation is not permitted. The number 0.01 might be OK according to this de nition, but 10<sup>5</sup> is not.

It should be emphasized that the ratio required between the doublet and triplet H iggs masses is roughly (M<sub>G</sub>=M<sub>P</sub>)<sup>4</sup>. So if one believes that the allowed P lanck suppressed operators are present, it is not su cient to suppress the H iggs mass in the renorm alizable Lagrangian. It must be true that the potential respects whatever symmetry is necessary to fourth order in M<sub>P</sub> suppression.

A third restriction is that the model should perm it successful gauge coupling uni cation within errors, both experim ental and theoretical. This is important since the major phenom enological motivation for GUTs is the uni cation of couplings. This is very constraining{ it does not allow for even one additional pair of doublets or triplets.

Now simply stated, GUT theories mandate simple relations between the elds in a single GUT representation. It is very dicult to see how elds from a single representation can have such di erent masses. Not suprisingly the \solutions" are often very complicated and are not often entirely adequate, though many are quite clever. It is very dicult to nd a model in which the doublet-triplet splitting is solved naturally and in which there is no ne tuning.

In M inim al SU (5), the best proposed solution is the M issing Partner M echanism [3,4]. The idea here is to give the triplet a mass through a D irac mass term involving a more complicated representation of SU (5) which has the property that it contains a triplet but not a doublet. The 50 is the sm allest representation with this feature. O ne therefore constructs the mass term s

W 
$$5_{\rm H} 50_{\rm H} h75_{\rm H} i + {}^{0}5_{\rm H} 50_{\rm H} h75_{\rm H} i$$

so that the triplets, but not the doublets are massive. A model with additional

symmetry to forbid a direct mass term for the 5 and 5 incorporates an additional 75 [4].

This is a very nice idea, but seems unlikely to be the resolution of the dilem ma. There are several problems with this model. First of all, the large rank of the representations is disturbing. From a theoretical perspective, one has yet to not string theory examples containing these large rank representations. A nother problem is that the gauge coupling grows very rapidly, so that the theory is strongly coupled not far above the GUT scale. A though this might be acceptable, it is certainly a problem at the level of nonrenorm alizable operators which we discuss shortly.

A further problem is that one cannot leave the states in the remainder of the 50 m assless, since they contribute like an extra doublet pair to unication, which we know is too much. One can solve this problem for example by adding a mass term M 5050 (though this is forbidden by the symmetry of Ref. [4]). But then there is nothing in the symmetry structure of the theory which could forbid the term (5) (5) (75) (75)=M  $_{\rm p}$  which is the product of two allowed terms in the superpotential divided by a third and is therefore allowed by the symmetry, no matter what it is. If one believes P lanck suppressed operators consistent with the symmetries are present, the doublet has much too big a mass. This problem is exacerbated in the case the coupling blows up at a low scale, because it is probably the associated strong scale which would suppress such operators.

A more compact in plementation of the M issing Partner M echanism was proposed for F lipped SU (5) [5]. The idea is again to pair up the H iggs with \som ething else". Here the som ething else is a  $10_H$  for the  $5_H$  and a  $10_H$  for the  $5_H$ , the subscript refers to the H iggs sector to distinguish these elds from the ordinary matter elds. These  $10_H$  and  $10_H$  elds are not dangerous because the nonsinglet nontriplet elds are eaten when SU (5) breaks. Hence one has eliminated the necessity for the additional m ass term.

The  $10_{\rm H}$  contains a 3 but no color singlet weak doublet. The  $10_{\rm H}$  and  $10_{\rm H}$  get VEVs breaking the SU (5) U (1) gauge group to the standard model. The triplet Higgs in  $5_{\rm H}$  pairs with the triplet in  $10_{\rm H}$  and the remaining elds in the  $10_{\rm H}$  are eaten by the massive vector bosons.

This model might work. However, ipped SU (5) is not really a uni ed model since the gauge group is SU (5) U (1) which is not a sem is in ple group. If it is embedded in a larger gauge group, the problem should be solved in the context of the larger gauge group. The other feature we ind disturbing is that there are many assumptions about the vacuum structure. At tree level, there is a D - at, F - at direction, and the loop corrections have to be such as to generate the desired minimum. In Ref. [5], the correct ratio of gauge and Yukawa couplings was assumed so that the VEV's for  $10_{\rm H}$  and  $10_{\rm H}$  were at the GUT scale while the  $5_{\rm H}$  and  $5_{\rm H}$  VEV's are small and the VEVs of an SU (5) singlet generated the  $\setminus$  " term . It is certainly easier to evaluate the vacuum when it is determined at tree level, as it will be in our preferred model. O ther solutions have been proposed form odels which incorporate SU (5) as a subgroup, for example SO (10). The  $\mbox{m}$  issing VEV " or D in opoulos-W ilczek m echanism [6] is probably the most popular SO (10) solution. The idea is again to pair the triplet and not the doublet H iggs with som ething else so that the triplet, but not the doublet is massive. In this model, the way this is done is that the VEV aligns so that the triplet, but not the doublet, gets a mass. (This would not have been possible in m inim al SU (5) due to the tracelessness of the ad pint.)

Again, this mechanism seems very nice at rst glance, but worrisom e at second. For if this were all there was, you would have four light doublets, not two. You need to give the extra doublets a mass, and the problem is how to do this without reintroducing a problem with proton decay. A series of papers by Babu, Barr and M ohapatra [7,8,9,10] showed possible ways to make the DW mechanism into a more com plete m odel.

The rst example [7] had two sectors giving VEVs aligning in di erent orientations, one responsible for the triplet m ass, and one responsible for the doublet m ass. They thereby achieved strong suppression of proton decay. There was an additional eld to complete the breaking of SO (10) to the standard m odel, and an additional adjoint to couple the two sectors together (elim ininating a m assless G oldstone) w ithout m isaligning the DW m echanism. The total eld content in this m odel is uncom fortably large { 3(16) + 3(10) + 3(45) + 2(54) + 16 + 16, leading to fairly big threshold corrections and the blow ing up of the gauge coupling before M <sub>P1</sub>. O ther problem s w ith this particular m odel was that som e operators which would have been allowed by the symmetries of the m odel needed to be forbidden, and that nonrenorm alizable P lanck m ass suppressed operators could be dangerous.

This last problem was addressed in their second model, where they sacrice strong suppression of proton decay but generate a natural model, in the sense that they include all operators permitted by their assumed symmetry structure. The eld content of this model was 3(16) + 2(10) + 3(45) + (54) + (126) + (126). Discrete symmetries were su cient to forbid any unwanted terms from the potential. However, the eld content was still quite large, and high rank representations were required.

The third model incorporated a smaller eld content and no high rank representations, so it should be more readily obtainable from string models. In this model, the authors achieved the DW form with higher dimension operators, so no (54) was required. There was a 16 + 16 to complete the breaking to the standard model.

However, without three adjoints, there were intermediate scale pseudo-Goldstone bosons. The authors resolved this problem by cancelling the fairly large corrections to unication of couplings (due to the light charged elds) by large threshold corrections. A lithough this might work, it is at the edge of parameter space.

A nother nice m odel based on SO (10) is the m odel of B abu and M ohapatra [10] which allows for a 10-16 m ixing and therefore a H iggs sector which distinguishes the up and down quark m asses. However, this m odel had a few small (but not very

sm all) parameters, a at direction and therefore vacuum degeneracy at tree level, extra singlets, and a complicated superpotential.

To sum m arize, there are som e interesting m odels in the literature, prim arily based on clever group theory structure. However m ost m odels su er from one of the follow ing problem s.

There is the problem of actually implementing the potential to get the desired m in imum and light Higgses. The m in imum can sometimes be destabilized with higher order terms. Also some models have at directions so the vacuum needs to be carefully thought through.

It is necessary to ensure the light particle spectrum is compatible with gauge coupling uni cation. Most solutions rely on pairing up the triplet higgsinos (not doublet) with \som ething else". \Som ething else" can be a problem (with gauge uni cation).

The particle representation is cumbersome. This leads to the questions of whether it is derivable from strings or whether the coupling blows up before the Planck scale. In any case, models with large particle content seem unappealing and unlikely.

The problem is clear. M inim al SU (5) relates doublets and triplets! A lm ost always, the solution relies on a comprom ise at the edge of parameter space or tuned parameters or setting some couplings to zero in the potential. This is a good introduction to the Higgses as pseudo-Goldstone bosons model which we will argue is an exception to the discussion above. Rather than relying on pairing the Higgs in complicated ways, the theory relies on a spontaneously broken symmetry under which the Higgses are Goldstone bosons. This distinguishes the doublets from the triplets in a very nontrivial way, so that it is natural to obtain light doublets when the remaining elds are heavy. The originally proposed model [11,12,13] involved gauged SU (5) symmetry and a global SU (6) symmetry which was implemented by tuning potential parameters. A better model [14,15,16,17] was later proposed which adm its the possibility for justifying the large global symmetry with discrete symmetries. In fact, as we will see, one can construct a simplem odel to implement this idea [18].

The general idea behind the H iggs as G oldstone scheme is to implement a global symmetry on the H iggs sector which is broken (explicitly) by the Y ukawa couplings to matter. The masses of the pseudo-G oldstone bosons are protected from large bop corrections by the nonrenormalization theorem. The H iggs sector can be distinguished by matter parity and consists of the adjoint (), the fundamental representation (H) and the antifundamental (H), where this refers to their representation under the gauge symmetry. Then additional global symmetry is ensured by assuming the superpotential is of the form

W (;H;H) = W () + W (H;H)

At the renorm alizable level, this can be achieved by forbidding the coupling H  $\,$  H . At higher orders, m any other couplings m ust be forbidden.

By far the nicest in plem entation of this idea (here we say how we would like the structure to work without explicitly in plem enting the potential which we do later) is based on extending m inim al SU (5) to the gauge group SU (6), with SU (6) SU (6) global sym m etry. W e will refer to this as the SU (6) m odel. The Higgses are then an adjoint, which is a 35 = 24 + 6 + 6 + 1, a fundam ental H, in a 6 = 5 + 1, and an H, in a 6 = 5 + 1 where we have given the SU (5) decom positions. The accidental sym m etry is realized if m ixing term s of the form H H are not present in the superpotential. If the elds and H; H develop VEV's of the form

then one of the global SU (6) factors breaks to SU (4) SU (2) U (1), while the other to SU (5). Together, the VEV's break the gauge group to SU (3) SU (2) U (1).

The G oldstone bosons (G B's) coming from the breaking SU (6) ! SU (4) SU (2) U (1) are (according to their SU (3) SU (2) U (1) transform ation properties):

$$(3;2)_{\frac{5}{2}} + (3;2)_{\frac{5}{2}} + (1;2)_{\frac{1}{2}} + (1;2)_{\frac{1}{2}};$$
 (3)

while from the breaking SU (6)! SU (5) the GB's are

$$(3;1) \quad \frac{1}{2} + \quad (3;1)_{\frac{1}{2}} + \quad (1;2)_{\frac{1}{2}} + \quad (1;2) \quad \frac{1}{2} + \quad (1;1)_0: \tag{4}$$

But the following GB's are eaten by the heavy vector bosons due to the supersymmetric Higgs mechanism (the gauge symmetry is broken from SU (6) to SU (3) SU (2) U (1)):

$$(3;1)_{\frac{1}{3}} + (3;1)_{\frac{1}{3}} + (3;2)_{\frac{5}{6}} + (3;2)_{\frac{5}{6}} + (1;2)_{\frac{1}{2}} + (1;2)_{\frac{1}{2}} + (1;1)_{0}:$$
(5)

Thus exactly one pair of doublets remains uneaten which can be identied with the Higgs elds of the MSSM. One can show that the uneaten doublets are in the following combinations of the elds ; H; H:

$$h_{1} = \frac{U h}{P} \frac{3V h_{H}}{9V^{2} + U^{2}};$$
(6)

$$h_{2} = \frac{U h}{P} \frac{3V h_{H}}{9V^{2} + U^{2}};$$
(7)

where  $h_H$  and  $h_H$  denote the two doublets living in the SU (6) eld H and H, while h and h denote the two doublets living in the SU (6) adjoint . In order to maintain the correct prediction for  $\sin^2$ , we need to have h i M<sub>GUT</sub>, hH i = hH i > h i.

It is important to note that the triplets in H and H are eaten and not dangerous. Only the triplet needs to be made heavy in the superpotential to avoid proton decay.

Having established the desired vacuum and symmetry structure, we now have to face the question of whether such a model exists. In fact, we would once again like to see a model with the more stingent requirement that the Higgs is massless up to order ( $M_G = M_{Pl}$ )<sup>4</sup>. Furthermore we will take the point of view that only gauge and discrete symmetries are exact. To obtain the desired structure therefore requires an accidental symmetry which is respected in nonrenormalizable terms to fourth order in 1=M<sub>Pl</sub>. We also require that the potential has no at directions so that the minimum is determined and is the desired one. In order to show that it is a viable model we also require that the model to fermion masses. We emphasize that the existence of such a model is important because there are no other complete consistent and simple models. The model [18] we constructed has simple eld content, no sm all parameters, and the above properties.

Before actually presenting an example of a model, let us rst understand the di culty. Suppose the potential is such that the minim a of and H are determined. We then require at least four terms in the potential (one can not have nonzero VEVs for and H; H with just three terms).

$$\frac{1}{M_{P1}^{a}} Tr^{a} + \frac{1}{M_{P1}^{b}} Tr^{b};$$
$$\frac{1}{M_{P1}^{2c}} (H H)^{c} + \frac{1}{M_{P1}^{2d}} (H H)^{d}$$

But then the symmetries allow

$$\frac{1}{M_{Pl}^{2c+ba3}}$$
 (H  $^{ba}$  H ) (H H )<sup>c1</sup>

Notice that the dangerous term involving both and H is suppressed by precisely the ratio of two superpotential terms. But at the minimum, all the terms in the potential should be of the same order of magnitude. This means that the ratio of elds is of the same order of magnitude as the ratio of the coe cients of the operators. To get

the required suppression of the Higgs mass would then require a ratio of coe cients of order  $(M_W = M_{Pl})!$  This is precisely what we are trying to avoid {that is tuning the Higgs mass to be small.

So we need to explore the possible loopholes in the above reasoning. One possibility is to add more elds, so that one cannot make a holom orphic function with positive powers of the elds which is allowed by the existing symmetries. But to eliminate the at directions then requires more terms in the superpotential. Furthermore we do not want very high dimension operators in the potential (because when the VEV is of order  $M_G$ , the triplets in the will get too small a mass, suppressed by ( $M_G = M_{Pl}$ ) to a large power). Also, the dimensions of the operators in the or H, H potential should not be very dimension of the ratio of couplings will be large (or small) so that all terms in the potential can be of the same order of magnitude at the minimum. But with su ciently many elds and low dimension terms in the potential to eliminate at directions, one can generally make invariant operators by more complicated extensions of the above argument. We found no examples where we obtained a satisfactory minimum at tree level and all dangerous operators were forbidden.

We conclude we need a small number to make a model. Fortunately we know there exists a small number, namely the ratio of the weak scale to the Planck scale. Even without knowing how this ratio arises, we know this small number is present in any satisfactory model.

The alternative for a small number is 0. One can also construct models which incorporate elds with zero expectation value, so that the dangerous terms involving the and H elds vanish.

Severalm odels involving one or the other of these ideas were presented in R ef. [18]. We present the simplest of the examples in the following section.

#### 3.A M odel

The eld content is the minimal eld content required, namely H , H , and . We impose an additional discrete symmetry H H : H H  $~!~~e^{2}~^{i=n}$  H H .

The superpotential is

$$W = \frac{1}{2}M Tr^{2} + \frac{1}{3} Tr^{3} + \frac{(H H)^{n}}{M_{pl}^{2n}}$$

and the potential after the inclusion of the soft breaking term s is given by

$$V(; H; H) = Tr_{M} + \frac{2}{6} \frac{1}{6} Tr^{2}f + \frac{n^{2}}{M_{Pl}^{4n}} (HH)^{2n} (H_{Pl}^{2} + H_{Pl}^{2}) + \frac{n^{2}}{M_{Pl}^{4n}} (HH)^{2n} (H_{Pl}^{2} + H_{Pl}^{2}) + \frac{n^{2}}{M_{Pl}^{4n}} (H_{Pl}^{2} + H_{Pl}^{2n}) + \frac{n^{2}}{M_{Pl}^{4n}} (H_{Pl}^{2} + H_{Pl}^{2n}) + \frac{n^{2}}{M_{Pl}^{4n}} (H_{Pl}^{2n} + H_$$

Am Tr 
$${}^{3}$$
 + A  ${}^{0}$ m  $\frac{(H H)^{n}}{M_{Pl}^{2n}{}^{3}}$  + BM m  ${}^{2}$   
+ m  ${}^{2}$  (Tr  ${}^{2}$  +  $H$   $J$  +  $H$   $J$ ) + D terms (8)

Notice that at the m inim um for H , a high dimension operator is balanced against the soft supersymmetry breaking terms. There is a m inim um with

$$hH i = hH i = \frac{m}{M_{Pl}} \prod_{m=1}^{\frac{1}{2n-2}} M_{Pl}$$

For n = 4;5;6, we get hH i 1:5  $1^{46}$ ;6  $1^{46}$ ;2  $1^{47}$  GeV. The rst mixing term allowed by  $Z_n$  is  $\frac{1}{M_{p1}^{2n-2}}$  (H H)<sup>n 1</sup> (H H). This gives mass to the pseudo-G oldstone bosons which is less than the weak scale, and therefore safe. Notice it was essential that H mass term was small for naturalness of the model. This was only permitted because the triplets in H and H were eaten.

We emphasize that this model had no elds other than an adjoint, fundam ental, and antifundam ental, which we expect to be present in any SU (n) model which breaks to the standard model and gives the necessary ferm ion masses. However, to accome odate ferm ion masses in our context and to generate a successful mass texture, we found it useful to incorporate an additional adjoint eld. W ith an additional discrete Z<sub>3</sub> symmetry (under which  $_1 ! e^{2 i=3} _{1, 2} ! e^{2 i=3} _{2}$ , H H ! H H), the superpotential takes the form

M Tr 
$$_{1}$$
  $_{2}$  +  $\frac{1}{3}$   $_{1}$ Tr  $_{1}^{3}$  +  $\frac{1}{3}$   $_{2}$ Tr  $_{2}^{3}$  +  $\frac{1}{M_{Pl}^{2n}}$  (H H )<sup>n</sup>

This is useful for constructing a massm odel in which the masses and mixing angles are naturally of order of magnitude of the ratio of VEVs and the Planck scale [18]. These are enforced through discrete symmetries acting on the fermions. One interesting aspect of these models is that the top quark is in a distinct representation from the other up type quarks [16,18]. This is required in order to give a renormalizable Y ukawa coupling to the top quark so it can get a su ciently large mass. A nother interesting aspect of the mass models is that b unication occurs naturally, because there is a unique operator which gives a mass to the b and (so there are no C lebsches distinguishing the mass).

### 4. Phenom enology

Because of the enlarged gauge group and the restricted Higgs sector, the phenom enology of the SU (6) model operates very dierently from the minimal SU (5) model. However, since there is probably not a minimal SU (5) model, since any GUT model which solves the doublet-triplet splitting problem is likely to have additional structure, it is in portant to explore the phenom enology of a model with a satisfactory Higgs sector. These predictions can dier signi cantly from \generic" results.

Here we focus on the violations of lepton avor in supersymmetric unied theories [20]. Barbieri, Hall, and Strum is point out that in supersymmetric GUTs there is large avor violation in the lepton sector due to  $t_{t}$  and that it is communicated to physics at low energies through the scalar partners. In principle, one can test SUSY GUTs through avor changing processes, such as ! e. They worked out the predictions for the minimal GUT theories. However, the SU (6) model predictions look very di erent as a function of the couplings and SUSY soft parameters from the minimal SU (5) case. Let us focus for the moment on what distinguishes the predictions of the SU (6) model from those of a \generic" GUT. First of all, the top quark is in a di erent gauge representation from the other up type quarks, so in principle there are additional avor changing e ects from gauge interactions. It turns out however these are smaller than those due to the top Yukawa coupling. Second, there are di erent possible contractions of higher dimension operators so there are di erent mixing angles for the leptons and down type quarks, reducing the predictability of the avor changing lepton process (related to unknown mixing angles). Another distinguishing aspect of our model is that there are additional potential avor dependent Yukawa couplings from interactions with heavy ferm ions (which are necessarily present in the theory for the reason of anom aly cancellation). These can however be naturally suppressed by discrete symmetries [19].

W hat turns out to be the most important num erical di erence to the prediction is the larger gauge group requiring larger representations. This makes everything run much faster, since essentially the counting factors on loop diagrams are bigger.

Let us now consider the predictions. First we bok at the top Yukawa coupling, t. There is an upper bound in this model from two things{ rst we require the top Yukawa to be perturbative up to the Planck scale, which gives a bound  $_{\rm t} < 1.12$ (compared to  $_{\rm t} < 1.56$  in m inim al SU (5). Second, we require that the stau m ass remains positive. In m inim al SU (5) the running is slower, so there is usually no such a constraint. There is also a lower bound on  $_{\rm t}$ . This comes from requiring that buni cation works. In our model, it is easy to make b- uni cation work to a level of a few percent, but di cult to make it work with greater than 10% error without com pletely destroying the prediction. W ith only this much leeway,  $_{\rm t}$  cannot be too sm all, greater than about 0.9.

Now with t constrained to be so sm all, m in in alSU (5) would predict unobservably sm all avor violations. However, because in SU (6) the masses run so much more quickly, we nd we still predict potentially testable avor changing lepton processes over most of parameter space. The details of this analysis will be given in Ref. [19]. We nd for comparable t, the rate for t e is an order of magnitude greater than form in in al SU (5).

#### 5. C onclusions

Supersymmetric GUTs seem very nice, but they are theoretically problematic. The most important issue from a model building perspective is to understand how the doublet-triplet splitting problem can be resolved. Without a solution, the existence of the theory is dicult to support.

M ost solutions have some problem. However, the Higgs as pseudo-Goldstone boson solution appears to be an exception. We have constructed a rather simplem odel where the Higgs is light because of additional accidental global symmetry present in the theory. This gives a natural resolution to the doublet-triplet splitting (and the ) problem. We have a complete consistent example. Moreover, with an explicit realization of the model, one can explore its phenom enological consequences. This gives a new and maybe even more realistic perspective on low energy phenem enology. It is important to understand the range of possibilities which follow from a complete model.

One might also interpret the di culty in constructing GUT models as indicative that the theory is not uni ed below the Planck scale. In this case, we will ultimately want to understand the discrepancy between the uni cation scale and the Planck scale. If we are to apply the same standards applied to the GUT models, we would also want to better understand the string vacuum and resolve the problem of moduli proliferation and the associated vacuum degeneracy. A fler all, all the solutions appeared beautiful until one tried to implement them explicitly in realistic non netuned natural models which gave the correct value for  $\sin^2$ .

#### 6. A cknow ledgem ents

W e thank our collaborator Zurab Berezhiani. W e would also like to thank G reg Anderson, M arcela Carena, D iego Castaro and Carlos W agner for useful conversations.

- 1. L.Randall, Nucl. Phys. B 403 (1993) 122.
- 2. H. Georgi, Nucl. Phys. B 416 (1994) 699.
- 3. S.D im opoulos and F.W ilczek, in Erice Sum m er Lectures, Plenum, New York, 1981;
  - H.Georgi, Phys. Lett. B 108 (1982), 283;
  - A.Masiero et al., Phys. Lett. B 115 (1982) 380.
- 4. B.Grinstein, Nucl. Phys. B 206 (1982), 387;
- 5. I. Antoniadis, J. Ellis, J.S. Hagelin, D.V. Nanopoulos, Phys. Lett. B194 (1987), 231.

- 6. S.D in opoulos and F.W ilczek, NSF-ITP-82-07 (unpublished);
  M.Srednicki, Nucl. Phys. B 202 (1982), 327.
- 7. K S. Babu and SM . Barr, Phys. Rev. D 48 (1993), 5354.
- 8. K S. Babu and SM . Barr, Phys. Rev. D 50 (1994), 3529.
- 9. K S.Babu and SM .Barr, Phys. Rev. D 51 (1995) 2463.
- 10. K.S.Babu and R.N.M ohapatra, Phys. Rev. Lett. 74 (1995) 2418.
- 11. K. Inoue, A. Kakuto and H. Takano, Prog. Theor. Phys. 75 (1986), 664.
- 12. A.Anselm and A.Johansen, Phys. Lett. B 200 (1988), 331;
  - A.Anselm, Sov. Phys. JETP 67 (1988), 663.
- 13. R. Barbieri, G. Dvali, A. Strum ia, Nucl. Phys. B 391 (1993), 487.
- 14. Z. Berezhiani and G. Dvali, Sov. Phys. Lebedev Inst. Rep. 5 (1989), 55.
- 15. R. Barbieri, G. Dvali, M. Moretti, Phys. Lett. B 312 (1993), 137.
- 16. R. Barbieri, G. Dvali, A. Strum ia, Z. Berezhiani, L. Hall, Nucl. Phys. B 432 (1994), 49.
- 17. Z.Berezhiani, INFN-FE-14-94, hep-ph/9412372;
  - Z.Berezhiani, INFN-FE-02-95, hep-ph/9503366.
- 18. Z. Berezhiani, C. Csaki, and L. Randall, Nucl. Phys. B 444 (1995) 61.
- 19. C.C saki and L.R and all, to appear
- 20. R. Barbieri, L. Halland A. Strum ia, FUP-TH-72-94, hep-ph/9501334.