

Up-dating for the paper “The mass of the
Higgs boson in the Standard Model from
precision tests” (MPI-Ph/95-39,
hep-ph/9505304). *

Piotr H. Chankowski
Institute of Theoretical Physics, Warsaw University
ul. Hoża 69, 00-681 Warsaw, Poland.

Stefan Pokorski [†]
Max-Planck-Institute für Physik
Werner – Heisenberg – Institute
Föhringer Ring 6, 80805 Munich, Germany

December 23, 2021

Abstract

The bounds on the Higgs mass in the Standard Model are re-analyzed using the precision electroweak data presented at the International Europhysics Conference on High Energy Physics, Brussels, 27 July – 2 August, 1995

*Supported in part by the Polish Committee for Scientific Research and European Union Under contract CHRX-CT92-0004.

[†]On leave of absence from Institute of Theoretical Physics, Warsaw University

After submission of the paper [1] for publication, new data from the SLD and LEP became available. We present here an up-date of our fits with new measurements included. The experimental input used in the present fits is summarized in ref [2]. The bulk of the LEP results show satisfactory agreement with the earlier reports (one should note, however, the new value $A_{FB}^{b,0} = 0.0997 \pm 0.0031$). For M_W we use (as previously) 80.33 ± 0.17 GeV which is the average value of the UA2 measurement and the new measurement reported by the CDF [3] (the D0 collaboration has not published the results of their new analysis yet). When the top quark mass is included in the fit we use the value $m_t = 181 \pm 12$ GeV. For $\Delta\alpha_{EM}^{hadr}$ we use the value ¹ [4] 0.0280 ± 0.0007 and include it into the χ^2 fit.

The most significant changes in the LEP data are $R_b = 0.2219 \pm 0.0017$, and $R_c = 0.1540 \pm 0.0074$. Since the identification of b quarks is much better than of the c quarks experimental collaborations also quote the value $R_b = 0.2206 \pm 0.0016$ which is obtained when the value of R_c is fixed to its SM prediction $R_c = 0.172$.

The new SLD result read [8, 2]: $A_{LR}^e \equiv \mathcal{A}_e = 0.1551 \pm 0.0040$ (which corresponds to $\sin^2 \theta_l^{eff} = 0.23049 \pm 0.00050$). The SLD collaboration also reported for the first time [2] the results $A_{LR}^b \equiv \mathcal{A}_b = 0.841 \pm 0.053$ and $A_{LR}^c \equiv \mathcal{A}_c = 0.606 \pm 0.090$.

The new SLD data for A_{LR}^e and the LEP value still remain more than 2σ apart: although the central values are now much closer to each other, the SLD error has significantly decreased.

The newly reported results for R_b , R_c , A_{LR}^b , A_{LR}^c have drastic effects on SM fits whose quality has significantly decreased. This indicates either large fluctuations in the present data or new physics (or both). Here we adopt the first point of view and assume that the SM is the correct low energy effective theory at the electroweak scale. In this framework we critically re-examine the bounds on the Higgs boson mass in several different versions of the fit.

We begin with the results of a fit to all the LEP, SLAC and Fermilab data described above (i.e. to M_W , $1 - M_W^2/M_Z^2$, Γ_Z , σ_h , \mathcal{A}_e , \mathcal{A}_τ , $\sin^2 \theta_{eff}^{lept} < Q_{FB} >$, \mathcal{R}_l , $A_{FB}^{0,l}$, R_b , R_c , $A_{FB}^{0,b}$, $A_{FB}^{0,c}$, A_{LR}^e , A_{LR}^b , A_{LR}^c , m_t and $\Delta\alpha_{EM}^{hadr}$). They are given in the first line of Table 1 and 1σ and 2σ bounds in the (m_t, M_h) plane are shown in Fig.1. Significant upper bounds on the Higgs boson mass are obtained but the quality of the fit is poor.

Table 1. Results of the fit to all the data [2, 3, 8]. All masses in GeV.

m_t	Δm_t	M_h	ΔM_h	$\alpha_s(M_Z)$	$\Delta\alpha_s(M_Z)$	χ^2	d.o.f
171.0	11.1	93	$^{+189}_{-63}$	0.122	0.005	25.0	14
170.5	10.9	82	$^{+181}_{-55}$	0.122	0.005	21.1	12

¹The result reported in the ref. [5] has been recently updated [7, 2] and is now close to the results reported in [4]. The result of ref. [6] is about 1σ lower than that of [4]; it is, however, based on more theoretical assumptions [2].

We now examine the impact of the observables which give the dominant contribution to χ^2 on the Higgs boson mass bounds. In the second line of Table 1 there are also given the results of a fit without the two new SLAC measurements of A_{LR}^b and A_{LR}^c . The bounds on M_h are slightly stronger and the fit is a bit better.

In the first row of Table 2 we show the results of a fit without R_b , R_c and without all the LR asymmetries measured by the SLD (A_{LR}^e and A_{LR}^b , A_{LR}^c). The χ^2 value is now excellent. The 1σ and 2σ contours in the (m_t, M_h) plane are shown in Fig.2a. The bounds on M_h are very weak and critically depend on the value of m_t measured in Fermilab. Without m_t in the fit the included data only correlate M_h with m_t (as shown also in Fig.2a) with, however, no $2[s]$ upper bound on M_h .

Another observable which depends on m_t but not on M_h is R_b . The measured value, to be consistent with the SM, requires very low value of m_t . Thus, inclusion of m_t and/or R_b gives the fitted value of m_t lower than the Fermilab value and, in consequence, stronger bounds on M_h . This is also shown in Table 2 and in Fig. 2b. Here we give the results of fits with $R_b = 0.2206 \pm 0.0016$ which is the experimental number obtained under the assumption of $R_c = 0.172$ (with the new values of R_b and R_c we get very similar bounds but the quality of the fit is poorer).

Table 2. Results of the fit without the SLD asymmetries.

m_t	R_b	m_t	Δm_t	M_h	ΔM_h	$\alpha_s(M_Z)$	$\Delta\alpha_s(M_Z)$	χ^2	d.o.f
+	−	180.1	11.9	335	$^{+529}_{-222}$	0.125	0.005	3.2	9
−	+	152.2	$^{+14.7}_{-12.9}$	52	$^{+115}_{-29}$	0.123	0.005	9.4	9
+	+	172.4	11.2	189	$^{+326}_{-122}$	0.124	0.005	11.6	10

As the next step we include in the fits the SLD value of A_{LR}^e . Its effect on the fits can be understood from Fig. 2a. We show there the contour of constant $A_{LR}^e = 0.1551$. It is almost parallel to the open contours which show the $m_t - M_H$ correlation without m_t , R_b and A_{LR}^e in the fit but for the same m_t the correlated M_h is much lower. It is clear therefore that to a very good approximation the inclusion of A_{LR}^e should not alter the fitted value of m_t but strengthen the bound on M_h . This is indeed seen in Table 3 and in Fig. 3 (small changes in m_t compared to Table 2 are due to the fact that the two discussed above contours in Fig.2 are not exactly parallel).

Table 3. Results of the fit as in Table 2 but with A_{LR}^e included.

m_t	R_b	m_t	Δm_t	M_h	ΔM_h	$\alpha_s(M_Z)$	$\Delta\alpha_s(M_Z)$	χ^2	d.o.f
+	−	176.7	12.0	125	$^{+271}_{-89}$	0.122	0.005	8.9	10
−	+	155.6	11.1	33	$^{+49}_{-16}$	0.122	0.005	14.8	10
+	+	169.6	10.7	79	$^{+160}_{-54}$	0.122	0.005	17.1	11

The third line in Table 3 can be compared with the second line in Table 1 to see the effect of using new values of R_b and R_c instead of the value of $R_b = 0.2206 \pm 0.0016$ obtained with R_c fixed to its SM value. We conclude that the limits on M_h are stable with respect to the treatment of R_b and R_c in the fit.

To a very good approximation, the upper bounds result from a combination of $m_t - M_h$ correlation given by a fit to all but R_b electroweak data and the upper bound on m_t obtained from R_b and the direct measurement of m_t . The SLD A_{LR}^e result has important impact on $m_t - M_h$ correlation in the direction of lowering the values of M_h for given m_t (as seen in Fig. 3a).

We also note that inclusion of the SLD result in the fits gives lower value of $\alpha_s(M_Z)$. This follows from the fact that the $m_t - M_h$ values required by the value of A_{LR}^e tend to increase Γ_h , i.e. a lower value of $\alpha_s(M_Z)$ is now needed.

Finally one can add a few remarks related to the new values of R_b and R_c . Following earlier references [9] we have discussed in the original version of the paper the possibility of new physics in R_b and studied the upper bounds on M_h in a toy model which reproduces all the results of the SM and has an *ad hoc* correction to the $Z \rightarrow \bar{b}b$ width, so that $R_b \approx 0.22$. Then the fitted value of $\alpha_s(M_Z) = 0.108$ and the observable R_b becomes irrelevant (its contribution to $\chi^2 \approx 0$) from the point of view of the limits on M_h . In this toy model the limits on M_h were similar to the SM limits without R_b in the fit (somewhat weaker due to the smaller value of $\alpha_s(M_Z)$). With the new data the discussion remains valid provided we use the value of R_b extracted under the assumption of R_c fixed to the SM value and disregard the new R_c measurement. If, however, we include both measurements, R_b and R_c , and consider a toy model with new physics in both variables, so that $R_c \approx 0.1540$ and $R_b \approx 0.222$ then the fitted value of $\alpha_s(M_Z) = 0.178 \pm 0.005$. This is easy to understand: with $\Gamma_{Z^0 \rightarrow hadr} = 1744.8 \pm 3.0$ the sum $\Gamma_{Z^0 \rightarrow \bar{b}b} + \Gamma_{Z^0 \rightarrow \bar{c}c}$ is now much smaller than in the SM and we get $\Gamma_{Z^0 \rightarrow hadr} - \Gamma_{Z^0 \rightarrow \bar{b}b} - \Gamma_{Z^0 \rightarrow \bar{c}c}$ large. Of course, also in this toy model the observables R_b and R_c become irrelevant for the bounds on M_h and we get them similar to the SM fit without R_b in the fit. However, the large value of the fitted $\alpha_s(M_Z)$ casts doubts on the measurement of R_c , unless new physics also alters the light quark sector. (We thank H.E. Haber and C. Wagner for the discussion on this point.)

Our results, in part which overlaps, are in very good agreement with recent fits presented in [10].

P.Ch. would like to thank Max Planck Institute in Munich for hospitality.

References

- [1] P.H. Chankowski, S. Pokorski preprint MPI-Ph/95-39, hep-ph/9505304, to be published in *Phys.Lett.* **B**.
- [2] A. Olchevski, plenary talk at the International Europhysics Conference on High Energy Physics, Brussels, 27 July – 2 August, 1995,
D. Charlton, talk at the International Europhysics Conference on High Energy Physics, Brussels, 27 July – 2 August, 1995.
- [3] U. Uwer talk at XXX Rencontres de Moriond, March 1995,
K. Einsweiler talk at the 1995 APS Meeting, Washington, D.C.,
F. Abe et al., the CDF Collaboration, Report FERMILAB-PUB-95-033 (1995), to appear in *Phys. Rev.* **D**.
- [4] S. Eidelman, F. Jegerlehner Paul Scherer Institute preprint, PSI-PR-95-1, January 1995,
H. Burkhard, B. Pietrzyk report LAPP-EXP-95-05.
- [5] M. Swartz report SLAC-PUB-6710.
- [6] A.D. Martin, D. Zeppenfeld *Phys.Lett.* **345B** (1995) 558.
- [7] M. Peskin plenary talk at the SUSY 95 Conference, Palaiseau, 15 – 20 May, 1995.
- [8] K. Abe et al, The SLD Collaboration *Phys. Rev. Lett.* **70** (1993) 2515, **73** (1994) 25,
SLD Collaboration, as presented at CERN by C. Baltay, June 1995.
- [9] A. Blondel, C. Verzegnassi *Phys.Lett.* **311B** (1993) 346,
P. Langacker, J. Erler *Phys. Rev.* **D52** (1995) 441,
Shifman Univ. of Minnesota preprint TPI-MINN-94/42-T (1994).
- [10] J. Ellis, G.L. Fogli, E. Lisi preprint CERN-TH/95-202, BARI-TH/211-95.

FIGURE CAPTIONS

Figure 1.

Contours of $\Delta\chi^2 = 1$ and $\Delta\chi^2 = 4$ for the χ^2 fit to all LEP, SLD and Fermilab data.

Figure 2.

Contours of $\Delta\chi^2 = 1$ and $\Delta\chi^2 = 4$ for the χ^2 fits without the asymmetries measured at SLD (A_{LR}^e , A_{LR}^b , A_{LR}^c):

a) without R_b , R_c ,

b) with $R_b = 0.2206 \pm 0.0016$ ($R_c = 0.172$ fixed) included.

In both figures results of the fits without/with the Fermilab value for m_t included are marked as $1, 2\sigma_{-t}/1, 2\sigma_{+t}$. In the right bottom corner of Fig. 2a the line of constant $A_{LR}^e = 0.1551$ as measured at SLAC is also shown.

Figure 3.

As in Fig. 2 but with the SLD result for A_{LR}^e included.