

Running coupling at small momenta, renormalization schemes and renormalons

N.V. Krasnikov and A.A. Pivovarov

*Institute for Nuclear Research of the Russian Academy of Sciences,
Moscow 117312, Russia*

Abstract

We suggest a method of summing the perturbation theory (PT) asymptotic series related to infrared (IR) renormalons in QCD by using special renormalization schemes in which the running coupling can be integrated over the region of small momenta. For the method to work, one should consider higher order PT corrections to the standard bubble-chain diagrams. High-order corrections allow one to choose a scheme in which the coupling evolution can smoothly be extrapolated to small momenta. In such schemes the sum of an (extended) IR-renormalon asymptotic series is defined as an integral of the running coupling over the IR region. We give explicit examples of renormalization schemes in QCD which can be used for summing IR-renormalon asymptotic series according to our definition.

High-order behavior of perturbation theory series in quantum field theory determines the analytic structure of amplitudes as functions of the coupling constant. This analytic structure encodes important information on the ultimate precision obtainable in the finite-order PT analysis for physical observables. A general approach to analyzing high-order PT expansions is based on the saddle-point approximation for the path integral evaluation [1, 2, 3]. The leading asymptotic behavior of the PT coefficients has been investigated in some models of quantum field theory while there are no exact results in QCD. The high-order behavior can also be studied by selecting large contributions of some particular subclass of diagrams [4, 5]. In QED a set of diagrams generated by inserting an infinite number of one-loop fermionic corrections into the photon propagator (“bubble-chain” diagrams) is usually considered as a relevant one in this context. In other gauge models (for instance, in QCD) the selection of relevant diagrams is more involved due to the necessity to retain gauge invariance. The divergent series [6] emerging from such a type of approximation for the high-order behavior – “bubble chains” or renormalons – have intensively been studied [7, 8, 9]. The ultraviolet (UV) renormalons refer to asymptotic series emerging from the integration over large energies while the infrared (IR) renormalons correspond to integration over small energies. Depending on the model, renormalons lead to alternating (Borel summable) asymptotic series or non-alternating (Borel non-summable) asymptotic series. It is generally assumed that the Borel summation procedure gives the proper value for the asymptotic series which means that the theory can cope with the renormalon singularities that give alternating asymptotic series. The appearance of Borel non-summable series is usually considered as a signal that a theory (or an approximation in which it is studied) may be inconsistent. In QCD the IR renormalons give an example of Borel non-summable series if the leading-order approximation for the QCD β -function with a positive coefficient β_0 (not too large number of flavors) is used.

The recent interest in IR renormalons in QCD has been connected with nonperturbative (nonPT) power corrections [10]. This interest is caused by a limiting accuracy of finite-order PT expressions that cannot presently compete with a high precision of experi-

mental data (e.g. [11]); the accuracy of theoretical formulae has to be improved. A regular way of going beyond a finite-order PT approximation consists in the inclusion of nonPT terms as power corrections to the ordinary PT expressions. For the two-point correlators of interpolating hadronic currents the nonPT terms can be represented phenomenologically as vacuum expectation values of some local operators [12, 13] within the operator product expansion (OPE) which is a general method of quantum field theory [14, 15, 16]. Such an approach has been successful in investigating hadron properties in QCD [17]. However, there is no well-established technique to calculate nonPT contributions in cases when OPE is not applicable. It has been suggested to consider the IR renormalons in QCD (or, generally, Borel non-summable asymptotic series of perturbation theory) as a source for power corrections in such cases.

In QED, the bubble-chain diagrams form a gauge invariant class and give dominant contributions in the framework of $1/N_F$ expansion [18]. In QCD the relevant diagrams are the gluon bubbles which do not form a gauge invariant set. Therefore, to find renormalon contributions in QCD one applies a method of "naive nonabelianization" to the same set of diagrams as in QED [19, 20]. This is equivalent to the substitution of the leading order running coupling constant into integrals over the phase space of the original diagram. Thus, the renormalon approximations are attractive for discussing power corrections from a practical point of view because the relevant diagrams are closely connected with the running of the coupling constant within the renormalization group analysis. The renormalon contributions can be obtained by either a direct summation of the selected set of diagrams or performing the integration of the running coupling constant over the phase space of the leading order diagram which serves as a source for a bubble-chain generation. These ways of obtaining the renormalon contributions (the direct summation of a given subclass of diagrams vs the integration of the running coupling constant) are similar to the methods by which the ultraviolet asymptotics of Green's functions in quantum electrodynamics was originally found. First the direct summation of the leading ultraviolet asymptotics in all orders was performed [21] and later the same results were obtained by

computing the one-loop β -function and solving the renormalization group (RG) equation for the invariant charge [22, 23]. Note that to use the direct summation in high orders of PT is technically difficult while to solve the RG equations with a high-order β -function is feasible. This observation is also valid for renormalons: to analyze corrections to the leading order renormalon expressions within the direct summation approach is technically difficult while the integration of the running coupling constant can practically be done at any order of the approximation for the evolution if the corresponding β -function is known. The technique for calculation of RG functions is well developed [24] and the high-order PT expressions for β -functions are available for a number of models.

In the asymptotically free QCD, the running coupling is regular at large momenta and integrals over large momenta are well defined which solves the problem of summing the UV-renormalon contributions. In the direct summation technique these contributions correspond to alternating asymptotic series which are Borel summable and can be obtained by a formal term-by-term integration of the leading order PT expansion for the running coupling. For determining the IR-renormalon contributions in QCD one should integrate the running coupling over small energies where it is singular in the leading order of PT (Landau pole). Therefore, the integral (and the contribution of IR renormalons) is not defined in the leading order of the coupling evolution. By expanding the running coupling in PT series under the integration sign and term-by-term integration of the resulting series one arrives at a divergent series which is not Borel summable. This corresponds to the direct summation method. Thus, the problem of summing the contributions stemming from the infrared region can be reduced (at least at the formal computational level) to defining an integral of the running coupling over the small momenta. The interpretation of the integral is necessary if there is a singularity in the IR region as in the leading order of PT in QCD where there is a Landau pole. A particular definition of the evolution in the IR region provides a recipe for resummation of the Borel non-summable asymptotic series emerging from the integration over the small momenta. Because the evolution in the IR region is not perturbative the extrapolation of the running to small momenta is not

unique and implies some nonPT assumptions. For instance, the Landau pole appears for the evolution generated by the leading order PT β -function. This singularity is unstable against inclusion of higher order PT corrections which are large in the IR region because it is a domain of strong coupling. Higher order corrections can strongly change the behavior of the running coupling at small energies. For instance, the pole can disappear (infrared fixed point) or the singularity can become an integrable one and the integration will lead to a physically acceptable result. In such cases the integration is possible that defines the way of resummation of the asymptotic series generated by the IR renormalons which means that high-order PT corrections stabilize the theory. An acceptance of a particular scenario for the IR evolution, however, cannot be justified in PT framework only and means that some nonPT assumptions have been made.

In the present paper we define the renormalon contributions using the method of the integration of the running coupling. The relevant representation can be written in a compact form and generalized to higher orders of PT. In the leading order it is equivalent to the "naive nonabelianization" trick. The IR contribution in QCD is not defined in the leading order because of the singularity of the formally extrapolated running coupling to small momenta. However, the method of integration allows one to generalize the renormalon expressions to higher orders of the coupling evolution. In higher orders there is a freedom of choosing renormalization schemes for the coupling and one can always find a scheme in which the evolution trajectory is integrable in the IR region [25]. With an integrable coupling one obtains a particular way of resummation of renormalon-type contributions. In QED the generalization to higher orders means that one inserts the full one-particle irreducible polarization function into the photon propagator. It is close in spirit to organization of the expansion in skeleton diagrams. In QCD, the renormalon chains beyond the leading order can be generated from the running coupling using Dyson-Schwinger equation [26].

Practical applications of the renormalon-type estimates are most frequent in the standard QCD [27]. An important consequence of the bubble-chain based analysis for heavy

quarks is an IR sensitivity of the pole mass [28]. Because the IR region is a nonperturbative one and the results based on the consideration of bubble chains are not rigorous (e.g. [29, 30]) there is no unique quantitative evaluation of such a sensitivity. In high precision analysis of heavy quark properties it is generally assumed that the uncertainty of the heavy quark mass due to the IR sensitivity is numerically small [31, 32].

In the following we explore some possibilities to define an integral of the running coupling over the IR region by using a freedom of renormalization scheme (RS) choice in high-order PT. This provides a particular method of summation of asymptotic series related to the IR renormalons in QCD. For the method to work one should consider higher order PT corrections to the standard bubble-chain diagrams.

First we consider an example of how higher order PT contributions can be used to give a definition of the IR renormalon contributions. Let the β -function of a theory in some particular renormalization scheme be summed into the expression

$$\beta(\alpha) = -\frac{\alpha^2}{1 + k\alpha^2}, \quad k > 0 \quad (1)$$

with the standard asymptotic β -function of the form $\beta^{as}(\alpha) = -\alpha^2$. The renormalization group equation for the running coupling constant $\alpha(z)$

$$z \frac{\partial}{\partial z} \alpha(z) = \beta(\alpha) \quad (2)$$

has a solution

$$\alpha(z) = \frac{-L_z + \sqrt{L_z^2 + 4k}}{2k}, \quad L_z = \ln \frac{z}{\Lambda^2}. \quad (3)$$

This solution has no singularities at positive z . However, the asymptotic solution

$$\alpha^{as}(z) = \frac{1}{\ln(z/\Lambda^2)} \quad (4)$$

has a pole at $z = \Lambda^2$. Thus, the particular way of summing an infinite number of perturbative terms for the β -function in eq. (1) has cured the Landau pole problem in a sense that the coupling becomes smooth in the IR region and the integration of the evolution trajectory can be performed explicitly. Let us stress that there are no nonPT

terms added but the freedom of choosing a renormalization scheme for an infinite series defining the running coupling was used instead. This is, however, beyond the formal framework of perturbation theory where only finite order polynomials in the coupling constant are considered for β -functions. The problem of RS ambiguity in high orders of PT was discussed in refs. [33, 34, 35, 36] where several methods to reduce this ambiguity have been proposed. In contrast to previous approaches where the main criterion for a good scheme is a weak dependence on the change of the scheme parameterization, in the present paper we use the RS freedom to find a scheme that allows one to perform the integration over the infrared region without encountering a singularity.

Eq. (1) can be considered either as a pure PT result in some particular renormalization scheme after an infinite resummation or as an approximation of some exact β -function (as obtained from a numerical study on the lattice, for instance) that might include nonperturbative terms as well. The essential point for us here is that the running coupling obeying the renormalization group equation with such a β -function can smoothly be extrapolated into the IR region. The running coupling given in eq. (3) has a correct asymptotic behavior at $z \rightarrow \infty$. It is a regular expansion parameter for physical observables that obey the dispersion relation because it has no Landau pole on the physical cut.

The generalization of eq. (1) to QCD is now straightforward. Keeping in mind that first two coefficients β_0 and β_1 of the QCD β -function are renormalization scheme invariant one could suggest a possible "improved" β -function as

$$\beta(\alpha) = \frac{-\beta_0\alpha^2 - \beta_1\alpha^3}{1 + k(\beta_0\alpha^2 + \beta_1\alpha^3)}, \quad k > 0 \quad (5)$$

with the running coupling constant $\alpha(z)$ implicitly determined by the equation

$$\beta_0 \ln \frac{z}{\Lambda^2} = \frac{1}{\alpha(z)} - \frac{\beta_1}{\beta_0} \ln \left(\frac{\beta_0 + \beta_1\alpha(z)}{\beta_1\alpha(z)} \right) - k\beta_0\alpha(z) \quad (6)$$

The β -function given in eq. (5) is bounded at large α and eq. (6) has a solution for $\alpha(z)$ that is defined on the positive semi-axis and has no singularities.

The solution given in eq. (6) can be rewritten through an intermediate energy scale μ without using the Λ parameter (the Λ parameter has no special meaning as a position of

the pole of the running coupling). For the sake of simplicity we do that for the leading order expression given in eq. (3) and find

$$\alpha(Q^2) = \frac{2\alpha}{1 + \alpha L_Q - k\alpha^2 + \sqrt{(1 + \alpha L_Q - k\alpha^2)^2 + 4k\alpha^2}}, \quad (7)$$

where $L_Q = \ln(Q^2/\mu^2)$ and $\alpha \equiv \alpha(\mu^2)$. This form of the evolution of the coupling constant is a particular regularization of the Landau singularity. This regularization introduces no imaginary part on the positive semi-axis (cf. ref. [37]). Thus, in the pure PT framework one can eliminate the Landau pole of the coupling using the freedom of the RS choice in high orders. The smooth coupling requires no nonperturbative effects to regularize the integrals in which it appears. The integral over the small momenta determines a particular sum for a series related to an (extended) IR renormalon contribution.

The expression given in eq. (5) for the β -function is not polynomial. Still even for a two-loop β -function there can exist an IR fixed point that allows one to make an extrapolation to small energies. Indeed, for n_f light flavors in QCD the first two coefficients of the β -function read

$$4\pi\beta_0 = 11 - \frac{2}{3}n_f, \quad (4\pi)^2\beta_1 = 102 - \frac{38}{3}n_f$$

and for

$$\frac{153}{19} < n_f < \frac{33}{2}$$

one has

$$\beta_0 > 0, \quad \beta_1 < 0.$$

Therefore, for $n_f = 9$, for instance, there exists an infrared fixed point and an extended IR-renormalon asymptotic series (generated with a two-loop β -function) can be explicitly summed according to our prescription, i.e. in the second order approximation for the evolution of the coupling in QCD.

The coupling constant defined in a particular renormalization scheme is not an immediate physical quantity. However, one can write a formal definition of the effective coupling through some physical quantity. This definition can implicitly include nonperturbative

terms. For instance, taking the expression for Adler function

$$D_{e^+e^-}(Q^2) = Q^2 \int \frac{R_{e^+e^-}(s)ds}{(s+Q^2)^2}$$

one can define

$$d^{as}(Q^2) = (D_{e^+e^-}(Q^2) - 1)\pi\beta_0 = \left(\ln \frac{Q^2}{\Lambda^2}\right)^{-1}$$

and [38, 39]

$$r_{e^+e^-}^{as}(s) = \frac{1}{\pi} \arctan\left(\frac{\pi}{\ln \frac{s}{\Lambda^2}}\right) \Theta(s) + \Theta(s+\Lambda^2)\Theta(-s). \quad (8)$$

The unphysical singularity (a cut along a part of the negative semi-axis) corresponds to the unphysical pole in $D(Q^2)$. Subtracting this singularity in a minimal way we find the expression

$$d(Q^2) = \frac{1}{\ln(Q^2/\Lambda^2)} - \frac{\Lambda^2}{Q^2 - \Lambda^2} \quad (9)$$

that is regular for positive Q^2 but has power corrections at large Q^2 and is not polynomial in terms of the asymptotic charge

$$d^{as}(Q^2) = \left(\ln(Q^2/\Lambda^2)\right)^{-1}.$$

Indeed, for large Q^2 one finds

$$d(Q^2) = d^{as}(Q^2) - \frac{1}{e^{\frac{1}{d^{as}(Q^2)}} - 1} = d^{as}(Q^2) - e^{-\frac{1}{d^{as}(Q^2)}} + \dots \quad (10)$$

The corresponding β -function written in terms of the asymptotic charge $d^{as} \equiv \xi$

$$Q^2 \frac{\partial}{\partial Q^2} d(Q^2) = -\xi^2 + e^{-\frac{1}{\xi}}(1 - e^{-\frac{1}{\xi}})^{-2} \Big|_{\xi=d^{as}(Q^2)} \quad (11)$$

also has an explicit nonperturbative term unlike eq. (5). It is unclear whether the β -function in eq. (11) can be rewritten in terms of the full charge $d(Q^2)$ from eq. (9).

Thus, there are various possibilities to eliminate the leading order pole and, thereby, to give an interpretation to the renormalon series as an integral of the running coupling over the IR region. A pure perturbative possibility means that higher order PT corrections make the evolution well-defined till the origin. This can be achieved with both finite-order and resummed expressions for the β -function. Other possibilities can be nonperturbative

ones when the evolution is made smooth in the IR region by some nonperturbative procedure.

Within the IR-renormalon approach the power corrections emerge as a result of integration of some sub-blocks of a whole diagram over the region of small momenta in the phase space of the diagram. The main generating sub-block in the IR-renormalon approximation is taken to be the running coupling. However, for defining the IR contributions there is no need to know the point-by-point evolution of the coupling in the IR domain. One has to define some integrals only. Finding only integrals is less demanding than determining the point-by-point evolution of the coupling in the IR domain. For phenomenological applications the information about a particular way of extrapolation of the coupling constant evolution into the IR domain can be encoded through just a nonPT parameter describing the leading order nonPT correction. In other words any particular prescription of summing the asymptotic series generated by the (extended) IR-renormalon chains corresponds to a numerical value of this parameter. For instance, the parameter k in eq. (1) describes a way of summing the IR-renormalon asymptotic series by using higher order corrections and changing the renormalization scheme in which the evolution is described. For phenomenological applications, however, it is convenient to retain a contact with the leading order asymptotic charge. By choosing the principal value (PV) prescription for the regularization of integrals of the asymptotic charge over the IR region one obtains a particular way of resumming the IR-renormalon contributions. This way is not unique. The residual freedom is parameterized by a nonPT quantity which is IR sensitive only [40]. The parameterization is written in the form

$$\beta_0 \alpha^{eff}(s) = \beta_0 \alpha_s(s)|_{\text{PV}} + A\Lambda^2 \delta(s) + B\Lambda^4 \delta'(s) + \dots \quad (12)$$

The normalization is chosen such that for large s

$$\alpha_s(s) = \frac{1}{\beta_0 \ln(s/\Lambda^2)}. \quad (13)$$

Eq. (12) is understood in a sense of distributions and $\alpha_s(s)|_{\text{PV}}$ is the asymptotic charge containing the pole regularized by taking the principal value. Such a distribution is a

natural generalization of the classical solution to the renormalization group equation with the asymptotic β -function $\beta^{as}(x) = -\beta_0 x^2$. The generalized solution $\alpha_s(s)|_{\text{PV}}$ is not unique and allows some arbitrariness at the point $s = \Lambda^2$ that should be a series of δ -function and its derivatives. Quantities A, B, \dots parameterize this arbitrariness. We have chosen the support of "nonperturbative terms" at $s = 0$ instead of $s = \Lambda^2$ which suffices for the discussion in the present paper. However, if eq. (12) is integrated with a weight function that is singular in the origin one has to retain the exact location of the support $s = \Lambda^2$ of the "nonperturbative terms". It is necessary, for instance, in case of finding renormalon corrections to the pole mass of a heavy quark. Note that multiplication is not defined for distributions therefore the non-leading PT corrections are supposed to be included in the leading order term in eq. (12) that results in the substitution $\Lambda_{\overline{\text{MS}}} \rightarrow \Lambda_{\text{eff}}$ (e.g. [41, 42, 43]). The integration of expression in eq. (12) over the IR region with some numerical values for the parameters A and B should give the same result as an integration of the solution to eq. (5) with some numerical value for the parameter k . For a while we retain two parameters in eq. (12).

Now we consider several models for $\alpha^{eff}(s)$ and compute numerical values for parameters A and B . In the spirit of the operator product expansion $A = 0$ (there are no gauge invariant operators of dimensionality two) and $B \sim \langle G^2 \rangle$, where $\langle G^2 \rangle$ is the gluon condensate. With the constraint $A = 0$ the parameter B is related to the parameter k from eq. (5). Within such a model one expects that the pattern of extrapolation of the running coupling to the infrared region can be found by comparing the numerical value for the parameter B with that of the gluon condensate. The gluon condensate is connected with the violation of dilatation symmetry which is closely related to dimensional transmutation and appearance of the scale Λ . Note that quark condensates are related to chiral symmetry breaking and we do not consider them within our approach.

Assuming a universality of extrapolation of the running coupling into the IR region we fix the parameter B from comparison with OPE. Taking the process of e^+e^- annihilation where the operator product expansion is applicable we find the relation between the B

parameter and the gluon condensate. From the expression for Adler's function at large Euclidean Q^2

$$D_{e^+e^-}(Q^2) = 1 + \frac{\alpha_s^{e^+e^-}(Q)}{\pi} + \frac{\langle g_s^2 G^2 \rangle}{6Q^4} \quad (14)$$

one obtains the (naive) asymptotic spectral density in the leading order

$$R_{e^+e^-}(s) = 1 + \frac{\alpha_s^{e^+e^-}(s)}{\pi} + \frac{\langle g_s^2 G^2 \rangle}{12} \delta'(s) \quad (15)$$

with

$$\alpha_s^{e^+e^-}(s) = \frac{1}{\beta_0 \ln(s/\Lambda_{e^+e^-}^2)}. \quad (16)$$

Note that π^2 -corrections are ignored (cf eq. (8)). Assuming the interpretation of the asymptotic charge $\alpha_s^{e^+e^-}(s)$ in the above equations as a distribution on the semi-axis $0 < s < \infty$ with PV regularization we have

$$\alpha_s^{eff}(s) = \alpha_s^{e^+e^-}(s)_{PV} + \frac{\pi}{12} \langle g_s^2 G^2 \rangle \delta'(s). \quad (17)$$

This equation fixes a particular extrapolation of the coupling evolution into the IR domain. The chosen extrapolation is represented through the PV regularized asymptotic charge and the gluon condensate contribution which has support only in the IR domain. This extrapolation is coordinated with OPE for the two-point correlator in e^+e^- annihilation. Comparing eq. (17) with eq. (12) one finds for three light quarks

$$16B\Lambda_{e^+e^-}^4 = 3\langle g_s^2 G^2 \rangle$$

that translates into the following expression for the scale parameter $\Lambda_{e^+e^-}$ through the gluon condensate

$$\Lambda_{e^+e^-} = \frac{1}{2} \left(\frac{3\langle g_s^2 G^2 \rangle}{B} \right)^{\frac{1}{4}}.$$

For the standard numerical value $\langle g_s^2 G^2 \rangle = 0.474 \text{ GeV}^4$ we find

$$\Lambda_{e^+e^-} = \frac{546}{B^{\frac{1}{4}}} \text{ MeV}. \quad (18)$$

In e^+e^- annihilation the parameter $\Lambda_{e^+e^-}$ is an effective scale of the process [42, 41]. The use of the effective scale allows one to absorb the next-to-leading order PT correction into

$\alpha_s^{e^+e^-}(s)$. The numerical value of the effective scale is rather sensitive to the process and can characterize the scale where PT is violated and nonPT term are important numerically [43, 44]. To find the relation between $\Lambda_{e^+e^-}$ and the standard parameter $\Lambda_{\overline{\text{MS}}}$ we consider the first perturbative correction to the coupling constant $\alpha_s^{e^+e^-}(s)$ in eq. (14)

$$\frac{\alpha_s^{e^+e^-}(\mu)}{\pi} = \frac{\alpha_{\overline{\text{MS}}}(\mu)}{\pi} + k_1 \left(\frac{\alpha_{\overline{\text{MS}}}(\mu)}{\pi} \right)^2 + O(\alpha_s^3) \quad (19)$$

where

$$k_1 = \frac{365}{24} - 11\zeta(3) + n_f \left(-\frac{11}{12} + \frac{2}{3}\zeta(3) \right).$$

Here $\zeta(z)$ is Riemann ζ -function. For the standard QCD with $n_f = 3$ we find numerically

$$\frac{\alpha_s^{e^+e^-}(\mu)}{\pi} = \frac{\alpha_{\overline{\text{MS}}}(\mu)}{\pi} + 1.64 \left(\frac{\alpha_{\overline{\text{MS}}}(\mu)}{\pi} \right)^2 + O(\alpha_s^3) \quad (20)$$

that leads to

$$\Lambda_{e^+e^-} = 1.44 \Lambda_{\overline{\text{MS}}}. \quad (21)$$

Using eqs. (18,21) one obtains a numerical prediction

$$\Lambda_{\overline{\text{MS}}} = 380 B^{-\frac{1}{4}} \text{ MeV}. \quad (22)$$

For the present experimental value of $\Lambda_{\overline{\text{MS}}}^{\text{exp}}$ extracted from the data on the τ -lepton decay [45]

$$\Lambda_{\overline{\text{MS}}}^{\text{exp}} = 349 \pm 61 \text{ MeV} \quad (23)$$

one finds

$$B = \left(\frac{380}{349 \pm 61} \right)^4 = 0.74 \div 3.03. \quad (24)$$

Below we consider several explicit models of extrapolation of the coupling evolution into the IR region that allow one to determine a numerical value for the parameter B . We find that the numerical value for the parameter B obtained from these models agrees with the result given in eq. (24). For the sake of technical simplicity we do not use sophisticated expressions for running coupling in the IR region stemming from RS redefinition. We

limit ourselves to models with rather a similar pattern of the IR behavior but expressible in simple analytical terms that makes the numerical analysis more transparent.

Model 1.

Switching the interaction off completely at small momenta:

$$\alpha^{eff}(z) = \alpha_s(z)\Theta(z - a\Lambda^2).$$

The system of equations for determining A and B is [46]

$$li(a) + A = 0, \quad li(a^2) - B = 0,$$

where $li(a)$ is a special function

$$li(a) = \int_0^a \frac{dt}{\ln(t)}$$

with the PV (principal value) prescription for the integration across the pole at $t = 1$ for real positive $a > 1$ [47]. The numerical solution to the system of equations with an additional constraint $A = 0$ reads $a = 1.45$, $B = li(2.1) = 1.19$. We retain only three significant figures.●

Model 2.

Freezing the running coupling at small momenta (e.g. [26])

$$\alpha^{eff}(s) = \alpha_s(a)\Theta(a\Lambda^2 - s) + \alpha_s(s)\Theta(s - a\Lambda^2).$$

The system of equations for determining A and B is

$$\frac{a}{\ln(a)} = li(a) + A, \quad \frac{a^2}{2\ln(a)} = li(a^2) - B.$$

The numerical solution with the constraint $A = 0$ reads $a = 3.85$, $B = 2.6$.●

Model 3.

Regularization of the low energy behavior by the minimal subtraction of the pole singularity

$$\beta_0\alpha^{eff}(s) = \frac{1}{\ln(s/\Lambda^2)} - \frac{\Lambda^2}{s - \Lambda^2}.$$

The system of equations to determine the parameters A and B has the form

$$li(a) - \ln(a - 1) = li(a) + A, \quad li(a^2) - a - \ln(a - 1) = li(a^2) - B.$$

The solution with $A = 0$ is $a = 2$, $B = 2$. Note that this model has recently got some special attention [48]. •

Model 4.

Renormalization scheme modification of the β -function and smooth extrapolation of the evolution to small momenta. The model is given by the pattern of running presented in eq. (3). The solution depends on the numerical value for the parameter k entering the expression for the β -function in eq. (1). For $k = 2$ the numerical solution is $a = 3.0$, $B = 3.0$ which is close to that of Models 2,3. For larger k the solution is rather stable. For smaller values of k there may be no solution at all ($k=1$). This shows that the ansatz chosen in eqs. (12,17) restricts the possible extrapolations of the effective coupling into the IR region in Minkowskian domain and can be used to fix the numerical value for the B parameter through the gluon condensate. •

Thus, the models of reasonably soft extrapolation of the running to small momenta give the value for the parameter B that agrees with the estimate given in eq. (24). Assuming that the numerical value for B has been found independently we can find a numerical value for the QCD scale parameter from eq. (22). Using the results of Models 1-3 $B = 1.2 - 2.6$ or $B^{\frac{1}{4}} = 1.05 - 1.27$ one obtains $\Lambda_{\overline{\text{MS}}} = 300 \div 362$ MeV, or $\Lambda_{\overline{\text{MS}}} = 331 \pm 31$ MeV that is in a reasonable agreement with the present data. In general, our result means that the numerical values of the gluon condensate and $\Lambda_{\overline{\text{MS}}}$ are compatible with each other for the smooth continuation of the running coupling into the IR region. The gluon condensate is interpreted as a quantity that corrects the PV-regularized leading-order running coupling at small momenta (in a sense of distributions) to provide a smooth continuation of the evolution into the IR domain.

To conclude, we have exploited the RS freedom in high orders of PT to show that contributions of some infinite subsets of diagrams that are represented as integrals of

the running coupling over the IR region can explicitly be summed in a generalized way through the proper definition of the integral. The lack of any parameter or sensible strict criterion for choosing a particular set of diagrams that dominate high-order behavior beyond the leading order expression for the running coupling allows one to use the RS freedom for interpretation of the IR-renormalon singularities. If in a particular renormalization scheme there are no singularities of the running coupling at small momenta, this scheme provides a particular recipe for the resummation of the asymptotic series related to the IR renormalons. As for phenomenological applications, the models with some extrapolation of the running coupling into the infrared region are used for practical calculation. Our results show that with an extrapolation chosen one cannot freely add the standard contribution of the gluon condensate to take into account nonperturbative effects and "improve" the computation – this contribution must be coordinated with the extrapolation of the coupling.

References

- [1] L.N. Lipatov, Zh. Eksp. Teor. Fiz. 72, 411 (1977).
- [2] A.P. Bukhvostov and L.N. Lipatov, Zh. Eksp. Teor. Fiz. 73, 1658 (1977).
- [3] J. Zinn-Justin, Quantum field theory and critical phenomena, Third edition, Oxford, 1996.
- [4] G.'t Hooft, The Whys of Subnuclear Physics, Proceedings of the 15th Inter.School on Subnuclear Physics, Erice, Sicily, 1977, edited by A.Zichichi (Plenum Press, New York, 1979), p.943.
- [5] B. Lautrup, Phys. Lett. B69, 109 (1977).
- [6] G.H. Hardy, Divergent series, Oxford 1973.
- [7] G. Parisi, Phys. Lett. 76B, 65 (1978); Nucl. Phys. B150, 163 (1979).

- [8] A.H. Mueller, Nucl. Phys. B250, 327 (1985).
- [9] I.I. Balitsky, Phys. Lett. B273, 282 (1991).
- [10] V.I. Zakharov, Nucl. Phys. B385, 452 (1992).
- [11] J.G. Korner, F. Krajewski and A.A. Pivovarov, Eur. Phys. J. C12, 461 (2000);
ibid C14, 123 (2000).
- [12] H.D. Politzer, Nucl. Phys. B117, 397 (1976).
- [13] A. Vainshtein, V. Zakharov, M. Shifman, JETP Lett. 27, 59 (1978).
- [14] K.G. Wilson, Phys. Rev. 179, 1399 (1969).
- [15] F. David, Nucl. Phys. B209, 433 (1982).
- [16] A.A. Pivovarov, A.N. Tavkhelidze and V.F. Tokarev,
Theor. Math. Phys. 60, 765 (1985).
- [17] M.A. Shifman, A.I. Vainshtein, V.I. Zakharov, Nucl. Phys. B147, 385 (1979).
- [18] R. Coquereaux, Phys. Rev. D23, 2276 (1981).
- [19] D.J. Broadhurst and A.G. Grozin, Phys. Rev. D52, 4082 (1995);
M. Beneke, V. Braun, Phys. Lett. B348, 513 (1995);
C.N. Lovett-Turner, C.J. Maxwell, Nucl. Phys. B452, 188 (1995).
- [20] G.P. Korchemsky and G. Sterman, Nucl. Phys. B437, 415 (1995).
- [21] L.D. Landau, A.A. Abrikosov, I.M. Khalatnikov,
Doklad. Akad. Nauk SSSR 95, 773 (1954).
- [22] M. Gell-Mann, F.E. Low, Phys.Rev. 95, 1300 (1954).
- [23] N.N. Bogoliubov, D.V. Shirkov, Doklad. Akad. Nauk SSSR 103, 391 (1955)
Nuovo Cim. 3, 845 (1956).

- [24] A.A. Vladimirov, Teor. Mat. Fiz. 43, 210 (1980).
- [25] N.V. Krasnikov and A.A. Pivovarov, INR-903-95, Oct 1995. 9pp., to be published in Yad. Fiz. [hep-ph/9512213];
INR-925-96, Jul 1996. 8pp. Talk given at 10th International Conference on Problems of Quantum Field Theory, Alushta, Ukraine, 13-17 May 1996. Proceedings of 10th International Conference on Problems of Quantum Field Theory, Eds. D.V.Shirkov, D.I.Kazakov, A.A.Vladimirov, Dubna 1996, p.178 - 183. [hep-ph/9607247];
Mod. Phys. Lett. A11, 835 (1996).
- [26] N.V. Krasnikov and A.A. Pivovarov, Sov. Phys. J. 25, 55 (1982).
- [27] M. Beneke, Phys. Rept. 317, 1 (1999).
- [28] I.I. Bigi, M.A. Shifman, N.G. Uraltsev, A.I. Vainshtein, Phys. Rev. D50, 2234 (1994).
- [29] S.V. Faleev, P.G. Silvestrov, Nucl. Phys. B507, 379 (1997).
- [30] I.M. Suslov, Zh. Eksp. Teor. Fiz. 116, 369 (1999).
- [31] A. Hoang *et al.*, Eur. Phys. J. direct C3, 1 (2000).
- [32] A.A. Penin and A.A. Pivovarov, Phys. Lett. B435, 413 (1998);
Nucl. Phys. B549, 217 (1999); MZ-TH-98-61, Dec 1998. 41pp., to be published in Yad. Fiz. [hep-ph/9904278].
- [33] P.M. Stevenson, Phys. Rev. D23, 2916 (1981).
- [34] J. Brodsky, G.P. Lepage, P.B. Mackenzie, Phys. Rev. D28, 228 (1983).
- [35] A. Dhar and V.Gupta, Phys. Rev. D29, 2822 (1984).
- [36] A.C. Mattingly and P.M. Stevenson, Phys. Rev. D49, 437 (1994).
- [37] G. Grunberg, Phys. Lett. B349, 469 (1995).

- [38] N.V. Krasnikov, A.A. Pivovarov, Phys. Lett. 116B, 168 (1982).
- [39] A.A. Pivovarov, Sov. J. Nucl. Phys. 54, 676 (1991); Z.Phys. C53, 461 (1992); Nuovo Cim. 105A, 813 (1992).
- [40] A.A. Penin and A.A. Pivovarov, Phys. Lett. B357, 427 (1995); *ibid* B367, 342 (1996); *ibid* B401, 294 (1997); Phys. Atom. Nucl. 60, 2056 (1997).
- [41] G. Grunberg, Phys. Lett. B90, 70 (1980).
- [42] N.V. Krasnikov, Nucl. Phys. B192, 497 (1981).
- [43] A.L. Kataev, N.V. Krasnikov and A.A. Pivovarov, Phys. Lett. B107, 115 (1981); Nucl. Phys. B198, 508 (1982); *ibid* B490, 505 (1997).
- [44] A.A. Pivovarov and E.N. Popov, Phys. Lett. B205, 79 (1988); Sov. J. Nucl. Phys. 49, 693 (1989).
- [45] J.G. Körner, F. Krajewski and A.A. Pivovarov, MZ-TH-00-03, Feb 2000. 11pp., Phys. Rev. D, to be published [hep-ph/0002166].
- [46] N.V. Krasnikov, A.A. Pivovarov, N.N. Tavkhelidze, JETP Lett. 36, 333 (1982); Z.Phys. C19, 301 (1983).
- [47] I.S. Gradshteyn, I.M. Ryzhik, Tables of integrals, sums and series. Moscow, 1951.
- [48] D.V. Shirkov and I.L. Solovtsov, Phys. Rev. Lett. 79, 1209 (1997).