

TOKUSHIMA 95-04

(hep-ph/9510269)

October 1995

EW Precision Analysis and the Higgs Mass¹⁾Zenro HIOKI²⁾

Institute of Theoretical Physics, University of Tokushima

Tokushima 770, JAPAN

ABSTRACT

Two topics on the standard electroweak theory are discussed based on its remarkable success in precision analyses. One is a test of structure of the radiative corrections to the weak-boson masses as a further precision analysis. The other is an indirect Higgs-boson search through the radiative corrections to the various quantities measured at LEP.

¹⁾ Invited Talk at the second German-Polish Symposium "New Ideas in the Theory of Fundamental Interactions", Zakopane, Poland, September 11-15, 1995 (to appear in the Proceedings).

²⁾ E-mail: hioki@ias.tokushima-u.ac.jp

x1. Introduction

The discovery of the top quark [1] has completed the fermion world in the three generation scheme. In the framework of the standard (minimal) electroweak theory, we now have only one yet-undiscovered ingredient left: the Higgs boson. Combining this with the fact that the electroweak theory (with the radiative corrections) has been quite successful in precision analyses through LEP, SLC, Tevatron and lots of other experimental information, we find ourselves in a position to proceed to further more detailed studies of this theory.

At this Symposium, I gave a talk about two topics under this circumstance based upon some of our recent works [2, 3]: One is a test of structure of the EW radiative corrections via $W \rightarrow Z$ masses. The other is a Higgs-boson search through the radiative corrections and precision LEP data. The latter has already been a popular subject, and there are a lot of related papers (see [4-6] and references cited therein). I do not mean that we developed some new technique to analyze the data. However, it is quite significant for future experiments to draw any information on the Higgs mass, and I showed some results which Consoli and I obtained lately.

Before stepping into actual discussions, let me briefly describe what we have been studying on these topics. First, EW corrections consist of several parts with different properties, and I examined via α , G_F and $M_{W,Z}$ what would happen if each of them would not exist. For example, there are top-quark corrections which do not decouple, i.e., become larger and larger as m_t increases. Studying them are significant not only because it is a test of the EW theory as a renormalizable field theory but also because the existence of such effects is a characteristic feature of theories in which particle masses are produced through spontaneous symmetry breakdown plus large Yukawa couplings.

Next, on the Higgs search. Stimulated by the first CDF report on the top-quark evidence, Najima and I considered if there is not any problem in the EW

theory. We then found that the Higgs mass needs to be 1.1-1.2 TeV in order for $M_W |_{m_t=174 \text{ GeV}}$ to reproduce the central value of M_W^{exp} , contrary to some other analyses using the LEP/SLC data which prefer a lighter Higgs boson: $m_H < 300 \text{ GeV}$ [4, 6]. At present, it is not that serious since such a lighter Higgs is also allowed if we take into account the size of m_t^{exp} and M_W^{exp} , but this motivated us to analyze the LEP data our own way.

The first subject is discussed in section 2, and the second one is in §3. Section 4 is for brief summary and discussions.

2. Structure of EW Corrections

Within the electroweak theory, the muon-decay width up to the $O(\alpha^2)$ corrections is calculated as

$$\Gamma = \Gamma^{(0)}(\alpha; M_W, M_Z) (1 + 2r); \quad (2.1)$$

where $\Gamma^{(0)}$ is the lowest-order width in terms of the fine-structure constant and the weak-boson masses $M_{W,Z}$, and r is the corrections to the amplitude. As mentioned in §1, r consists of several parts with different properties: the leading-log terms, the non-decoupling top-quark terms $r[m_t]$ and the other terms including the bosonic effects. On the other hand, its experimental data, Γ^{exp} , is usually expressed by the Fermi coupling constant G_F . Therefore, by solving $\Gamma = \Gamma^{\text{exp}}$ on M_W , we get

$$M_W = M_W(\alpha; G_F; M_Z; r); \quad (2.2)$$

This formula, the M_W - M_Z relation, is the main tool of my analyses in this section.¹¹

¹¹Over the past several years, some corrections beyond the one-loop approximation have been computed. They are two-loop top-quark corrections [7] and QCD corrections up to $O(\alpha_s^2)$ for the top-quark loops [8] (see [9] as reviews). As a result, we have now a formula including $O(\alpha_s^2)$ and $O(\alpha_s^2 m_t^4)$ effects. In the following, M_W is always computed by incorporating all of these higher-order terms as well, although I will express the whole corrections with these terms also as r for simplicity.

Let me show first by using this formula how the theory with the full corrections is successful, though it is already a well-known fact. We thereby have

$$M_W^{(0)} = 80.9404 \pm 0.0027 \text{ GeV} \quad \text{and} \quad M_W = 80.36 \pm 0.09 \text{ GeV} \quad (2.3)$$

for $M_Z^{\text{exp}} = 91.1887 \pm 0.0022 \text{ GeV}$ [10], where $M_W^{(0)} = M_W(\alpha; G_F; M_Z; r=0)$ and M_W is for $m_t^{\text{exp}} = 180 \pm 12 \text{ GeV}$ [1], $m_b = 300 \text{ GeV}$ and $\alpha_{\text{QED}}(M_Z) = 0.118$. Concerning the uncertainty of M_W , 0.09 GeV , I have a little overestimated for safety. From these results, we can find that the theory with the corrections is in good agreement with the experimental value $M_W^{\text{exp}} = 80.26 \pm 0.16 \text{ GeV}$ [11], while the tree prediction fails to describe it at about 4.3 (99.998 % C.L.).

We are now ready. First, let us see if taking only $(\ln(m_f/M_Z))$ into account is still a good approximation ("QED \rightarrow improved-Born approximation"), which was shown to be quite successful in [12]. The W mass is calculated within this approximation by putting $r=0$ and replacing with $(M_Z)(=1)$ in Eq.(2.2), where $(M_Z) = 1/(128.92 \pm 0.12)$.¹² The result is

$$M_W[\text{Born}] = 79.964 \pm 0.017 \text{ GeV}; \quad (2.4)$$

which leads to

$$M_W^{\text{exp}} - M_W[\text{Born}] = 0.30 \pm 0.16 \text{ GeV}; \quad (2.5)$$

This means that $M_W[\text{Born}]$ is in disagreement with the data now at 1:9, which corresponds to about 94.3 % C.L.. Although the precision is not yet sufficiently high, it indicates some non-Born terms are needed which give a positive contribution to the W mass. It is noteworthy since the electroweak theory predicts such positive non-Born type corrections unless the Higgs is extremely heavy (beyond TeV scale). Similar analyses were made also in [15].

¹² Recently three papers appeared in which (M_Z) is re-evaluated from the data of the total cross section of $e^+e^- \rightarrow \text{hadrons}$ [13] (their updated results are given in [14]). Here I simply took the average of the maximum and minimum among them.

The next test is on the non-decoupling top-quark effects. Except for the coefficients, their contribution to r is

$$r[m_t] = (m_t - M_Z)^2 + \ln(m_t - M_Z): \quad (2.6)$$

According to my strategy, I computed the W mass by using the following r^0 instead of r in Eq.(2.2):

$$r^0 = r - r[m_t]: \quad (2.7)$$

The resultant W mass is denoted as M_W^0 . The important point is to subtract not only m_t^2 term but also $\ln(m_t - M_Z)$ term, though the latter produces only very small effects as long as m_t is not extremely large. r^0 still includes m_t dependent terms, but no longer diverges for $m_t \rightarrow +1$ thanks to this subtraction. I found that M_W^0 takes the maximum value for the largest m_t and the smallest m . That is, we get an inequality

$$M_W^0 \leq M_W^0[m_t^{\max}; m^{\min}]: \quad (2.8)$$

We can use $m_t^{\exp} = 180 \pm 12$ GeV [1] and $m^{\exp} > 65.1$ GeV [16] in the right-hand side of the above inequality, i.e., $m_t^{\max} = 180 + 12$ GeV and $m^{\min} = 65.1$ GeV, but I first take $m_t^{\max} \rightarrow +1$ and $m^{\min} = 0$ in order to make the result as data-independent as possible. The accompanying uncertainty for M_W^0 is estimated at most to be about 0.03 GeV. We have then

$$M_W^0 < 79.950 (\pm 0.030) \text{ GeV} \quad \text{and} \quad M_W^{\exp} - M_W^0 > 0.31 \pm 0.16 \text{ GeV}; \quad (2.9)$$

which show that M_W^0 is in disagreement with M_W^{\exp} at about 1.9%. This means that 1) the electroweak theory is not able to be consistent with M_W^{\exp} whatever values m_t and m take if $r[m_t]$ would not exist, and 2) the theory with $r[m_t]$ works well, as shown before, for experimentally-allowed m_t and m . Combining them, we can summarize that the latest experimental data of $M_{W,Z}$ demand, independent of m , the existence of the non-decoupling top-quark corrections.

The confidence level of this result becomes higher if we use $m_t^{\text{max}} = 180 + 12 \text{ GeV}$ and $m_t^{\text{min}} = 65.1 \text{ GeV}$:

$$M_W^0 < 79.863 (\pm 0.030) \text{ GeV} \quad \text{and} \quad M_W^{\text{exp}} - M_W^0 > 0.40 \pm 0.16 \text{ GeV} ; \quad (2.10)$$

that is, 2.5 level.

Finally, let us look into the bosonic contribution. It was pointed out in [17] by using various high-energy data that such bosonic electroweak corrections are now required. I studied whether we could observe a similar evidence in the $M_W - M_Z$ relation. For this purpose, we have to compute M_W taking account of only the pure-fermionic corrections $\delta r[f]$. Since $\delta r[f]$ depends on m_t strongly, it is not easy to develop a quantitative analysis of it without knowing m_t . Therefore, I took into account m_t^{exp} from the beginning in this case. I express thus-computed W mass as $M_W[f]$. The result became

$$M_W[f] = 80.48 \pm 0.09 \text{ GeV} ; \quad (2.11)$$

This value is of course independent of the Higgs mass, and leads to

$$M_W[f] - M_W^{\text{exp}} = 0.22 \pm 0.18 \text{ GeV} ; \quad (2.12)$$

which tells us that some non-fermionic contribution is necessary at 1.2 level.

It is of course too early to say from Eq.(2.12) that the bosonic effects were confirmed in the $M_W - M_Z$ relation. Nevertheless, this is an interesting result since we could observe nothing before: Actually, the best information on m_t before the first CDF report (1994) was the bound $m_t^{\text{exp}} > 131 \text{ GeV}$ by D0 [18], but we can thereby get only $M_W[f] > 80.19 (\pm 0.03) \text{ GeV}$ while M_W^{exp} [94] was $80.23 \pm 0.18 \text{ GeV}$ (i.e., $M_W[f] - M_W^{\text{exp}} > -0.04 \pm 0.18 \text{ GeV}$). We will be allowed therefore to conclude that "the bosonic effects are starting to appear in the $M_W - M_Z$ relation thanks to the discovery of the top-quark".

x3. Indirect Higgs Search

Here I wish to discuss what information on the Higgs we can get from precision LEP data. As a matter of fact, it is not that easy to draw its indirect information from existing experimental data since the Higgs mass enters EW radiative corrections only logarithmically at one-loop level [19]. Therefore, at present, one can only hope to separate out the heavy Higgs-mass range (say $m_H = 500-1000$ GeV) from the low mass regime $m_H = 100$ GeV as predicted, for instance, from supersymmetric theories. Such analyses are, however, still very important and indispensable for future experiments at, e.g., LHC/NLC.

For our analysis, we used in [3] the disaggregated data, just as presented by the experimental Collaborations, without taking any average of the various results. This type of analysis is interesting by itself to point out the indications of the various sets of data since even a single measurement, if sufficiently precise, can provide precious information. At the same time, since the LEP data are becoming so precise, before attempting any averaging procedure one should first analyze the various measurements with their errors and check that the distribution of the results fulfills the requirements of Gaussian statistics. Without this preliminary analysis, one may include uncontrolled systematic effects which can sizably affect the global averages.

We first restricted to a fixed value of the top-quark mass $m_t = 180$ GeV. As input data, we used the available, individual results $R_Z, R_{had}, R_V, A_{FB}(\ell)$ and $A_{e\ell}$ from the four Collaborations as quoted in [10], where $R_V = R_{had} = 0$, and $A_V = 2g_V^e g_A^e = (g_V^e)^2 + (g_A^e)^2$ ($\ell = e, \mu, \tau$), and $g_{V,A}^f$ are the vector and axial-vector couplings of ℓ to Z).¹³ The theoretical computations have been performed with the computer code TOPAZ0 [21], and the main results are given in Table 1. There we do not see any specific indication on m_H , but a heavy Higgs seems to be a

¹³We did not consider the LR asymmetry by SLD [20] since it is already known that it demands a very heavy top (around 240 GeV) when the lower bound on m_H is taken into account, or conversely m_H must be much lower than this bound when m_t^{exp} is used within the standard EW theory (see, e.g., Ellis et al. in [4]).

little bit favored by the total χ^2 .

--	--	--	--	--	--	--	--	--	--

Table 1

--	--	--	--	--	--	--	--	--	--

We, however, found some problems in the forward-backward asymmetry as shown below. Let us consider the global averages

$$A_{FB}^{\exp}(e) = 0.0154 \pm 0.0030; \quad (3.1)$$

$$A_{FB}^{\exp}(\mu) = 0.0160 \pm 0.0017; \quad (3.2)$$

$$A_{FB}^{\exp}(\tau) = 0.0209 \pm 0.0024; \quad (3.3)$$

and transform the averages for A_e and A_μ

$$A_e^{\exp} = 0.137 \pm 0.009; \quad A_\mu^{\exp} = 0.140 \pm 0.008 \quad (3.4)$$

into "effective" F-B asymmetries by using

$$A_{FB}^{\text{eff}}(e) = \frac{3}{4}(A_e)^2; \quad A_{FB}^{\text{eff}}(\mu) = \frac{3}{4}A_e A_\mu; \quad (3.5)$$

which hold in the electroweak theory. We find

$$A_{FB}^{\text{eff}}(e) = 0.0141 \pm 0.0019; \quad A_{FB}^{\text{eff}}(\mu) = 0.0144 \pm 0.0018 \quad (3.6)$$

in very good agreement with Eqs.(3.1) and (3.2) but not with Eq.(3.3). Therefore, there might be some problem in the direct measurement of $A_{FB}(\mu)$ since all other measurements are in excellent agreement with each other.

Just to have an idea of the effect, we computed the χ^2 without $A_{FB}^{\exp}(\mu)$. The results are illustrated in Table 2, which should be compared with Table 1. We find that the tendency toward a heavy Higgs becomes stronger and the best values of the χ^2 are obtained for a large value of m_H , just as in the case of the W mass mentioned in x1. It is still not easy to get a definite conclusion from this, but the "bulk" of the LEP data, namely those well consistent with each other, show no preference for a light Higgs boson, to say the least of it.

--	--	--	--	--	--	--	--	--	--

Table 2

--	--	--	--	--	--	--	--	--	--

Finally, to see the m_t -dependence of σ^2 , I show in Tables 3 and 4 the total σ^2 for $m_t = 170, 180$ and 190 GeV including all data or excluding $A_{FB}^{exp}(\)$. By increasing (decreasing) the top-quark mass, a larger (smaller) value of m_t comes to be favored. This is because the leading top and Higgs terms in the radiative corrections have opposite signs to each other. For a heavier top $m_t > 180$ GeV, however, Tables 3 and 4 give rather different information and it becomes crucial to include the problematic data for $A_{FB}^{exp}(\)$ to accommodate $m_t = 100$ GeV.

--	--	--	--	--	--	--	--	--	--

Tables 3 and 4

--	--	--	--	--	--	--	--	--	--

We have no mind to claim that Tables 2 and 4 represent a more faithful representation of the real physical situation than Tables 1 and 3. Most likely, our results suggest only that further improvement in the data taking is necessary for a definitive answer. We may, however, conclude thereby that it is not a good idea to focus on a light-mass region in Higgs searches at future experiments.

x4. Summary and Discussions

Let us briefly summarize and discuss what I have talked. In section 2, I have shown that we can now test not only (1) the whole EW corrections but also their various parts separately: (2) the light-fermion leading-log corrections which lead to the improved-Born approximation, (3) the non-decoupling m_t corrections and (4) the bosonic corrections. Studying corrections (1) is a test of the theory as a renormalizable field theory, while (2)–(4) are more detailed tests.

The improved-Born approximation succeeded to a certain extent, which is related to the fact that the EW theory unifies the weak interaction (with $q^2 \sim M_W^2$ scale) and the electromagnetic interactions (with $q^2 \sim 0$ scale). We, however,

have seen that some non-Born corrections are now starting to appear. Next we observed that the non-decoupling m_t corrections are also required, which gives a strong support to the mechanism that m_t is produced via spontaneous symmetry breakdown plus large Yukawa couplings. Similar way we also tested the bosonic corrections and found some small indication for them.

Based on this excellent success, we are able to explore the remaining unknown area, i.e., the Higgs sector, which I discussed in section 3. Concerning such a Higgs search, there are already a lot of papers. Unfortunately, the m -dependence of one-loop quantities are only logarithmic in the minimal scheme and therefore it is not easy to get any strong restriction on m , but we have so far obtained some quantitative information. Such information is of course extremely important for future experimental projects like LHC/NLC. Several papers pointed out that the Higgs will be rather light, say less than about 300 GeV [4]. We have found, however, there is also an indication for a rather heavy Higgs through our analysis of LEP and W -mass data.

Due to the reason mentioned above, our results cannot be strong either, but at least we can say it is risky to concentrate our attention on a light-mass region in Higgs searches at future experiments, though I am not a fan of a heavy Higgs boson. More precise measurements of the top-quark and W -boson masses are considerably significant for studying this problem (and also for searching any new-physics effects), and I wish to expect that the Tevatron and LEP II will give us a good answer for it in the very near future.

ACKNOWLEDGEMENTS

I am grateful to R. Rączka and the organizing committee of the Symposium for the invitation and their warm hospitality. I also would like to thank M. Consoli and R. Najima for collaboration, on which many parts of this talk is based. During the Symposium, I enjoyed valuable conversation with R. Rączka,

G. Weinberg and many other participants, which I appreciate very much. I must not forget to thank Jan Fischer, without whom I could not have reached the Symposium site on the first day. This work is supported in part by the Grant-in-Aid for Scientific Research (No. 06640401) from the Ministry of Education, Science, Sports and Culture, Japan.

References

- [1] CDF Collaboration : F. Abe et al, Phys. Rev. Letters 73 (1994), 225; Phys. Rev. D 50 (1994), 2966; Phys. Rev. Letters 74 (1995), 2626;
D0 Collaboration : S. Abachi et al, Phys. Rev. Letters 74 (1995), 2632.
- [2] Z. Hioki, Phys. Letters B 340 (1994), 181; Z. Hioki and R. Najima, Mod. Phys. Letters A 10 (1995), 121; Z. Hioki, Int. J. Mod. Phys. A 10 (1995), 3803.
- [3] M. Consoli and Z. Hioki, Mod. Phys. Lett. A 10 (1995), 845; Preprint TOKUSHIMA 95-03 (hep-ph/9505249) (to appear in Mod. Phys. Lett. A).
- [4] J. Ellis, G. L. Fogli and E. Lisi, Phys. Letters B 333 (1994), 118;
G. Montagna, O. Nicrosini, G. Passarino and F. Piccinini, Phys. Letters B 335 (1994), 484;
S. Matsumoto, Preprint KEK-TH-418 (hep-ph/9411388);
P. H. Chankowski and S. Pokorski, Preprint MPI-Ph/95-39 (IFT-95/6 (hep-ph/9505308)).
- [5] P. Langacker, in: Proceedings of 22nd INS International Symposium "Physics with High Energy Colliders", March 8-10, 1994, INS, Univ. Tokyo, Japan, ed. by S. Yamada and T. Ishii (World Scientific, 1995), p.107;
K. Hagiwara, S. Matsumoto, D. Haidt and C. S. Kim, Zeit. für Phys. C 64 (1994), 559;

K .K ang and S .K .K ang, P reprint BROW N -HET -968 (hep-ph/9412368) and BROW N -HET -979 (hep-ph/9503478).

[6] J .Ellis, G .L .Fogli and E .Lisi, P reprint CERN -TH /95-202 { BAR I-TH /211-95 (hep-ph/9507424).

[7] R .Barbieri, M .Beccaria, P .C iafaloni, G .Curci and A .V icere, Nucl. Phys. B 409 (1993), 105;
J .F leischer, O .V .Tarasov and F .Jegerlehner, Phys. Letters B 319 (1993), 249.

[8] F .Halzen and B .A .K niehl, Nucl. Phys. B 353 (1991), 567;
K .G .Chetyrkin, J .H .K uhn and M .Steinhauser, Phys. Letters B 351 (1995), 331;
L .A vdeev, J .F leischer, S .M ikhailov and O .Tarasov, Phys. Letters B 336 (1994), 560;
B .A .K niehl and A .Sirlin, Nucl. Phys. B 371 (1992), 141; Phys. Rev. D 47 (1993), 883;
B .H .Sm ith and M .B .Voloshin, Phys. Rev. D 51 (1995), 5251.

[9] S .Fanchiotti, B .K niehl and A .Sirlin, Phys. Rev. D 48 (1993), 307;
W .Hollik, in: Precision tests of the Standard M odel, A dvanced series on directions in high-energy physics, ed. by Paul Langacker (W orld Scienti c, 1995), p.37;
B .A .K niehl, Int. J. M od. Phys. A 10 (1995), 443.

[10] The LEP Collaborations, CERN report LEPEW W G /95-01;
T .M ori, P reprint UT -ICEPP 95-07.

[11] CDF Collaboration : F .Abe et al., Phys. Rev. Letters 75 (1995), 11;
M .D em arteau, H .Frisch, U .Heintz, R .Keup and D .Saltzberg, Joint CDF note/D 0 note CDF /PHYS/CDF /PUBLIC /2552 and D 0NOTE 2115.

- [12] V . A . Novikov, L . B . O kun and M . I . V ysotsky, M od . P hys . Letters A 8 (1993), 2529;
M . B ilenky, K . K obodziej, M . K uroda and D . Schildknecht, P hys . Letters B 319 (1993), 319;
G . A ltarelli, P reprint CERN -TH .7045/93 (Talk at the EPS Conference on High Energy Physics, M arseille, France, July 1993) .
- [13] M . L . Swartz, P reprint SLAC -PUB -6710 and SLAC -PUB -95-7001 (hep-ph/9411353 and 9509248);
A . D . M artin and D . Zeppenfeld, P hys . Letters B 345 (1995), 558;
S . E idelman and F . Jegerlehner, P SI-P R -95-1 (hep-ph/9502298) .
- [14] T . Takeuchi, P reprint FERM ILAB -CONF -95/173-T (hep-ph/9506444) .
- [15] V . A . Novikov, L . B . O kun, A . N . Rozanov and M . I . V ysotsky, M od . P hys . Letters A 9 (1994), 2641;
K . K ang and S . K . K ang, P reprint BROW N-HET -940 (hep-ph/9403406 and 9509431) .
- [16] A . Sopczak, P reprint CERN -PPE /95-46 (hep-ph/9504300) .
- [17] S . D ittm aier, D . Schildknecht, K . K obodziej and M . K uroda, Nucl . P hys . B 426 (1994), 249;
P . G ambino and A . Sirlin, P hys . Rev . Letters 73 (1994), 621 .
- [18] D 0 Collaboration : S . Abachi et al., P hys . Rev . Letters 72 (1994), 2138 .
- [19] M . V eltm an, A cta P hys . Pol . B 8 (1977), 475; P hys . Letters 70B (1977), 253;
M . E inhom and J . W udka, P hys . Rev . D 39 (1989), 2758; D 47 (1993), 5029 .
- [20] SLD Collaboration : K . Abe et al., P hys . Rev . Letters 73 (1994), 25 .
- [21] G . M ontagna, O . N icrosini, G . Passarino, F . P iccinini and R . P ittau, Nucl . P hys . B 401 (1993), 3; C omput . P hys . C om m un . 76 (1993), 328 .

$\alpha_{\text{QCD}}(M_Z)$	0.113	0.125	0.127	0.130
m (GeV)	100	100	500	1000
ALEPH	11.2	15.2	13.9	14.7
DELPHI	5.1	7.9	6.7	7.2
L3	11.6	6.0	8.0	9.2
OPAL	19.4	13.9	8.5	6.9
Total χ^2	47.3	43.0	37.1	38.1

Table 1. Total χ^2 for the four Collaborations.

$\alpha_{\text{QCD}}(M_Z)$	0.113	0.125	0.127	0.130
m (GeV)	100	100	500	1000
ALEPH	10.2	14.3	12.1	12.5
DELPHI	4.7	7.5	5.9	6.2
L3	8.4	2.8	3.9	4.8
OPAL	19.4	13.8	8.0	6.1
Total χ^2	42.7	38.4	29.9	29.6

Table 2. Total χ^2 for the four Collaborations by excluding the data for $A_{\text{FB}}^{\text{exp}}(\)$.

ALEPH+DELPHI+L3+OPAL

$\alpha_{\text{QCD}}(M_Z)$	0.113	0.125	0.127	0.130
m_t (GeV)	100	100	500	1000
m_t (GeV)= 170	46.3	38.4	38.3	41.2
= 180	47.3	43.0	37.1	38.1
= 190	51.8	50.4	38.9	37.5

Table 3. Total χ^2 for the four Collaborations at various values of m_t .

ALEPH+DELPHI+L3+OPAL

$\alpha_{\text{QCD}}(M_Z)$	0.113	0.125	0.127	0.130
m_t (GeV)	100	100	500	1000
m_t (GeV)= 170	40.7	32.8	29.7	31.2
= 180	42.7	38.4	29.9	29.6
= 190	50.1	46.6	32.9	30.3

Table 4. Total χ^2 for the four Collaborations at various values of m_t by excluding the data for $A_{\text{FB}}^{\text{exp}}(\)$.