

October 23, 1995

UMN-TH-1412/95

# Massive multi-flavor Schwinger model at finite temperature and on compact space<sup>1</sup>

YUTAKA HOSOTANI

*School of Physics and Astronomy, University of Minnesota  
Minneapolis, Minnesota 55455, U.S.A.  
yutaka@mnhep.hep.umn.edu*

The multi-flavor Schwinger model on  $R^1$  at finite temperature  $T$  is mathematically equivalent to the model on  $S^1$  at  $T = 0$ . The latter is reduced to a quantum mechanical system of  $N - 1$  degrees of freedom. Physics sensitively depends on the parameter  $m/T$ . Finite temperature behavior of the massive Schwinger model is quite different from that of the massless Schwinger model.

## 1. Introduction

The Schwinger model is QED in two dimensions, described by

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \sum_{a=1}^N \bar{\psi}_a \left\{ \gamma^\mu (i\partial_\mu - eA_\mu) - m_a \right\} \psi_a . \quad (1)$$

Massless theory ( $m_a = 0$ ) is exactly solvable. In the  $N = 1$  (one flavor) model the gauge boson acquires a mass without breaking the gauge invariance.<sup>1</sup> It has the  $\theta$  vacuum and a non-vanishing chiral condensate  $\langle \bar{\psi}\psi \rangle \neq 0$ .<sup>2</sup>

The  $N \geq 2$  model is distinctively different from the  $N = 1$  model. The spectrum contains  $N - 1$  massless bosons. The chiral condensate vanishes,  $\langle \bar{\psi}\psi \rangle = 0$ ,<sup>3</sup> as in two dimensions continuous symmetry,  $SU(N)$  chiral symmetry in this case, cannot be spontaneously broken. Nonvanishing  $\langle \bar{\psi}\psi \rangle$  in the  $N=1$  theory is allowed, since the  $U(1)$  chiral symmetry is broken by an anomaly.

With massive fermions the model is not exactly solvable. The effect of the fermion mass in the  $N = 1$  theory is minor for  $m/e \ll 1$ , except that it necessitates the  $\theta$  vacuum. For  $N \geq 2$  the situation is quite different. Coleman showed<sup>4</sup> that in the  $N = 2$  model with  $m_a = m$  two resultant bosons acquire masses given by

$$\mu_1 \sim \frac{\sqrt{2}e}{\sqrt{\pi}} , \quad \mu_2 \propto m^{2/3} e^{1/3} \left| \cos \frac{1}{2}\theta \right|^{2/3} . \quad (2)$$

Surprising is the fractional power dependence on  $m$ ,  $e$  and  $|\cos \frac{1}{2}\theta|$  of the second boson mass  $\mu_2$ , resulting from the self-consistent re-alignment of the vacuum against fermion masses. The effect of fermion masses is always non-perturbative even if  $m/e \ll 1$ .

---

<sup>1</sup>To appear in the Proceedings of the 4th International Workshop on Thermal Field Theory and their Application, Dalian P.R. China, August 5 - 10, 1995.

How does  $\mu_2$  depend on  $m$  in the  $N$  flavor model? How does it change at finite temperature? Does the chiral condensate vanish at sufficiently high temperature? Can the fermion mass be treated as a small perturbation at high temperature?

These are questions addressed in this article. We present a powerful method to evaluate various physical quantities at zero and finite temperature. We shall recognize the importance of a dimensionless parameter  $m/T$ . The behavior at  $m/T \gg 1$  is quite different from that at  $m/T \ll 1$ . Coleman's result (2) corresponds to  $m/T \gg 1$  as  $T \rightarrow 0$ . At high temperature  $m/T \ll 1$ , one finds  $\mu_2 \propto m$ , a result obtained in perturbation theory. This work is based on the result obtained in ref. 5.

## 2. At finite temperature and on a circle

In Matsubara's formalism the model at finite temperature  $T$  in equilibrium is equivalent to an Euclidean field theory, or a theory with an imaginary time  $\tau$ , satisfying boundary conditions

$$\psi_a(\tau + \frac{1}{T}, x) = -\psi_a(\tau, x) \quad , \quad A_\mu(\tau + \frac{1}{T}, x) = A_\mu(\tau, x) \quad . \quad (3)$$

If one, instead, places the model on a circle of circumference  $L$  (at zero temperature) with boundary conditions

$$\psi_a(t, x + L) = -\psi_a(t, x) \quad , \quad A_\mu(t, x + L) = A_\mu(t, x) \quad , \quad (4)$$

then one obtains a theory which is, after Wick's rotation, mathematically equivalent to the finite temperature field theory defined by (3). Various physical quantities in the Schwinger model at  $T \neq 0$  are related to corresponding ones in the model on  $S^1$  by substitution of  $T$  by  $L^{-1}$ .

Our strategy is to solve the model on a circle  $S^1$  with an arbitrary size  $L$ . There is powerful machinery which specifically works on  $S^1$ .<sup>5-8</sup>

## 3. Reduction to a quantum mechanical system

Fermion operators on  $S^1$  can be expressed in terms of bosonic operators: Take  $\gamma^\mu = (\sigma_1, i\sigma_2)$  and write  $\psi_a^T = (\psi_+^a, \psi_-^a)$ . In the interaction picture defined by free massless fermions

$$\psi_\pm^a(t, x) = \frac{1}{\sqrt{L}} C_\pm^a e^{\pm i\{q_\pm^a + 2\pi p_\pm^a(t \pm x)/L\}} : e^{\pm i\sqrt{4\pi}\phi_\pm^a(t, x)} : \quad (5)$$

$$e^{2\pi i p_\pm^a} |\text{phys}\rangle = |\text{phys}\rangle \quad . \quad (6)$$

The Klein factors are given by  $C_+^a = e^{i\pi \sum_{b=1}^{a-1} (p_+^b + p_-^b)}$  and  $C_-^a = e^{i\pi \sum_{b=1}^a (p_+^b - p_-^b)}$ . Here  $[q_\pm^a, p_\pm^b] = i\delta^{ab}$  and  $\phi_\pm^a(t, x) = \sum_{n=1}^\infty (4\pi n)^{-1/2} \{c_{\pm, n}^a e^{-2\pi i n(t \pm x)/L} + \text{h.c.}\}$  where  $[c_{\pm, n}^a, c_{\pm, m}^{b\dagger}] = \delta^{ab}\delta_{nm}$ . The  $:$   $:$  indicates normal ordering with respect to  $(c_n, c_n^\dagger)$ . The antiperiodic boundary condition is ensured by a physical state condition (6). In physical states  $p_\pm^a$  takes integer eigenvalues.

After substituting the bosonization formula (5), the total Hamiltonian in the Schrödinger picture becomes

$$\begin{aligned}
H_{\text{tot}} &= H_0 + H_\phi + H_{\text{mass}} \\
H_0 &= -\frac{e^2 L}{2} \frac{d^2}{d\Theta_W^2} + \frac{\pi}{2L} \sum_{a=1}^N \left\{ (p_+^a - p_-^a)^2 + (p_+^a + p_-^a + \frac{\Theta_W}{\pi})^2 \right\} \\
H_\phi &= \int_0^L dx \frac{1}{2} \left[ \sum_{a=1}^N \left\{ \Pi_a^2 + (\phi'_a)^2 \right\} + \frac{e^2}{\pi} \left( \sum_{a=1}^N \phi_a \right)^2 \right]
\end{aligned} \tag{7}$$

Here  $p_+^a - p_-^a$  and  $p_+^a + p_-^a$  correspond to the charge and chiral charge operators, respectively.  $\Theta_W$  is the phase of the Wilson line around the circle, the only physical degree of freedom of the gauge field on the circle,  $A_1 = \Theta_W(t)/eL$ . The coupling between  $p_\pm^a$  and  $\Theta_W$  is induced through the chiral anomaly.<sup>7</sup>  $\phi_a = \phi_+^a + \phi_-^a$  and  $\Pi_a$  is its canonical conjugate.  $H_{\text{mass}}$  is the fermion mass term.

Notice that (7) is an exact operator identity. In the absence of fermion masses  $H_{\text{tot}} = H_0 + H_\phi$ . The zero modes  $(\Theta_W, q_\pm^a)$  decouple from the oscillatory modes  $\phi_a$ , and the Hamiltonian is exactly solvable. The spectrum contains one massive field  $N^{-1/2} \sum_{a=1}^N \phi_a$  with a mass  $\mu = (N/\pi)^{1/2} e$ , and  $N - 1$  massless fields.

To examine effects of  $H_{\text{mass}}$ , first note that  $H_{\text{mass}}$ , and therefore  $H_{\text{tot}}$ , commutes with  $p_+^a - p_-^a$ . Hence we can restrict ourselves to states with  $(p_+^a - p_-^a) |\text{phys}\rangle = 0$ . With this restriction a complete set of eigenfunctions and eigenvalues of  $H_0$  is

$$\begin{aligned}
\Phi_s^{(n_1, \dots, n_N)} &= \frac{1}{(2\pi)^N} u_s \left[ \Theta_W + \frac{2\pi}{N} \sum_a n_a \right] e^{i \sum_a n_a (q_+^a + q_-^a)} \\
E_s^{(n_1, \dots, n_N)} &= \mu \left( s + \frac{1}{2} \right) + \frac{2\pi}{L} \sum_{a=1}^N n_a^2 - \frac{2\pi}{NL} \left\{ \sum_{a=1}^N n_a \right\}^2
\end{aligned} \tag{8}$$

where a harmonic oscillator wave function  $u_s$  satisfies  $\frac{1}{2}(-\partial_x^2 + x^2)u_s = (s + \frac{1}{2})u_s$  with  $x = (\pi e^2 L^2/N)^{-1/4} \Theta_W$ . The ground states of  $H_0$  are infinitely degenerate for  $n_1 = \dots = n_N$  due to the invariance under a large gauge transformation  $\Theta_W \rightarrow \Theta_W + 2\pi$  and  $\psi_a \rightarrow e^{2\pi i x/L} \psi_a$ .

$H_{\text{mass}}$  induces transitions among  $\Phi_s^{(n_1, \dots, n_N)}$ 's. It also gives finite masses to the  $N - 1$  previously massless fields. The structure of the vacuum sensitively depends on  $H_{\text{mass}}$ . The effect of  $H_{\text{mass}}$  turns out quite nonperturbative so long as  $m_a \neq 0$ .

It is more convenient to work in a coherent state basis given by

$$\Phi_s(\varphi_a; \theta) = \frac{1}{(2\pi)^{N/2}} \sum_{\{n, r_a\}} e^{in\theta + i \sum_{a=1}^{N-1} r_a \varphi_a} \Phi_s^{(n+r_1, \dots, n+r_{N-1}, n)} \tag{9}$$

Transitions in the  $s$  index may be ignored to a very good approximation. We seek the vacuum in the form

$$\Phi_{\text{vac}}(\theta) = \int_0^{2\pi} d\varphi_1 \cdots d\varphi_{N-1} f(\varphi_a; \theta) \Phi_0(\varphi_a; \theta). \tag{10}$$

Let  $\chi_\alpha = U_{\alpha a} \phi_a$  and  $\mu_\alpha$  be a mass eigenstate field and its mass. Then matrix elements of  $H_{\text{mass}}$  in the coherent state basis are

$$\begin{aligned} \langle \Phi_0(\varphi'_a; \theta') | H_{\text{mass}} | \Phi_0(\varphi_a \theta) \rangle &= -\delta_{2\pi}(\theta' - \theta) \prod_{b=1}^{N-1} \delta_{2\pi}(\varphi'_b - \varphi_b) \sum_{a=1}^N A_a \cos \varphi_a \\ A_a &= 2m_a e^{-\pi/N\mu L} \prod_{\alpha=1}^N B(\mu_\alpha L)^{(U_{\alpha a})^2} \quad , \quad \varphi_N = \theta - \sum_{a=1}^{N-1} \varphi_a \quad . \end{aligned} \quad (11)$$

$B(z)$  is given by

$$\begin{aligned} B(z) &= \exp \left\{ \sum_{n=1}^{\infty} \left( \frac{1}{n} - \frac{1}{\sqrt{n^2 + (z/2\pi)^2}} \right) \right\} \\ &= \frac{z}{4\pi} \exp \left\{ \gamma + \frac{\pi}{z} - 2 \int_1^{\infty} \frac{du}{(e^{uz} - 1)\sqrt{u^2 - 1}} \right\} . \end{aligned} \quad (12)$$

The eigenvalue equation  $(H_0 + H_{\text{mass}}) \Phi_{\text{vac}}(\theta) = E \Phi_{\text{vac}}(\theta)$  becomes

$$\begin{aligned} \left\{ -\Delta_N^\varphi + V_N(\varphi) \right\} f(\varphi) &= \epsilon f(\varphi) \\ \Delta_N^\varphi &= \sum_{a=1}^{N-1} \frac{\partial^2}{\partial \varphi_a^2} - \frac{2}{N-1} \sum_{a < b}^{N-1} \frac{\partial^2}{\partial \varphi_a \partial \varphi_b} \\ V_N(\varphi) &= -\frac{NL}{2(N-1)\pi} \sum_{a=1}^N A_a \cos \varphi_a \quad . \end{aligned} \quad (13)$$

Here  $\epsilon = NEL/2\pi(N-1)$ . Eq. (13) is nothing but the Schrödinger equation with the kinetic and potential terms given by  $-\Delta_N^\varphi$  and  $V_N(\varphi)$ , respectively.

The potential  $V_N(\varphi)$  depends, through  $A_\alpha$  defined in (11), on  $\mu_\alpha$  and  $U_{\alpha a}$  which are to be self-consistently determined from the ground state wave function  $f(\varphi_a; \theta)$  of the Schrödinger equation (13).  $\mu_\alpha$ 's and  $U_{\alpha a}$ 's are determined by

$$\begin{aligned} \frac{\mu^2}{N} \begin{pmatrix} 1 & \cdots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \cdots & 1 \end{pmatrix} + \begin{pmatrix} R_1 & & \\ & \ddots & \\ & & R_N \end{pmatrix} &= U^T \begin{pmatrix} \mu_1^2 & & \\ & \ddots & \\ & & \mu_N^2 \end{pmatrix} U \\ R_a &= \frac{4\pi}{L} A_a \langle \cos \varphi_a \rangle_f = -4\pi m_a \langle \bar{\psi}_a \psi_a \rangle_\theta \quad . \end{aligned} \quad (14)$$

We have denoted  $\langle g(\varphi) \rangle_f = \int [d\varphi] g(\varphi) |f(\varphi)|^2$ . We need to solve (11), (13), and (14) self-consistently.

We have shown that the  $N$  flavor massive Schwinger model is reduced to quantum mechanics of  $N-1$  degrees of freedom in which the potential has to be fixed self-consistently with its ground state wave function.




Figure 1: Chiral condensate  $-\langle\bar{\psi}\psi\rangle/\mu$  and  $\kappa_0$  as a function of temperature  $T$  at  $\theta = 0$  in the  $N = 1$  and  $N = 3$  models. In the  $N = 3$  model  $m/\mu = 10^{-1}, 10^{-2}, 10^{-3}$  and  $10^{-4}$ . The crossover in  $\langle\bar{\psi}\psi\rangle/\mu$  takes place when  $\kappa_0 \sim 0.2$ .

#### 4. N=1 (one flavor)

One flavor case is special.  $\varphi_1 = \theta$  and there is only one massive boson with a mass  $\mu_1$ . This case was analysed in detail in refs. 7 and 9.

The vacuum on  $S^1$  is  $\Phi_{\text{vac}}(\theta) = \Phi_0(\theta)$ . Converting all expressions to finite temperature case, we find the boson mass and chiral condensate to be

$$\begin{aligned}\mu_1^2 &= \mu^2 + 8\pi m T \cos \theta e^{-\pi T/\mu} B\left(\frac{\mu_1}{T}\right) \\ \langle\bar{\psi}\psi\rangle_\theta &= -2T \cos \theta e^{-\pi T/\mu} B\left(\frac{\mu_1}{T}\right)\end{aligned}\tag{15}$$

where  $\mu = e/\sqrt{\pi}$ . At  $T = 0$

$$\begin{aligned}\mu_1 &= \sqrt{\mu^2 + m^2 e^{2\gamma} \cos^2 \theta} + m e^\gamma \cos \theta \\ \langle\bar{\psi}\psi\rangle_\theta &= -\frac{e^\gamma}{2\pi} \mu_1 \cos \theta.\end{aligned}\tag{16}$$

Notice that the condensate is nonvanishing even for  $m = 0$ . As  $T$  is raised, it shows a crossover around  $T = \mu$ , and approaches zero at high temperature. See fig. 1. The correction due to the fermion mass  $m \ll \mu$  is minor.

## 5. $N \geq 2$ (multi-flavors) at low and high $T$

The situation in the multi-flavor case is quite different. When  $m_a = 0$ ,  $V_N(\varphi) = 0$  and  $f(\varphi) = \text{const}$  so that  $\langle \cos \varphi_a \rangle_f = 0$ . Consequently  $\langle \bar{\psi}_a \psi_a \rangle_\theta = 0$ .

Suppose that fermion masses are degenerate:  $m_a = m$ . There results one heavy boson and  $N - 1$  light bosons with masses  $\mu_1$  and  $\mu_2$ , respectively. The potential in (13) becomes

$$V_N(\varphi) = -\kappa_0 \sum_{a=1}^N \cos \varphi_a$$

$$\kappa_0 = \frac{N}{(N-1)\pi} \frac{m}{T} B\left(\frac{\mu_1}{T}\right)^{\frac{1}{N}} B\left(\frac{\mu_2}{T}\right)^{1-\frac{1}{N}} e^{-\pi T/N\mu} \quad (17)$$

where  $\mu_1$  and  $\mu_2$  are determined by

$$\mu_1^2 = \mu^2 + \mu_2^2$$

$$\mu_2^2 = \frac{8\pi^2(N-1)}{N} \kappa_0 T^2 \langle \cos \varphi \rangle_f = -4\pi m \langle \bar{\psi}_a \psi_a \rangle_\theta \quad (18)$$

Recognize that two parameters  $\kappa_0$  and  $\theta$  fix the potential  $V_N(\varphi)$ .

If  $m \neq 0$ ,  $\kappa_0$  becomes very large at low temperature  $T \rightarrow 0$ . In this regime the potential term dominates over the kinetic energy term in the Schrödinger equation (13). The wave function  $f(\varphi)$  has a sharp peak around the minimum of the potential. The minimum is located at  $\varphi_a = \bar{\theta}/N$  where  $\bar{\theta} = \theta - 2\pi[(\theta/2\pi) + \frac{1}{2}]$ . As  $\theta$  varies from  $-\pi$  to  $+\pi$ , the minimum moves from  $\varphi_a = -\pi/N$  to  $\varphi_a = +\pi/N$ , and jumps back to  $\varphi_a = -\pi/N$ .

In the  $T \rightarrow 0$  limit,  $\langle \cos \varphi \rangle_f = \cos \bar{\theta}$  so that

$$\frac{1}{\mu} \langle \bar{\psi} \psi \rangle_\theta = -\frac{1}{4\pi} \left( 2e^\gamma \cos \frac{\bar{\theta}}{N} \right)^{\frac{2N}{N+1}} \left( \frac{m}{\mu} \right)^{\frac{N-1}{N+1}} \quad \text{for } T \ll m^{\frac{N}{N+1}} \mu^{\frac{1}{N+1}} \quad (19)$$

Two important observations follow. First, the dependence of the condensate on  $m$  is non-analytic. It has fractional power dependence. The effect of fermion masses is nonperturbative in this limit. Secondly, as a function of  $\theta$ , the condensate has a cusp at  $\theta = \pm\pi$ , which originates from the discontinuous jump in the location of the minimum of the potential.

In the opposite limit  $\kappa_0 \ll 1$ , which includes both  $m \rightarrow 0$  (with  $T > 0$  kept fixed) and  $T \gg \mu$ , the potential  $V_N(\varphi)$  can be treated as a small perturbation in (13). One finds

$$\langle \cos \varphi_a \rangle_f = \begin{cases} (1 + \cos \theta) \kappa_0 & \text{for } N = 2 \\ \kappa_0 & \text{for } N \geq 3. \end{cases} \quad (20)$$




Figure 2: Wave function  $|f(\varphi)|^2$  in the  $N = 3$  model.

There appears no  $\theta$ -dependence for  $N \geq 3$  to this order. The condensate for  $N \geq 3$  is found to be

$$\frac{1}{\mu} \langle \bar{\psi} \psi \rangle_{\theta} = -\frac{2N}{\pi(N-1)} \frac{m}{\mu} \begin{cases} \left( \frac{\mu e^{\gamma}}{4\pi T} \right)^{2/N} & \text{for } m^{\frac{N}{N+1}} \mu^{\frac{1}{N+1}} \ll T \ll \mu \\ e^{-2\pi T/N\mu} & \text{for } T \gg \mu \end{cases} \quad (21)$$

For  $N = 2$ , the expressions for  $\langle \bar{\psi} \psi \rangle_{\theta}$  in (21) must be multiplied by a factor  $2 \cos^2 \frac{1}{2} \theta$ . Notice that the condensate is linearly proportional to  $m$  in this regime.

## 6. Crossover in $\langle \bar{\psi} \psi \rangle_{\theta, T}$

For general values of  $T$  and  $m$ , we need to solve the set of equations numerically. In the two flavor case the Schrödinger equation is

$$\left\{ -\frac{\partial^2}{\partial \varphi_1^2} - 2\kappa_0 \cos \frac{1}{2} \theta \cos(\varphi_1 - \frac{1}{2} \theta) \right\} f(\varphi_1) = \epsilon f(\varphi_1) . \quad (22)$$

This is the equation for a quantum pendulum. The strength of the potential is  $2\kappa_0 \cos \frac{1}{2} \theta$ , which changes the sign at  $\theta = \pm \pi$ . In particular, the potential vanishes at  $\theta = \pm \pi$  and  $f(\varphi_1) = \text{constant}$ . Consequently the chiral condensate vanishes at  $\theta = \pm \pi$ . This is a special feature of  $N = 2$ .

For  $N = 3$  the equation is

$$\left\{ -\frac{\partial^2}{\partial \varphi_1^2} - \frac{\partial^2}{\partial \varphi_2^2} + \frac{\partial^2}{\partial \varphi_1 \partial \varphi_2} - \kappa_0 [\cos \varphi_1 + \cos \varphi_2 + \cos(\varphi_1 + \varphi_2 - \theta)] \right\} f(\varphi) = \epsilon f(\varphi) . \quad (23)$$




Figure 3: The location of the minimum of the potential in the  $N = 3$  model is displayed with various values of  $(q, r)$  in (23). For  $(q, r) = (10, 1.3)$  the minimum at  $\theta = \pm\pi$  is located at the origin.

The potential term never vanishes unless  $\kappa_0 = 0$ , or equivalently  $m = 0$  or  $T \rightarrow \infty$ . The equation can be solved numerically for an arbitrary  $\kappa_0$ . The ground state wave function  $|f(\varphi)|^2$  has been displayed for various  $\kappa_0$  and  $\theta$  in fig. 2.

With given values of  $m$  and  $T$  the chiral condensate is determined by solving (17), (18), and (23) simultaneously. We developed an iteration procedure which yields a consistent set of values of  $m/\mu$ ,  $T/\mu$ ,  $\mu_a/\mu$ , and  $\kappa_0$ . It takes less than ten iterations even for moderate values of  $\kappa_0 \sim 1$  to achieve four digits accuracy.

In the top figure of fig. 1 we have displayed the condensate  $\langle \bar{\psi}\psi \rangle_{\theta=0}/\mu$  as a function of  $T/\mu$  and  $m/\mu$  in the  $N=3$  case too. With a given  $m/\mu$  the condensate is almost constant at low  $T/\mu$ , and sharply drops to a small value around  $T_*$ . This crossover takes place when  $\kappa_0(m, T_*) \sim 0.2$  for a wide range of the value of  $m/\mu$ . In the bottom figure we have shown a plot for  $\kappa_0$  in the same region of the  $m$ - $T$  space.

The asymptotic formulas (19) and (21) are quite accurate for  $\kappa_0 > 1$  and  $\kappa_0 < 0.1$ . The important parameter is  $\kappa_0$ .

## 7. The singularity at $\theta = \pm\pi$ and fermion masses

As shown in (19), the chiral condensate at  $T = 0$  shows a cusp singularity in its  $\theta$  dependence when fermion masses are degenerate. When fermion masses are not degenerate, the potential  $V_N(\varphi)$  in (13) is deformed from the symmetric one in (17). Accordingly the location of the minimum is shifted. The cusp singularity appears when the location of the minimum of the potential makes a discontinuous jump.

With given  $\{m_a\}$ ,  $A_a$  in  $V_N(\varphi)$  is determined self-consistently. We have determined the location of the minimum of the potential in the three flavor case. The potential is



proportional to

$$F(\varphi_1, \varphi_2; \theta) = -q \cos \varphi_1 - r \cos \varphi_2 - \cos(\theta - \varphi_1 - \varphi_2) \quad (24)$$

where  $q = A_1/A_3$  and  $r = A_2/A_3$ .

In the symmetric case  $q=r=1$  the location of the minimum moves from  $(-\frac{1}{3}\pi, -\frac{1}{3}\pi)$  to  $(+\frac{1}{3}\pi, +\frac{1}{3}\pi)$  as  $\theta$  varies from  $-\pi$  to  $+\pi$ , and makes a jump. At  $\theta=0$  the minimum is located at the origin for arbitrary  $(q, r)$ . We have plotted trajectories of the location of the minimum for several typical values of  $(q, r)$  in fig. 3.

So long as asymmetry is small, there is a discontinuous jump at  $\theta = \pm\pi$ . However, a sufficiently large asymmetry restores continuity. For instance, with  $(q < 0.5, r = 1)$  the minimum at  $\theta = \pm\pi$  is located at  $(\varphi_1, \varphi_2) = (\pm\pi, 0)$ . The trajectory makes a closed loop on the  $\varphi_1$ - $\varphi_2$  torus. For  $(q \gg 1, r = 1)$ , the minimum at  $\theta = \pi$  is around  $(0, \frac{1}{2}\pi)$  so that the discontinuity remains. However, if one adds a small asymmetry in “ $r$ ”, the minimum is pushed back to the origin. For instance, for  $(q, r) = (10, 1.3)$ , the minimum starts to turn back at  $\theta \sim 0.8\pi$  and reaches the origin at  $\theta = \pi$ . The implication to QCD physics is profound. As  $m_s \gg m_d > m_u$ , we are facing at a case  $q \gg r > 1$  in which there is no singularity in  $\theta$  any more.<sup>10</sup>

We conclude that a sufficiently large asymmetry in the fermion masses removes the cusp singularity at  $\theta = \pm\pi$  in  $\langle \bar{\psi}\psi \rangle$  at  $T = 0$ .

## Acknowledgments

This work was supported in part by the U.S. Department of Energy under contracts DE-AC02-83ER-40105. The author would like to thank the Aspen Center for Physics for its hospitality where a part of the work was carried out.

## References

1. J. Schwinger, *Phys. Rev.* **125** (1962) 397; **128** (1962) 2425;
2. J.H. Lowenstein and J.A. Swieca, *Ann. Phys. (N.Y.)* **68** (1971) 172; A. Casher, J. Kogut and L. Susskind, *Phys. Rev.* **D10** (1974) 732; S. Coleman, R. Jackiw, and L. Susskind, *Ann. Phys. (N.Y.)* **93** (1975) 267.
3. M.B. Halpern, *Phys. Rev.* **D13** (1976) 337; I. Affleck, *Nucl. Phys.* **B265** [FS15] (1986) 448.
4. S. Coleman, *Ann. Phys. (N.Y.)* **101** (1976) 239.
5. J.E. Hetrick, Y. Hosotani, and S. Iso, *Phys. Lett.* **B350** (1995) 92; ‘The interplay between mass, volume,  $\theta$ , and  $\langle \bar{\psi}\psi \rangle$  in  $N$ -flavor  $QED_2$ ’, hep-th/9510090.
6. N. Manton, *Ann. Phys. (N.Y.)* **159** (1985) 220;
7. J.E. Hetrick and Y. Hosotani, *Phys. Rev.* **D38** (1988) 2621.
8. R. Link, *Phys. Rev.* **D42** (1990) 2103; S. Iso and H. Murayama, *Prog. Theoret. Phys.* **84** (1990) 142.
9. I. Sachs and A. Wipf, *Helv. Phys. Acta.* **65** (1992) 652.
10. Creutz, hep-th/9505112.