

ZU-TH 16/95
 IPNO /TH 95-53
 October 1995

Electromagnetic Corrections to the decays $\pi^0 \rightarrow 3\pi^0$ ¹

R. Baur^a, J. Kambor^b and D. Wyler^a

^a Theoretische Physik, Universität Zurich, CH-8057 Zurich, Switzerland

^b Division de Physique Théorique², Institut de Physique Nucleaire,
 F-91406 Orsay Cedex, France

Abstract

We calculate the electromagnetic corrections to the decays $\pi^0 \rightarrow 3\pi^0$ in next to leading order in the chiral expansion. We find that the corrections are small in accordance with Sutherland's theorem and modify neither rate nor the Dalitz plot distributions noticeably.

¹partially supported by Schweizerischer Nationalfonds.

²Unité de Recherche des Universités Paris 11 et Paris 6, associée au CNRS.

1 Introduction

In the isospin limit $m_u = m_d$ and $e = 0$, the decay $\eta \rightarrow 3\pi$ is forbidden. Consequently, the decay amplitude receives contributions from the QCD isospin violating interaction

$$H_{\text{QCD}}(x) = \frac{(m_d - m_u)}{2} (dd - uu)(x); \quad (1.1)$$

and from the electromagnetic interactions

$$H_{\text{QED}}(x) = \frac{1}{2} e^2 \int^Z dy D(x-y) T(j(x)j(y)); \quad (1.2)$$

where $D(x-y)$ is the photon propagator. Using soft pion techniques, it has been shown that the electromagnetic contribution is much too small to account for the experimentally observed rate [1, 2]. Much work has been devoted to study the effects of the QCD contribution, equation (1.1). Gasser and Leutwyler [3] carried out the one-loop calculation within chiral perturbation theory [4, 5, 6, 7]. Although they found large corrections to the rate³ their result failed to reproduce the experimental rates [8]. Recently, the complete unitary corrections were evaluated [9]. Despite considerable uncertainties, also these corrections are not quite sufficient to account for the observed rate, if the usually assumed value of $(m_u - m_d)$ [10] is taken. Instead, a somewhat smaller up-quark mass is required.

In this situation, it is of interest to reconsider the electromagnetic corrections. Strictly speaking, they may be divided into the indirect ones, affecting the parameters such as F_π which enter the calculation and the direct ones, specific for the process under consideration. In the following we will be concerned with the latter only. Corrections to F_π were calculated previously [11] and change its value by about 1%, leading to a noteworthy increase of the eta decay rate by about 4%.

Within the framework of chiral perturbation theory, electromagnetic correction can also be described by a series of effective operators of increasing power in momentum or masses of the mesons [12]. Sutherland's theorem [1] states that the first correction of order $e^2 p^0$ vanishes where p is a typical momentum. Thus, the electromagnetic corrections are at most of order $e^2 p^2$. If the decay amplitude is assumed to be linear in the Mandelstam variables, these corrections are further suppressed, i.e. p^2 is of order M^{-2} rather than M^2 [2, 13].

A similar result (Dashen's theorem [14]) for the electromagnetic mass differences states that they are equal for the pions and for the Kaons to leading order in the chiral expansion. Recently, it was argued that this equality could be violated [15] (see, however [16]), and one might expect then that also Sutherland's theorem receives larger corrections than previously thought which could enhance the decay rate of the η . In this article we determine the $O(e^2 p^2)$ electromagnetic corrections to the decay amplitudes $\eta \rightarrow 3\pi$. We show that they can be safely neglected compared

³The Dalitz plot distribution does not get equally large corrections from higher orders in the chiral expansion. It is therefore possibly more sensitive to electromagnetic corrections.

to the QCD isospin violating contributions, both for the rates and the Dalitz plot distribution. The paper is organized as follows: in sections 2 and 3 we review the chiral lagrangian for electromagnetic interactions up to order p^2 . In section 4 we calculate the electromagnetic corrections to $\pi^0 \rightarrow \pi^+ \pi^-$ to one-loop chiral perturbation theory. The numerical results are given in section 5 and section 6 contains our conclusions.

2 Effective low-energy Lagrangian

To lowest order in Chiral Perturbation Theory (ChPT), including the electromagnetic interactions, the Lagrangian L_2 is given by [5, 7, 17]

$$L_2 = \frac{1}{4} f_\pi^2 \text{Tr} [\partial_\mu U^\dagger \partial^\mu U] - \frac{1}{2} (\partial_\mu a)^2 + \frac{F_0^2}{4} \text{Tr} [h_R U_R + U_R^\dagger h_L] + \frac{F_0^2}{4} \text{Tr} [h_V U + U^\dagger h_A] + C h_Q U Q_L U^\dagger h_Q : \quad (2.3)$$

Here, f_π is the pion decay constant, the photon field a

$$F_{\mu\nu} = \partial_\mu a_\nu - \partial_\nu a_\mu : \quad (2.4)$$

The parameter χ is the gauge fixing parameter, which is set to $\chi = 1$ henceforth. The pseudoscalar meson fields are contained in the usual way in the matrix U , and the field h_Q incorporates the coupling of the pseudoscalar mesons to the scalar and pseudoscalar currents s and p

$$h_Q = 2B_0 (s + ip) = 2B_0 M + \dots : \quad (2.5)$$

where the quark mass matrix is

$$M = \frac{B_0}{C} \begin{pmatrix} m_u & 0 & 0 \\ 0 & m_d & 0 \\ 0 & 0 & m_s \end{pmatrix} : \quad (2.6)$$

The covariant derivative $D_\mu U$ defines the coupling of the pseudoscalar mesons to the photon field a , the external vector current V , and the external axial current A

$$D_\mu U = \partial_\mu U - i(V_\mu + Q_R a_\mu + A_\mu)U - iU(V_\mu + Q_L a_\mu - A_\mu) : \quad (2.7)$$

The two spurion fields Q_R and Q_L are introduced to construct a $SU(3)_L \times SU(3)_R$ invariant Lagrangian. In the later calculation they will be fixed by $Q_R = Q_L = Q$, where Q is the charge matrix of the three light quarks

$$Q = \frac{e}{3} \begin{pmatrix} 2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} : \quad (2.8)$$

The constant F_0 is up to order $O(m_q)$ the pion decay constant.

The low energy constant C determines the electromagnetic part of the masses $M_\pi; M_K$; of the charged pion and Kaon in the chiral limit

$$ChQUQU^yi = \frac{2e^2 C}{F^2} (\pi^+ \pi^- + K^+ K^-) + \dots \quad (2.9)$$

Clearly, $ChQUQU^yi$ contributes equally to the square of the masses of π and K , in agreement with Dashen's theorem [14].

Since the external currents V and A are of order $O(p)$, where p is their external momentum, the product Qa must have dimension $O(p)$. In order to formally maintain consistent chiral counting, it is convenient to set [12]

$$e = O(p); a = 1; \quad (2.10)$$

such that L_2 is $O(p^2)$.

3 Effective Lagrangian to order $O(p^4)$

At order $O(p^4)$ the generating functional has three different types of contributions:

An explicit local action of order $O(p^4)$

The one-loop graphs associated with the lowest order Lagrangian L_2 .

The anomaly is of order $O(p^4)$; its contributions will not be discussed here.

For a discussion, the reader is referred to the literature [5, 7].

Since we are not concerned here with the strong interactions, [3], we only give the electromagnetic terms. The most general chiral and Lorentz invariant, P and C symmetric Lagrangian of order $O(p^4)$ is [12]

$$\begin{aligned} L_4 = & K_1 F_0^2 \text{tr } U^\dagger \text{tr } U i h Q^2 i + K_2 F_0^2 \text{tr } U^\dagger \text{tr } U i h Q U Q U^i \\ & + K_3 F_0^2 \text{tr } U^\dagger Q U i \text{tr } U^\dagger Q U i + \text{tr } U Q U^\dagger i \text{tr } U Q U^i \\ & + K_4 F_0^2 \text{tr } U^\dagger Q U i \text{tr } U Q U^i + K_5 F_0^2 \text{tr } U^\dagger U^\dagger U^\dagger Q^2 i \\ & + K_6 F_0^2 \text{tr } U^\dagger \text{tr } U Q U^\dagger Q U + \text{tr } U \text{tr } U^\dagger Q U Q U^i \\ & + K_7 F_0^2 \text{tr } U^\dagger U + U^\dagger i h Q^2 i + K_8 F_0^2 \text{tr } U^\dagger U + U^\dagger i h Q U Q U^i \\ & + K_9 F_0^2 \text{tr } U^\dagger U + U^\dagger Q^2 + \text{tr } U^\dagger + U^\dagger Q^2 i \\ & + K_{10} F_0^2 \text{tr } U^\dagger U + U^\dagger Q U^\dagger Q U + \text{tr } U^\dagger + U^\dagger Q U Q U^i \\ & + K_{11} F_0^2 \text{tr } U^\dagger U^\dagger Q U^\dagger Q U + \text{tr } U^\dagger U^\dagger Q U Q U^i \\ & + K_{12} F_0^2 \text{tr } U^\dagger \text{tr } Q^R; Q U + \text{tr } U \text{tr } Q^L; Q U^i \\ & + K_{13} F_0^2 \text{tr } Q^R U \text{tr } Q^L U^i + K_{14} F_0^2 \text{tr } Q^R \text{tr } Q^R + \text{tr } Q^L \text{tr } Q^L i \\ & + K_{15} F_0^4 \text{tr } Q U Q U^i{}^2 + K_{16} F_0^4 \text{tr } Q U Q U^\dagger i h Q^2 i + K_{17} F_0^4 \text{tr } Q^2 i{}^2; \end{aligned} \quad (3.11)$$

where F^R and F^L are the field strength tensors of F^R and of F^L respectively

$$\begin{aligned} F^R &= V + Q^R a + A \\ F^L &= V + Q^L a - A \\ F^I &= \partial F^I - \partial F^I - i F^I; F^I; \quad I = R, L \end{aligned} \quad (3.12)$$

and the covariant derivative r_Q^I is defined as

$$r_Q^I = \partial Q^I + i Q^I; F^I; \quad I = R, L \quad (3.13)$$

Under a chiral transformation the covariant derivative r_Q^I transforms like the corresponding Q^I

$$\begin{aligned} r_Q^I &\rightarrow V_I r_Q^I V_I^\dagger \quad I = R, L \\ V_{R,L} &\in SU(3)_{R,L} \end{aligned} \quad (3.14)$$

We have set $Q^R = Q^L = Q$. Only in the covariant derivative r_Q^I have we kept the index $I = R, L$ in order to make explicit the proper chiral transformation. The coupling constants $K_1 :: K_{17}$ are not determined by chiral symmetry. While in principle calculable from the fundamental lagrangian of QCD and QED, we consider them as phenomenological constants. $K_1; K_7; K_{16}$ are electromagnetic corrections to F_0^2, B_0 and C respectively.

In the next step, the loops generated by the Lagrangian L_2 are calculated. They lead to terms proportional to p^4 . Their divergencies are cancelled by finite counterterms in the constants K_i . The procedure is standard and will not be displayed here in detail [12]. Using the usual dimensional regularization scheme with scale μ , one finds that the ultraviolet divergencies can be absorbed in the coupling constants K_i with the following renormalization of the low-energy coupling constants [12]:

$$\begin{aligned} K_i &= K_i^r(\mu) + \delta K_i \\ &= \frac{d-4}{16\pi^2} \frac{1}{d-4} - \frac{1}{2} \ln(4\pi) + \gamma(1) + 1 \end{aligned} \quad (3.15)$$

and

$$\begin{aligned}
g_1 &= \frac{3}{4} & g_2 &= Z & g_3 &= \frac{3}{4} \\
g_4 &= 2Z & g_5 &= \frac{9}{4} & g_6 &= \frac{3}{2}Z \\
g_7 &= 0 & g_8 &= Z & g_9 &= \frac{1}{4} \\
g_{10} &= \frac{3}{2}Z + \frac{1}{4} & g_{11} &= \frac{1}{8} & g_{12} &= \frac{1}{8} \\
g_{13} &= 0 & g_{14} &= 0 & g_{15} &= 20Z^2 + 3Z + \frac{3}{2} \\
g_{16} &= 4Z^2 - \frac{3}{2}Z - 3 & g_{17} &= 2Z^2 - \frac{3}{2}Z + \frac{3}{2} \\
Z &= \frac{C}{F_0^4}
\end{aligned} \tag{3.16}$$

Here, the $K_i^r(\mu)$ are the renormalized couplings at the scale μ .

The scaling behaviour of $K_i^r(\mu)$ is determined by the requirement

$$\frac{dK_i^r}{d\mu} = 0 \tag{3.17}$$

which implies for $K_i^r(\mu)$ in the limit $d = 4$

$$K_i^r(\mu) = K_i^r(\mu_0) - \frac{i}{16\pi^2} \ln\left(\frac{\mu}{\mu_0}\right) \tag{3.18}$$

4 Electromagnetic contributions to $\pi^0 \rightarrow 3\pi$

The electromagnetic contributions to the decay $\pi^0 \rightarrow 3\pi$ have been considered long ago [2, 13]. The transition amplitude is suppressed due to a soft-pion theorem [1]. At next-to-leading order in a low energy expansion, there is a further suppression factor $m_\pi^2 = m_K^2$, if linear dependence on the energy of the odd pion is assumed. This led to the conclusion that the electromagnetic interactions alone fail completely to explain the observed rate.

Here the problem is reinvestigated in the framework of effective chiral lagrangians, as reviewed in section 2. We calculate the corrections of order $e^2 p^2$, where p^2 is a generic low energy momentum. We work in the isospin limit, as corrections of order $e^2 (m_d - m_u)$ are very small.

As usual, we define the decay amplitude A by

$$\langle \pi^0 + \pi^+ \pi^- | T | \pi^0 \rangle = i(2\pi)^4 \delta^4(p_f - p_i) A(s; t; u) \tag{4.19}$$

with the Mandelstam variables

$$s = (p_1 + p_2)^2 \quad t = (p_1 - p_3)^2 \quad u = (p_1 - p_4)^2 \tag{4.20}$$

Due to G-parity, the three pions emerge in a $I = 1$ state. Hence, the decay amplitude into three neutral pions, $A(s; t; u)$, is also determined by $A(s; t; u)$

$$\begin{aligned} \langle 0^+ 0^+ 0^+ \text{out} | j \rangle &= i (2^4)^{-4} (P_f - P_i) A(s; t; u) \\ A(s; t; u) &= A(s; t; u) + A(t; u; s) + A(u; s; t): \end{aligned} \quad (4.21)$$

Charge conjugation invariance requires $A(s; t; u)$ to be symmetric with respect to interchanges of t and u .

The electromagnetic contributions to $A(s; t; u)$ may be written as

$$A_{\text{QED}}(s; t; u) = e^2 f(p^2; m_u; m_d; m_s; \dots) \quad (4.22)$$

where e is the scale of the theory and p^2 stands for any of the variables s, t, u . Keeping ratios $p^2/m_u, m_d, m_s$ fixed, f can be expanded according to

$$f = f_0 + f_1 + \dots : \quad (4.23)$$

f_0 is of order p^0 , f_1 of order p^2 etc. The lowest order term f_0 vanishes, as was shown by Sutherland [1]. As noted, f_1 is also suppressed, if the energy dependence of the amplitude is linear. This follows from the soft-pion theorem [1]

$$\lim_{p_0 \rightarrow 0} A_{\text{QED}}^{+0} = \lim_{p_0 \rightarrow 0} A_{\text{QED}}^{000} = 0 (e^2 \hat{m}): \quad (4.24)$$

Hence, for a linear amplitude

$$A_{\text{QED}}^{+0} = a + b s; \quad (4.25)$$

it follows that at the soft-pion point $s = M^2 + 3M^2$, and $t = u = 0$

$$A_{\text{QED}}^{+0}(p_0 = 0) = a + b(M^2 + 3M^2) \quad (4.26)$$

$$A_{\text{QED}}^{000}(p_0 = 0) = 3a + b(M^2 + 3M^2); \quad (4.27)$$

and therefore

$$a = 0; \quad b = O\left(\frac{\hat{m}}{m_s}\right): \quad (4.28)$$

In chiral perturbation theory the term f_1 can be calculated in a systematic manner. The assumption (4.25) is seen to be violated by non-local terms arising from one-loop graphs. However, only kaons appear in the loop, therefore the resulting amplitude is smooth throughout the physical region. Equation (4.25) is then still a good approximation and the $\frac{\hat{m}}{m_s}$ suppression is effectively at work. This is the main reason why the next-to-leading order term f_1 is small.

The relevant diagrams are shown in figure 1. The graphs a) and b) are the non-local unitary and the tadpole graphs respectively, and c) contains the direct and pole graphs.




Figure 1: Graphs contributing to the decay $\pi^0 \rightarrow 3\pi$. The filled circles denote vertices of order $O(e^2)$ and the filled boxes those of order $O(e^2 p^2)$.

The electromagnetic vertex in these diagrams is given by the leading order term in eq. (2.3) and by eq. (3.11). There are no graphs with pions in the loop as they are in addition suppressed by the η mixing angle θ , which is of the order

$$O(\theta) \sim 10^{-2} = O\left(\frac{m_u - m_d}{m_s}\right):$$

Also, the mass difference $(M_K^2 - M_{K^0}^2)_{\text{QCD}}$ contributes only to the order $O(\frac{m_u - m_d}{m_s})$ and can be neglected.

We then obtain for the unitary corrections

$$U(s; t; u) = \frac{C}{12 \frac{p}{3F_0^6}} \left[3s(3s - 4M_K^2) \frac{1}{i} C_{kk}(s) + 6(3s - 4M_K^2) J_{kk}(s) \right. \\ \left. + 3(3t - 4M_K^2) J_{kk}(t) + 3(3u - 4M_K^2) J_{kk}(u) \right. \\ \left. + (5s - 4M_K^2) J_{kk}(0) \right]; \quad (4.29)$$

for the tadpoles

$$T(s) = \frac{C}{9 \frac{p}{3F_0^6}} \left[(s - s_0) \left(\frac{9}{4} + \frac{2M^2}{M^2 - M_K^2} \right) \right. \\ \left. + \frac{2M^2}{3} \frac{5s}{12} + \frac{M_K^2}{3} J_{kk}(0) \right]; \quad (4.30)$$

and finally for the sum of the pole and direct graphs

$$D(s) = \frac{4M^2}{9 \frac{p}{3F_0^6}} \left[1 + \frac{3(s - s_0)}{M^2 - M_K^2} \right] \\ + f \left[3K_3 + \frac{3}{2} K_4 + K_5 + K_6 - K_9 - K_{10} g \right]; \quad (4.31)$$

We have introduced the following functions C_{kk} and J_{kk} :

$$\frac{1}{i} C_{kk}(s) = \frac{1}{16^2} \frac{1}{M_K^2} \frac{p^2}{p - s} \arctan \frac{s}{p - s} \\ J_{kk}(s) = \frac{1}{16^2} \left[2 - \frac{2}{s} \arctan \frac{s}{p - s} \right] + J_{kk}(0) \\ J_{kk}(0) = \frac{1}{2} - 2k + O(d^{-4}) \quad (4.32)$$

with

$$s = \frac{s}{M_K^2} = s(4 - s) \\ 3s_0 = s + t + u = m^2 + 3m^2; \quad (4.33)$$

and

$$= \frac{d^{-4}}{16^2} f \frac{1}{d^{-4}} - \frac{1}{2} (\ln(4) + {}^0(1) + 1) g \\ k = \frac{1}{32^2} f \ln \frac{M_K^2}{2} + 1g; \quad (4.34)$$

With equation (3.15) and the following definition:

$$J_{kk}(s) = J_{kk}(s) - J_{kk}(0) \quad (4.35)$$

we obtain the finite result

$$\begin{aligned}
A_{QED}(s;t;u) &= e^2 (U(s;t;u) + T(s) + D(s)) \\
&= \frac{C e^2}{3F_0^6} \left[\frac{1}{4} s (3s - 4M_K^2) \frac{1}{i} C_{kk}(s) + \frac{1}{2} (3s - 4M_K^2) J_{kk}(s) \right. \\
&\quad \left. + \frac{1}{4} (3t - 4M_K^2) J_{kk}(t) + \frac{1}{4} (3u - 4M_K^2) J_{kk}(u) \right] \\
&\quad + \frac{4C e^2 M^2}{3 \cdot 3F_0^6} f_K \frac{F_0^4}{3C} K^r(\cdot) g \left[1 + \frac{3(s - s_0)}{M^2 - M^2} \right]
\end{aligned} \tag{4.36}$$

with

$$\begin{aligned}
K^r(\cdot) &= 3K_3^r(\cdot) + \frac{3}{2} K_4^r(\cdot) + K_5^r(\cdot) \\
&\quad + K_6^r(\cdot) - K_9^r(\cdot) - K_{10}^r(\cdot);
\end{aligned} \tag{4.37}$$

Some remarks concerning the structure of this result are in order:

1. The scale dependence of k is cancelled by the running of $K^r(\cdot)$. Using equations (3.16) and (3.18) one gets

$$K^r(\cdot) = K^r(s_0) - \frac{3C}{F_0^4 32^2} \ln\left(\frac{s}{s_0}\right); \tag{4.38}$$

2. As expected, the amplitude vanishes in the soft pion point

$$s \rightarrow M^2; \quad t, u \rightarrow 0 \tag{4.39}$$

if the lowest order relation $3m^2 = 4m_K^2$ is used. This is in accord with Sutherland's theorem which predicts a vanishing amplitude if the pion mass is sent to zero.

3. The final result (4.36) consists of a nonlocal piece (first two lines) and a polynomial to first order in s (last line). The polynomial part is in proportion to the current algebra amplitude of strong interactions. It exhibits the M^2 suppression as implied by Sutherlands theorem. Since the counterterm K^r enters only this part of the amplitude, it's contribution is suppressed too. The nonlocal piece circumvents this suppression as it is clearly not linear in s . However, since only Kaon loops contribute, this part is kinematically suppressed.
4. The rate is to a large extent fixed by the amplitude at the center of the Dalitz plot. We therefore expand (4.36) according to

$$A_{QED}(s;t;u) = a_0 + a_1(s - s_0) + a_2(s - s_0)^2 + a_3(t - u)^2; \tag{4.40}$$

Explicitly, the constant term is given by

$$a_0 = \frac{C e^2}{3 F_0^6} \left(\frac{3}{4} s_0 (3s_0 - 4M_K^2) \frac{1}{i} C_{kk}(s_0) + 4M_K^2 k \frac{F_0^4}{3C} K^r(\cdot) \right) : \quad (4.41)$$

Note that at $s = s_0$ the remaining nonlocal piece in proportion to C_{kk} is numerically less suppressed than for instance J_{kk} .

5. The linear slope a_1 is given in terms of the same counterterm K^r as is the amplitude at the center of the physical region. Thus, if isospin breaking in the quark masses is switched off, the electromagnetic rate would be fixed in terms of the linear slope. The quadratic coefficients a_2, a_3 get contributions only from the nonlocal part of A_{QED} . They are given in terms of the known coupling constant C .

5 Numerical Results

As mentioned, the one-loop result of Gasser and Leutwyler yields a rate of 167 eV, much below the experimental value. Although the unitary corrections are available [9], we will use the one-loop result as reference. This is inessential, since the electromagnetic corrections are very small.

We set F_0 equal to the pion decay constant F , whose experimental value is $F = 92.4$ MeV if electromagnetic corrections are included [11]. The low energy constant C can be determined by using the underlying vector and axial-vector resonances. Using the Weinberg sum rules [18], one obtains [17]

$$C = \frac{3}{32} M^2 F^2 \ln \left(\frac{F^2}{F^2 - F^2} \right); \quad (5.42)$$

where M is the mass of the ρ -meson, $M = 770$ MeV and F is its decay constant, $F = 154$ MeV. Numerically,

$$\frac{C}{F^4} = 0.84; \quad (5.43)$$

The values of the coupling constants K_i^r are so far unknown. The naive chiral counting law [19] gives

$$jK_i^r \approx \frac{1}{16^2} \approx 6.3 \cdot 10^{-3}; \quad (5.44)$$

while an estimate for K_8^r yields [12]

$$K_8^r(\sqrt{s} = 0.5 \text{ GeV}) = (1.0 - 1.7) \cdot 10^{-3}; \quad (5.45)$$

consistent with the naive rule (5.44).

We conservatively use eq. (5.44) and vary the absolute value of K^r between the bounds

$$jK^r \approx \frac{8.5}{16^2}; \quad (5.46)$$

To get a feeling of the numerical size of the electromagnetic corrections, we expand the amplitude (4.36) around the center of the Dalitz plot and compare it to the ChPT one-loop result for the strong interactions. Writing the Taylor expansion of the sum of these amplitudes as

$$A_{QCD}(s;t;u) + A_{QED}(s;t;u) = a_0 + a_1(s - s_0) + a_2(s - s_0)^2 + a_3(t - u)^2 \quad (5.47)$$

we find

$$\begin{aligned} a_0 &= a_0^{QCD} + a_0^{n.l.} + a_0^1 \\ &= (0.17 + 0.02i) + 0.14 \cdot 10^{-2} - 0.30 \cdot 10^{-2}; \end{aligned} \quad (5.48)$$

where a_0^{QCD} denotes the strong one-loop value, $a_0^{n.l.}$ the contribution from the non-local terms in eq.(4.36), and a_0^1 that of the local terms which depend on the low energy coupling constants K_i . The variation of a_0^1 are due to the changing value of K^π . Similarly, we obtain

$$\begin{aligned} a_1 &= a_1^{QCD} + a_1^{n.l.} + a_1^1 \\ &= (2.25 + 0.75i) + 0.62 \cdot 10^{-3} - 0.30 \cdot 10^{-1} [\text{GeV}^{-2}]; \end{aligned} \quad (5.49)$$

$$\begin{aligned} a_2 &= a_2^{QCD} + a_2^{n.l.} \\ &= (1.33 - 0.91i) - 0.25 \cdot 10^{-1} [\text{GeV}^{-4}]; \end{aligned} \quad (5.50)$$

$$\begin{aligned} a_3 &= a_3^{QCD} + a_3^{n.l.} \\ &= (0.93 - 0.41i) - 0.85 \cdot 10^{-2} [\text{GeV}^{-4}]; \end{aligned} \quad (5.51)$$

Here, we used everywhere the new value of F_π [11]; the other relevant factors for the QCD amplitude are taken from ref [3]. From this we see that for the extreme values of K^π the local contribution dominates over the non-local contribution. Both are, however, small compared to the ChPT one-loop result. We therefore expect no significant changes from QED to both, the decay rates as well as the Dalitz plot distributions for the decays $\pi^0 \rightarrow \pi^0 \pi^0$ and $\pi^0 \rightarrow 3\pi^0$.

The decay rate $\Gamma_{\pi^0 \rightarrow \pi^0 \pi^0}$ is obtained by evaluating the phase space integral⁴

$$\begin{aligned} d\Gamma_{\pi^0 \rightarrow \pi^0 \pi^0} &= \frac{(2\pi)^4}{M} (P_i - P_f) j A_{QCD}(s;t;u) + A_{QED}(s;t;u) j^2 d\Phi; \\ d\Phi &= \frac{d^3p_0}{(2\pi)^4 2p_0^0} \frac{d^3p_+}{(2\pi)^4 2p_+^0} \frac{d^3p_-}{(2\pi)^4 2p_-^0}; \end{aligned} \quad (5.52)$$

We get for the integrated rate

$$\Gamma_{\pi^0 \rightarrow \pi^0 \pi^0} = 165 \cdot 5\text{eV} \quad (5.53)$$

where the uncertainty is due to our lack of knowledge on K^π , and for the ratio r

$$1.42 \leq r = \frac{\Gamma_{\pi^0 \rightarrow 3\pi^0}}{\Gamma_{\pi^0 \rightarrow \pi^0 \pi^0}} \leq 1.43 \quad (5.54)$$

⁴Note that the phase space is very small and thus very sensitive to the mass difference $M_{\pi^0} - M_{\pi^\pm}$.

In both cases electromagnetism gives, as expected, small corrections to the results obtained by [3], where the purely strong effects have been considered

$$\Gamma(K^0 \rightarrow \pi^0 \pi^0)_{\text{QCD}} = 167\text{eV} \quad (5.55)$$

$$r^{\text{QCD}} = 1.43 \quad (5.56)$$

Even though QED can give rise to a correction of up to 4 per cent to the decay rate $\Gamma(K^0 \rightarrow \pi^0 \pi^0)$, the Dalitz plot distribution remains unchanged. This is due to the fact that the non-local contributions are small and that the functional dependence of the local term in eq.(4.36) on the Mandelstam variables and that of the current algebra amplitude are the same. Since the Dalitz plot distribution is a measure of the s -dependence of the amplitude $A(s;t;u)$, the $K^*(\pi^0)$ contribution only leads to a different normalization of the distribution, but there is no significant change of the slopes.

6 Conclusions

In this paper we have calculated the direct electromagnetic corrections to the decay $K^0 \rightarrow \pi^0 \pi^0$. Using the general framework of chiral perturbation theory, we determine the first nontrivial term of order $e^2 p^2$. This includes mesonic loops and a combination of counterterms introduced earlier [12]. Unlike for meson mass differences, the corrections are tiny, and amount to at most two percent of the ChPT one-loop amplitude at the center of the decay region.⁵ The Dalitz plot parameters are also rigid with respect to electromagnetic corrections; from the results in section 5 we see that both linear and quadratic slope parameters are practically unaffected by the electromagnetic corrections. This is clear from the form of the amplitude: First, the nonlocal part which circumvents the Bell-Sutherland suppression arises from Kaon loops, which are kinematically strongly suppressed. The larger (by a factor of 13) pion loops multiply the small quantity $(m_d - m_u)$ and can be neglected. Furthermore, the polynomial part is suppressed by the small factor $M^2 = M_K^2$. This includes in particular the contributions from the coupling constants K_i . Thus, the dependence on these little known constants is weak; varying them between the rather large values $\frac{8.5}{16^2}$ results only in a variation of less than two percent relative to the one-loop ChPT amplitude of the strong interactions.

Thus, our result confirms the usual picture, namely that the large decay rate requires rather large isospin violations in the strong interactions. Moreover, any precise measurement of the $K^0 \rightarrow \pi^0 \pi^0$ Dalitz plot yields important information on the strong interactions without contamination from the electromagnetic ones.

Acknowledgements.

⁵There are other, indirect, electromagnetic corrections which influence the fundamental parameters, such as F_π . They are known [11] and change the normalization of the QCD amplitude noticeably, resulting in a 8 % increase of the rate.

We thank R. Urech for discussions and G.-J. Van Oldenborgh for the help in the numerical part. J.K. would like to acknowledge support from the Department of Physics and Astronomy, University of Massachusetts, where the work described herein was started.

References

- [1] D.G. Sutherland, Phys. Lett. 23 (1966) 384.
- [2] J.S. Bell and D.G. Sutherland, Nucl. Phys. B 4 (1968) 315.
- [3] J. Gasser and H. Leutwyler, Nucl. Phys. B 250 (1985) 539.
- [4] S. Weinberg, Physica A 96 (1979) 327.
- [5] J. Donoghue, E. Golowich, and B. Holstein, Dynamics of the Standard Model, CUP, Cambridge 1992; H. Leutwyler, Ann. Phys. 235 (1994) 165.
- [6] H. Leutwyler, Principles of Chiral Perturbation Theory', BUTP-94/13 (1994).
- [7] J. Gasser and H. Leutwyler, Nucl. Phys. B 250 (1985) 465.
- [8] Review of Particle Properties, Phys. Rev. D 50 (1994).
- [9] J. Kambor, C. Wiesendanger, and D. Wyler, preprint ZU-TH 41/94; A.V. Anisovich and H. Leutwyler, Bern University preprint BUTP 95/1.
- [10] J. Gasser and H. Leutwyler, Phys. Rep. 87 (1982) 77; H. Leutwyler, Nucl. Phys. B 337 (1990) 108; J. Donoghue and D. Wyler, Phys. Rev. D 45 (1992) 892; D. Wyler, Light Quark Masses beyond Leading Order', Proc. XVI Kazimierz Meeting, World Scientific 1994, Edts. Z. Ajduk et al.; H. Leutwyler, Masses of the Light Quarks', BUTP-94/8 (1994).
- [11] B. Holstein, Phys. Lett. 244 (1990) 83; Review of Particle Properties, Phys. Rev. D 50 (1994).
- [12] R. Urech, Nucl. Phys. B 433 (1995) 234.
- [13] P. Dittner, P.H. Dondi, and S. Eliezer, Phys. Rev. D 8 (1973) 2253.
- [14] R. Dashen, Phys. Rev. 183 (1969) 1245.
- [15] J. Donoghue, B. Holstein, and D. Wyler, Phys. Rev. D 47 (1993) 2089; J. Bijnens, Phys. Lett. B 95 (1993) 343.
- [16] R. Baur and R. Urech, preprint ZU-TH 22/95, hep-ph/9508393.
- [17] G. Ecker et al., Nucl. Phys. B 321 (1989) 311.

[18] S. Weinberg, Phys. Rev. Lett. 18 (1967) 507.

[19] A. Manohar and H. Georgi, Nucl. Phys. B 234 (1984) 189.