

DOUBLE SCALING OF ANGULAR CORRELATIONS INSIDE JETS ^a

W olfgang O chs

M ax-Planck-Institut für Physik, Fohringer Ring 6, D-80805 München, Germany

Jacek W osiek

Jagellonian University, Reymonta 4, PL-30-059 Cracow, Poland

Angular correlations of partons have been derived for high energy jets in QCD. Interesting new scaling properties with two redundant variables (jet energy P and jet opening angle θ) emerge which can be tested within a parton-hadron duality picture. Recent results from LEP support the scaling in θ ; the scaling in both P and θ could be tested with jets from deep inelastic scattering or with high p_T jets.

1 Introduction

There are indications that there is a soft confinement mechanism which does not lead to large rearrangements of momenta during the hadronisation process, so that QCD predictions on multiparton final states can be meaningfully compared with predictions on hadronic final states^{1,2}. An extreme possibility is the hypothesis of "Local Parton-Hadron Duality" (LPHD)³, according to which the parton cascade

The energy spectra of particles gave initial support for this idea, in that they showed the same depletion for soft particles as did the spectra of particles due to the soft gluon interference^{3,4}. Also the effect of the running of α_s is clearly visible in the energy spectra⁵. It is therefore important to test these ideas for other quantities as well, in particular multiparticle observables.

Recently we have studied the properties of angular correlations in jets within the double logarithmic approximation (DLA) of QCD which provides the leading order results at high energies⁶. We derived explicit expressions for various two- and multiparton observables. Here we will report on some general scaling and universality properties of our results which follow from the QCD cascade with angular ordering.

Results partly overlapping with ours have been also obtained by other groups using somewhat different theoretical approaches, on angular correlations, in particular azimuthal correlations⁷ and on multiplicity moments^{8,9}.

2 Theoretical scheme

We start from an integral equation for the generating functional in DLA^{10,2} from which the properties of the cascade can be derived. It takes into account the leading collinear and soft divergencies but neglects recoil effects.

^a presented by W. Ochs at the International Europhysics Conference on High Energy Physics, Bussels, July 1995

This yields the correct high energy asymptotic predictions and we investigated to what extent these predictions apply to present energies by comparing with the more complete MC¹¹ calculations. We derive from the generating functional the general n -parton cumulant correlation function $\langle n \rangle(\theta_1, \dots, \theta_n)$ in the spherical angles θ_i . Simpler quantities are obtained by partial integration. Finally we checked the backward consistency of our results by full integration, so our two-parton correlation $\langle 2 \rangle$ integrates to the global multiplicity moment $C_2 = F_2 - 1 = \langle n(n-1) \rangle = \langle n \rangle^2 - 1 = 1=3$, the well-known DLA result.

3 Scaling results

We consider the distribution $\langle 2 \rangle(\theta_{12}; P; \theta)$ in the relative angle θ_{12} between two partons inside a cone of half opening angle θ around the primary parton of momentum P , also the quantity $\hat{r} = \langle 2 \rangle / n^2$ normalized by the average multiplicity n in the cone or $r = \langle 2 \rangle / \langle n \rangle^2$, where $\langle n \rangle$ is constructed from uncorrelated pairs ("event mixing"). We find that this correlation function, after an appropriate rescaling depends on the three variables θ_{12}, P and θ only through the single variable $\theta(\theta_{12}; P; \theta)$

$$\frac{\ln \langle 2 \rangle(\theta_{12}; P; \theta)}{2 \ln P} = \theta(\theta_{12}; 2) \frac{P}{2} \frac{1}{\theta} \quad (1)$$

$$\theta = \frac{\ln \frac{\theta_{12}}{P}}{\ln \frac{P}{\theta}} \quad (2)$$

The function $\theta(\theta_{12}; 2)$ determines the exponential growth of $\langle 2 \rangle$ and the second term in (1) of $\langle n \rangle$. The function $\theta(\theta_{12}; n)$ enters the n -parton correlation function and is exactly calculable, also various approximate forms are known⁶. The rescaling factor in (1) determines the exponential growth of multiplicity and can also be written as $\ln n = 2 \ln \frac{P}{\theta}$ where $2 = 4N_c = (11N_c - 3) - 2N_f = 3$

