

Twist-three distributions and their appearance in the doubly-polarized Drell-Yan process¹

R.D. Tangerman and P.J. Mulders

National Institute for Nuclear Physics and High Energy Physics
(NIKHEF), P.O. Box 41882, 1009 DB Amsterdam, The
Netherlands

Abstract

The twist-three distributions $g_2(x)$ and $h_2(x)$ are defined as quark-field matrix elements between polarized hadron states. They can be written in terms of quark-mass and gluonic operators, after which the Burkhardt-Cottingham sum rule for g_2 can be derived and a similar sum rule for h_2 . Their role in the Drell-Yan double-spin asymmetry A_{LT} is explained.

¹Talk presented by R.D. Tangerman at 2nd Meeting on Possible Measurements of Singly Polarized pp and pn Collisions at HERA (DESY-TH Zeuthen, August 31 – September 2, 1995).

1 Introduction

Although the basic ingredients for high-energy scattering experiments, the quark distribution functions defined as quark-eld matrix elements, can not be calculated from QCD directly, certain sum rules have a more general origin. The Adler sum rule, for instance, follows from current conservation. At sub-leading twist, one has the Burkhardt-Cottingham (BC) sum rule for g_2 , and a similar sum rule for h_2 , which follow from rotational invariance [1, 2]. First attempts have been made to measure the BC sum rule [3]. The measurement of the h_2 sum rule may be possible in the doubly-polarized Drell-Yan process.

Within classes of equal twist, parton distribution functions can be distinguished by their spin-content. For unpolarized hadrons, for instance, one has the light-cone correlator²

$$\int \frac{d}{4} e^{i \cdot x} \langle P | \bar{\psi} \psi | 0 \rangle = f_1(x) p + f_4(x) M^2 n; \quad (1)$$

where the hadron momentum reads $P = p + (M^2=2)n$ in terms of the vectors p and n , satisfying $p^2 = n^2 = 0$ and $p \cdot n = 1$. Not an unimportant detail is the use of the light-cone gauge $A \cdot n = 0$, such that the path-ordered link operator $P \exp[i \int_0^1 d\lambda A(n) \cdot n]$, inserted between the quark-elds to ensure color gauge invariance, becomes unity. The twist-two momentum distribution $f_1(x)$ will contribute to the leading DIS structure function $F_T(x_B; Q^2)$, the logarithmic Q^2 dependence coming from radiative corrections (GLAP evolution).

If the hadron is polarized with spin vector $S = (S \cdot n)p + S_T \cdot (M^2=2)(S \cdot n)n$, the axial vector matrix element is parametrized like

$$\int \frac{d}{4} e^{i \cdot x} \langle P | \bar{\psi} \gamma_5 \psi | 0 \rangle = g_1(x) (S \cdot n)p + g_T(x) S_T + g_3(x) M^2 (S \cdot n)n; \quad (2)$$

where $g_1(x)$ is the twist-two helicity distribution. In this note we will concentrate on the combination $g_2(x) = g_T(x) - g_1(x)$, containing the transverse spin distribution, and its longitudinal counterpart $h_2(x)=2 = h_L(x) - h_1(x)$ defined by the light-cone correlator

$$\int \frac{d}{4} e^{i \cdot x} \langle P | \bar{\psi} \gamma_5 \psi | 0 \rangle = \frac{h_1(x)}{M} (S_T \cdot p - S_T \cdot p) + h_L(x) M (S \cdot n)(p \cdot p - p \cdot n) + h_3(x) M (S_T \cdot n - S_T \cdot n) \quad (3)$$

where $h_1(x)$ is the twist-two transversity distribution.

²We use the notation and conventions of Jaffe and Ji in Ref. [4].

2 Sum rules

The parton interpretation of the higher-twist distribution function is much more involved than for the twist-two case. This arises because of the mixing in the operator-product-expansion of gluonic and quark-mass operators [5, 6]. Consider first the transverse-spin distribution $g_2(x)$. In Ref. [2] it is shown to be closely related to the existence of intrinsic quark transverse momentum in the nucleon. Using the Dirac equation and Lorentz covariance, it can be rewritten as

$$g_2(x) = g_1(x) + \int_x^1 \frac{dy}{y} g_1(y) + \frac{m}{M} \frac{h_1(x)}{x} \int_x^1 \frac{dy}{y} \frac{h_1(y)}{y} + g_T(x) \int_x^1 \frac{dy}{y} g_T(y); \quad (4)$$

where m is the current-quark mass. The distribution g_T contains explicit gluon fields, and is defined by

$$x g_T(x) S_T = \frac{d}{8} e^{i x \cdot p} \langle p | S \cdot j^-(0) g A_T(0) [g_T \not{e}_5 + i \not{e}_T \not{e}] | n \rangle \rangle + h.c.; \quad (5)$$

using the transverse tensors $g_T = g_{\perp} p_{\perp} p_{\perp}$ and $\not{e}_T = \not{e}_{\perp} p_{\perp}$. The decomposition (4) is interesting, since it gives rise to the Burkhardt-Cottingham sum rule [7]

$$\int_0^1 dx g_2(x) = 0; \quad (6)$$

provided the order of x and y integrations may be interchanged. This sum rule has received some attention recently. Specifically, it was shown that for a free-quark target it acquires no $O(1)$ corrections [8].

In a similar fashion, the longitudinal-spin distribution $h_2(x)$ can be decomposed as

$$\begin{aligned} \frac{h_2(x)}{2} = & h_1(x) + 2x \int_x^1 \frac{dy}{y^2} h_1(y) + \frac{m}{M} \frac{g_1(x)}{x} \int_x^1 \frac{dy}{y^2} \frac{g_1(y)}{y} \\ & + h_L(x) \int_x^1 \frac{dy}{y^2} h_L(y); \end{aligned} \quad (7)$$

with the interaction-dependent function

$$x h_L(x) M (S \cdot n) = \frac{d}{8} e^{i x \cdot p} \langle p | S \cdot j^-(0) g A_T(0) \not{e}_5 | n \rangle \rangle + h.c.; \quad (8)$$

Equation (7) leads to the sum rule [1, 2]

$$\int_0^1 dx h_2(x) = 0; \quad (9)$$

for the distribution function h_2 . The validity of this sum rule for the structure function h_2 was questioned recently by Burkardt [1, 9].

3 Experiments

Writing down sum rules for parton distribution functions is one thing, extracting them from experiment is another. The distribution $g_2(x)$ appears cleanest in the cross section for the scattering of a longitudinally polarized lepton on a transversely polarized hadron, measuring only the scattered lepton momentum. From the handbag diagram in Fig. 1 and two diagrams involving a quark-gluon-quark correlator, one finds that the structure function $g_2(x_B; Q^2)$ at tree-level equals $(1/2) \sum_a e_a^2 g_2^a(x_B)$, the a denoting quark and antiquark flavors. Since $h_2(x)$ is chirally odd, one needs a second hadron to flip helicities. Thus we are forced to consider, for instance, Drell-Yan scattering. The analogue to the handbag diagram is depicted in Fig. 2. After averaging over the transverse momentum q_T of the produced massive photon, the distributions appear pair-wise in the double-spin asymmetry [4, 2]

$$A_{LT} = \frac{(\sum_a e_a^2 g_1^a(x) y [g_T^a(y) + g_T^a(y)] + x [h_L^a(x) + \bar{h}_L^a(x)] h_1^a(y)}{(\sum_a e_a^2 f_1^a(x) f_1^a(y))} : (10)$$

The helicity of the longitudinally polarized hadron is given by $\lambda_A = S_A \sin \theta_A$. The angles θ_A and θ_B are of those the lepton axis in the photon rest-frame. Also, $x = P_B \cdot q / P_B \cdot P$ and $y = P_A \cdot q / P_A \cdot P$. It is clear that $h_2(x)$ cannot easily be extracted from the above expression, since it pre-supposes knowledge of the other distributions. Also, the combinations $g_T + \bar{g}_T$ and $h_L + \bar{h}_L$ appear, making life even more complicated. Simply neglecting quark transverse momentum such that (for zero quark mass) $\bar{g}_T = g_T$ and $\bar{h}_L = h_L$, as is done in Ref. [4], is an inconsistent approximation [2].

This work is part of the research program of the foundation for Fundamental Research of Matter (FOM) and the National Organization for Scientific Research (NWO).

References

- [1] M .Burkardt, in Proceedings of 10th International Symposium on High Energy Spin Physics, Nagoya, Japan, Nov. 1992, edited by Hasegawa et al. (Universal Academy Press, Tokyo, 1993).
- [2] R . D . Tangeman and P . J . Mulders, Polarized twist-three distribution functions g_2 and h_2 and the role of intrinsic transverse momentum , 1994, preprint NIKHEF-94-P 7, hep-ph/9408305.
- [3] D . Adams et al., Phys. Lett. B 336, 125 (1994).
- [4] R . L . Jaffe and X . Ji, Nucl. Phys. B 375, 527 (1992).
- [5] R . L . Jaffe, Comments Nucl. Part. Phys. 19, 239 (1990).
- [6] R . L . Jaffe and X . Ji, Phys. Rev. D 43, 724 (1991).
- [7] H . Burkhardt and W . N . Cottingham , Annals of Physics 56, 453 (1970).
- [8] G . Altarelli, B . Lampe, P . Nason, and G . Ridol , Phys. Lett. B 334, 187 (1994).
- [9] M . Burkardt, On the possible violation of sum rules for higher-twist parton distributions, 1995, hep-ph/9505226.




Fig. 1. Handbag diagram .




Fig. 2. Drell-Yan tree-level diagram .