

The Goldstone boson equivalence theorem with fermions

Loyal Durand*

Department of Physics, University of Wisconsin-Madison,
1150 University Avenue, Madison, WI 53706

and

Kurt Riesselmann†

Physik Department T30, Technische Universität München,
James-Franck Strasse, 85747 Garching b. München, Germany

Abstract

The calculation of the leading electroweak corrections to physical transition matrix elements in powers of M_H^2/v^2 can be greatly simplified in the limit $M_H^2 \gg M_W^2, M_Z^2$ through the use of the Goldstone boson equivalence theorem. This theorem allows the vector bosons W^\pm and Z to be replaced by the associated scalar Goldstone bosons w^\pm, z which appear in the symmetry breaking sector of the Standard Model in the limit of vanishing gauge couplings. In the present paper, we extend the equivalence theorem systematically to include the Yukawa interactions between the fermions and the Higgs and Goldstone bosons of the Standard Model. The corresponding Lagrangian \mathcal{L}_{EQT} is given, and is formally renormalized to all orders. The renormalization conditions are formulated both to make connection with physical observables and to satisfy the requirements underlying the equivalence theorem. As an application of this framework, we calculate the dominant radiative corrections to fermionic Higgs decays at one loop including the virtual effects of a heavy top quark. We apply the result to the decays $H \rightarrow t\bar{t}$ and $H \rightarrow b\bar{b}$, and find that the equivalence theorem results including fermions are quite accurate numerically for Higgs-boson masses $M_H > 400$ (350) GeV, respectively, even for $m_t = 175$ GeV.

*Electronic address: ldurand@wishep.physics.wisc.edu

†Electronic address: kurtr@physik.tu-muenchen.de

1 Introduction

The physics of the Higgs-sector is one of the most important subjects of high-energy experiments in the near future. To distinguish the Standard Model (SM) Higgs-sector from other theoretical models, the calculation of radiative corrections to high-energy processes may be very important. In addition, the effects of a Higgs boson on low-energy observables such as the ρ -parameter already need to be included in present precision measurements. However, the calculation of the complete higher-order corrections is a difficult task. It is therefore important if one can calculate the most significant corrections relatively simply within a well-defined approximation. In the case of a heavy Higgs boson, such an approximation is provided by the equivalence theorem (EQT) [1, 2]. This theorem shows that the leading correction to a physical process in powers of M_H^2/v^2 can be obtained at any order by replacing the vector bosons W^\pm and Z of the SM by the associated scalar Goldstone bosons w^\pm, z which appear in the symmetry breaking sector of the theory in the limit of vanishing gauge couplings. This results in a substantial simplification of the calculations. The equivalence theorem is known to hold for $M_H \gg M_W, M_Z$. An important question, however, is how heavy the Higgs boson must be before the equivalence theorem gives a good approximation to the full electroweak theory.

The existence of a heavy top quark leads to a large Yukawa coupling g_t in the SM. In the case of a not-so-heavy Higgs boson, one can expect the radiative corrections which involve g_t to be comparable in magnitude to the corrections which involve the quartic Higgs coupling $\lambda = M_H^2/2v^2$. It is therefore desirable to take these contributions into account, and to calculate them in a simple way. This can again be done using the equivalence theorem: in the limit of vanishing gauge couplings, the fermions couple only to the Higgs boson and the Goldstone bosons in the symmetry breaking sector of the theory. However, the specific calculational scheme used has to be consistent both with the EQT as formulated for the bosons [1, 2], and with the complete SM. In the present paper we show how this can be achieved by extending the equivalence theorem, including fermions, and implementing consistent renormalization conditions. The resulting framework is a foundation for future calculations of leading higher-order electroweak corrections, and it is applicable to all orders in λ and g_f . Gauge interactions are explicitly set to zero.

Previous calculations based on the EQT have mostly been concerned with processes in which both the external and internal particles are Higgs bosons H or longitudinally polarized gauge bosons W_L and Z_L [1, 2, 3, 4, 5, 6, 7, 8, 9, 10], and have neglected corrections from internal fermion loops. An exception is the work of Barbieri *et al.* [11], who calculated the m_t^4 corrections to the decays $Z \rightarrow \mu^+\mu^-$ and $Z \rightarrow b\bar{b}$ using a method based on the equivalence theorem, but with the fermion fields renormalized only to one loop. The framework presented here allows the systematic inclusion of internal fermion loops at higher orders, as well as the calculation of radiative corrections to processes involving external fermions.

We illustrate our method at the one-loop level using the fermionic decay of the Higgs boson, $H \rightarrow f\bar{f}$. The equivalence theorem result approximates the full electroweak correction to $\Gamma(H \rightarrow t\bar{t})$ very well for $M_H > 400$ GeV, but fails near the threshold at 350 GeV for $m_t = 175$ GeV, where the effects of the gauge interactions become important. The result for $\Gamma(H \rightarrow b\bar{b})$ is also quite good for $M_H > 350$ GeV, though the accuracy is reduced by a cancellation between the Yukawa and Higgs-boson contributions. The gauge couplings give the dominant, very small, corrections for lower values of M_H .

2 The Lagrangian and renormalization scheme

2.1 Framework

We will be concerned in later sections with the calculation of the leading contributions to the decay rate of the Higgs boson to fermions, $H \rightarrow f\bar{f}$, in the limit of large Higgs-boson mass, $M_H \gg M_W$. Our tool for extracting the leading contributions in powers of M_H/M_W will be the Goldstone boson equivalence theorem [1, 2, 12, 13, 14, 15]. This theorem states that the dominant electroweak contribution to a Feynman graph can be calculated by replacing the gauge bosons W^\pm, Z of the full electroweak theory by the would-be Goldstone bosons w^\pm, z of the symmetry-breaking sector of the theory, and ignoring the gauge couplings. The result is an expansion for the dominant contributions in powers of $G_F M_H^2$, or equivalently, in powers of the quartic Higgs coupling λ . In the limit $g, g' \rightarrow 0$, the transverse gauge bosons decouple from the fermions and the scalar fields, and can be ignored. The remainder of the standard electroweak model reduces to a theory defined by the equivalence-theorem Lagrangian $\mathcal{L}_{EQT} = \mathcal{L}_H + \mathcal{L}_F$, where \mathcal{L}_H is the Lagrangian for the scalar fields in the symmetry-breaking sector, and \mathcal{L}_F is the Lagrangian for the fermions which includes their Yukawa interactions with the scalar fields. This reduced theory gives a good approximation to the full electroweak theory for $M_W^2 \ll M_H^2$, provided that the couplings are properly defined and the Goldstone modes are renormalized at a momentum scale $p^2 \ll M_H^2$ [13].

We will work entirely in the equivalence theorem limit using \mathcal{L}_{EQT} ,

$$\mathcal{L}_{EQT} = \mathcal{L}_H^0 + \mathcal{L}_F^0 + \text{counterterms}, \quad (1)$$

where

$$\mathcal{L}_H^0 = \frac{1}{2} (\partial_\mu \Phi)^\dagger (\partial^\mu \Phi) - \frac{1}{4} \lambda (\Phi^\dagger \Phi)^2 + \frac{1}{2} \mu^2 \Phi^\dagger \Phi, \quad (2)$$

$$\begin{aligned} \mathcal{L}_F^0 &= i \bar{\Psi}_L \not{\partial} \Psi_L + i \bar{\psi}_{t,R} \not{\partial} \psi_{t,R} + i \bar{\psi}_{b,R} \not{\partial} \psi_{b,R} \\ &\quad - \frac{1}{\sqrt{2}} g_t \bar{\Psi}_L \tilde{\Phi} \psi_{t,R} - \frac{1}{\sqrt{2}} g_b \bar{\Psi}_L \Phi \psi_{b,R} + \text{h.c.} + \dots \end{aligned} \quad (3)$$

We have written only the top- and bottom quark contributions to \mathcal{L}_F^0 ; the remaining fermionic contributions have the same form, with no righthanded contributions for the neutrinos. In these expressions, Ψ_L, Φ , and $\tilde{\Phi}$ are $SU(2)_L$ doublets with normalizations defined by

$$\Psi_L = \begin{pmatrix} \psi_{t,L} \\ \psi_{b,L} \end{pmatrix}, \quad \psi_{f,L} = \frac{1}{2} (1 - \gamma^5) \psi_f, \quad (4)$$

$$\Phi = \begin{pmatrix} w_1 + iw_2 \\ h + iz \end{pmatrix} = \begin{pmatrix} \sqrt{2} w^+ \\ h + iz \end{pmatrix}, \quad (5)$$

$$\tilde{\Phi} = i\sigma_2 \Phi^* = \begin{pmatrix} h - iz \\ -w_1 + iw_2 \end{pmatrix} = \begin{pmatrix} h - iz \\ -\sqrt{2} w^- \end{pmatrix}. \quad (6)$$

The righthanded fields are $SU(2)_L$ singlets, with

$$\psi_{f,R} = \frac{1}{2} (1 + \gamma^5) \psi_f. \quad (7)$$

The expressions in Eqs. (2) and (3) include all possible $SU(2)_L \times U(1)_Y$ symmetric terms consistent with renormalizability of the Lagrangian. The counterterms necessary to effect

the renormalization without breaking the symmetry must have the same forms, and can be introduced by independent multiplicative rescalings of each term above. Thus, in the case of the Higgs Lagrangian, all possible symmetric counterterms are generated by multiplying the kinetic energy term by a factor Z_Φ , replacing λ by $\lambda + \delta\lambda$, and μ^2 by $\mu^2 + \delta\mu^2$. Because the minimum in the Higgs potential is at a nonzero value of $\Phi^\dagger\Phi$ for $\mu^2 > 0$, Φ has a nonzero expectation value in the physical vacuum,

$$\langle \Omega | \Phi^\dagger \Phi | \Omega \rangle = v^2. \quad (8)$$

We will absorb the vacuum expectation value by rewriting the field h as $h \rightarrow H + v$. This results in an expansion around the physical vacuum in which

$$\langle \Omega | H | \Omega \rangle = \langle \Omega | \mathbf{w} | \Omega \rangle = 0. \quad (9)$$

The renormalized Higgs Lagrangian then has the form

$$\begin{aligned} \mathcal{L}_H &= \frac{1}{2} Z_\Phi (\partial_\mu \mathbf{w} \cdot \partial^\mu \mathbf{w} + \partial_\mu H \partial^\mu H) - \frac{1}{4} (\lambda + \delta\lambda) (\mathbf{w}^2 + H^2 + 2vH)^2 \\ &\quad - \frac{1}{2} ((\lambda + \delta\lambda) v^2 - \mu^2 - \delta\mu^2) (\mathbf{w}^2 + H^2 + 2vH), \end{aligned} \quad (10)$$

where \mathbf{w} is the SO(3) vector (w_1, w_2, w_3) with $w_3 = z$, and we have dropped an additive constant.

It is convenient for calculation to define a set of rescaled or “bare” fields H_0 , \mathbf{w}_0 and the corresponding SU(2)_L doublet Φ_0 such that the kinetic terms in \mathcal{L}_H have the customary unit normalization,

$$H_0 = Z_\Phi^{1/2} H, \quad \mathbf{w}_0 = Z_\Phi^{1/2} \mathbf{w}, \quad \Phi_0 = Z_\Phi^{1/2} \Phi, \quad (11)$$

and to introduce a corresponding bare vacuum expectation value v_0 , a bare coupling λ_0 , and a parameter δm_0^2 ,

$$v_0 = Z_\Phi^{1/2} v, \quad (12)$$

$$\lambda_0 = \frac{\lambda}{Z_\Phi^2} \left(1 + \frac{\delta\lambda}{\lambda} \right), \quad (13)$$

$$\delta m_0^2 = Z_\Phi^{-1} (\mu^2 + \delta\mu^2 - (\lambda + \delta\lambda) v^2). \quad (14)$$

With this rescaling, we obtain the form of \mathcal{L}_H that we will use,

$$\begin{aligned} \mathcal{L}_H &= \frac{1}{2} (\partial_\mu \mathbf{w}_0 \cdot \partial^\mu \mathbf{w}_0 + \partial_\mu H_0 \partial^\mu H_0) - \frac{1}{4} \lambda_0 (\mathbf{w}_0^2 + H_0^2 + 2v_0 H_0)^2 \\ &\quad + \frac{1}{2} \delta m_0^2 (\mathbf{w}_0^2 + H_0^2 + 2v_0 H_0). \end{aligned} \quad (15)$$

The fermion Lagrangian \mathcal{L}_F can be treated in a similar fashion. There is a separate SU(2)_L × U(1)_Y-symmetric counterterm for each term in Eq. (3). These counterterms can be absorbed in the definitions of a set of bare fields and couplings to bring \mathcal{L}_F into the form

$$\begin{aligned} \mathcal{L}_F &= i\bar{\Psi}_L^0 \not{\partial} \Psi_L^0 + i\bar{\psi}_{t,R}^0 \not{\partial} \psi_{t,R}^0 + i\bar{\psi}_{b,R}^0 \not{\partial} \psi_{b,R}^0 \\ &\quad - \frac{1}{\sqrt{2}} g_t^0 \bar{\Psi}_L^0 \tilde{\Phi}_0 \psi_{t,R}^0 - \frac{1}{\sqrt{2}} g_b^0 \bar{\Psi}_L^0 \Phi_0 \psi_{b,R}^0 + \text{h.c.}, \end{aligned} \quad (16)$$

where

$$\Psi_L^0 = Z_L^{1/2} \Psi_L, \quad (17)$$

$$\psi_{f,R}^0 = Z_{f,R}^{1/2} \psi_{f,R}, \quad f = t, b, \quad (18)$$

$$g_f^0 = \frac{g_f}{Z_\Phi^{1/2}} \left(1 + \frac{\delta g_f}{g_f} \right). \quad (19)$$

Note that there are separate renormalization constants for the $\psi_{t,R}$ and $\psi_{b,R}$, and a single renormalization constant for the left-handed doublet Ψ_L [16]. The factor $Z_\Phi^{-1/2}$ in g_f^0 has been introduced for later convenience.

2.2 Fixing the renormalization of the Higgs Lagrangian

In order that the theory defined by \mathcal{L}_{EQT} correspond to the equivalence theorem limit of the standard electroweak model, the definitions of the couplings and the renormalization scheme must be chosen to be consistent with that limit. Such a choice is not automatic. It is necessary that: (i), the w^\pm and z fields be renormalized at a momentum scale p^2 with $|p^2| \ll M_H^2$ [13]; and (ii), the couplings be defined in terms of physical standard-model quantities which have well-defined limits for $g, g' \rightarrow 0$. In the following, we will use an on-mass-shell renormalization scheme, define the quartic Higgs coupling in terms of the Fermi constant G_F and the physical Higgs-boson mass M_H , and relate the Yukawa couplings to the physical masses of the fermions.

The w^\pm and z bosons are guaranteed to be massless by the Goldstone theorem [17]. We will therefore renormalize the w_0^\pm and z_0 fields at $p^2 = 0$, thus satisfying condition (i), and will renormalize the Higgs field at $p^2 = M_H^2$. The quartic coupling λ will be defined [18] to be given *exactly* by the relation

$$\lambda = M_H^2/2v^2 = G_F M_H^2/\sqrt{2}, \quad (20)$$

where G_F is the Fermi constant obtained from the muon decay rate using the standard electromagnetic radiative corrections, and v is the physical vacuum expectation value, $v = 2^{-1/4} G_F^{-1/2}$. While this definition involves a process at a low momentum transfer, it connects smoothly with equivalence theorem limit as shown in [18, 19].

The determination of the renormalization constants proceeds as follows. The real part of the two-point function or inverse propagator $\Gamma^{(2)}(p^2)$ for each of the particles must vanish for p^2 equal to the square of the physical mass of that particle, $\Gamma^{(2)}(m^2) = 0$. The two-point functions calculated using the bare fields are easily seen to have the form

$$\Gamma_{w_0}^{(2)}(p^2) = p^2 - \Pi_w^0(p^2) + \delta m_0^2, \quad (21)$$

$$\Gamma_{z_0}^{(2)}(p^2) = p^2 - \Pi_z^0(p^2) + \delta m_0^2, \quad (22)$$

$$\Gamma_{H_0}^{(2)}(p^2) = p^2 - \Pi_H^0(p^2) - 2\lambda_0 v_0^2 + \delta m_0^2, \quad (23)$$

where the Π 's are the bare self-energy functions. Since there is only a single mass counterterm δm_0^2 in the Higgs Lagrangian, Eq. (15), the vanishing of the renormalized masses m_w and m_z required by the Goldstone theorem leads to the relations

$$\delta m_0^2 = \Pi_w^0(0) = \Pi_z^0(0), \quad (24)$$

and determines δm_0^2 . The self-energy functions for the bare w^\pm and z fields must therefore be equal at $p^2 = 0$, an identity that holds to all orders in perturbation theory. This would not be surprising in the absence of Yukawa couplings since the Goldstone fields then have an SO(3) symmetry, and $\Pi_w^0(p^2)$ and $\Pi_z^0(p^2)$ are necessarily identical for all values of p^2 . However, the presence of fermions with unequal Yukawa couplings in \mathcal{L}_{EQT} breaks the SO(3) symmetry, and destroys the equality of the self-energy functions except at the special point $p^2 = 0$ [20]. The equality $\Pi_w^0(0) = \Pi_z^0(0)$ can be checked to one loop using the results we give later. In the following, we will replace δm_0^2 by $\Pi_w^0(0)$.

The bare coupling λ_0 can be determined by using the definition $\lambda = M_H^2/2v^2$ and Eq. (23). The on-mass-shell condition for the Higgs boson, $\Gamma_{H_0}^{(2)}(M_H^2) = 0$, gives the relation

$$M_H^2 = 2\lambda_0 v_0 + \Pi_H^0(M_H^2) - \delta m_0^2. \quad (25)$$

Upon replacing v_0^2 by $Z_\Phi v^2 = Z_\Phi M_H^2/2\lambda$ and δm_0^2 by $\Pi_w^0(0)$ in this expression and solving for λ_0 , we find that [21]

$$\lambda_0 = \lambda \left(1 + \frac{\delta\lambda}{\lambda}\right) \frac{1}{Z_\Phi^2} = \frac{\lambda}{Z_\Phi} \left(1 - \frac{\text{Re}\Pi_H^0(M_H^2) - \Pi_w^0(0)}{M_H^2}\right). \quad (26)$$

The condition that v be the physical vacuum expectation value requires the vanishing of the truncated one-point function for the Higgs field,

$$\Gamma_{H_0}^{(1)}(0) = -iT_0 + iv_0\delta m_0^2 = 0, \quad (27)$$

where T_0 is the sum of all Higgs tadpole graphs (see Fig. 1) calculated using the bare fields. This gives the further relation $\delta m_0^2 = T_0/v_0$, so δm_0^2 can be calculated either as a self energy or as a tadpole contribution. The resulting identity,

$$\Pi_w^0(0) = T_0/v_0, \quad (28)$$

provides a useful check on the calculations [22]. The vanishing of the one-point function, Eq. (27), implies that the tadpole diagrams and the tadpole counterterm in Eq. (15) cancel order-by-order in the perturbation expansion and can be dropped together, as discussed by Taylor [23]. We will adopt this simplification in the following calculations.

Finally, the wave function renormalization constants Z_w , Z_z , and Z_H which relate the bare fields w_0^\pm , z_0 , H_0 to physical fields,

$$w_0^\pm = Z_w^{1/2} w_{\text{phys}}^\pm, \quad z_0 = Z_z^{1/2} z_{\text{phys}}, \quad H_0 = Z_H^{1/2} H_{\text{phys}}, \quad (29)$$

are determined by the condition that the propagators for the physical fields have unit residue at the particle poles. The bare propagators do not, but instead have residues Z_i given by

$$Z_i = \left(\frac{d}{dp^2} \Gamma_i^{(2)}(p^2) \Big|_{p^2=m_i^2} \right)^{-1}. \quad (30)$$

Thus, using Eqs. (21-23),

$$\frac{1}{Z_w} = 1 - \frac{d}{dp^2} \Pi_w^0(p^2) \Big|_{p^2=0}, \quad (31)$$

$$\frac{1}{Z_z} = 1 - \frac{d}{dp^2} \Pi_z^0(p^2) \Big|_{p^2=0}, \quad (32)$$

$$\frac{1}{Z_H} = 1 - \frac{d}{dp^2} \Pi_H^0(p^2) \Big|_{p^2=M_H^2}. \quad (33)$$

In the presence of fermions with different Yukawa couplings or masses, e.g., top and bottom quarks with $m_t \neq m_b$, the self-energy functions Π_w^0 and Π_z^0 are not equal except at the isolated point $p^2 = 0$, and, as a result, $Z_z \neq Z_w$.

The single wave function renormalization constant Z_Φ introduced earlier is sufficient, along with the other renormalization constants, to absorb the divergences in the Higgs sector of the theory. The constants Z_w , Z_z , and Z_H can therefore differ from Z_Φ only by finite multiplicative factors \tilde{Z}_i ,

$$Z_i = \tilde{Z}_i Z_\Phi. \quad (34)$$

The problem which remains is that of determining Z_Φ , an essential step in connecting the scalar theory to the full electroweak theory through the equivalence theorem. The crucial observation is that the Fermi constant G_F , and therefore the vacuum expectation value v , is defined through charged-current processes, i.e., muon decay and superallowed nuclear beta decays. These involve the W rather than the Z boson. The connection can then be made using the Ward identities for the electroweak charged current which underlie the equivalence theorem [13, 11]. In particular, Bagger and Schmidt [13] show that W^\pm scattering amplitudes calculated in the full electroweak theory are related to those calculated using the equivalence theorem by powers of the ratio [24]

$$C = \frac{M_W^0}{M_W} \frac{Z_W^{1/2}}{Z_w^{1/2}} = \frac{g_0 v_0}{g v} \frac{Z_W^{1/2}}{Z_w^{1/2}} = \frac{Z_\Phi^{1/2}}{Z_w^{1/2}} (1 + O(g^2)). \quad (35)$$

They then use the Ward identity to establish that $C = 1$ up to corrections of order g^2 under conditions satisfied by our renormalization scheme. Thus, $Z_\Phi = Z_w$ and $\tilde{Z}_w = 1$ in the limit $g, g' \rightarrow 0$.

2.3 Fixing the renormalization of the fermion Lagrangian

When written with the $SU(2)_L$ spinor products expanded, the fermion Lagrangian \mathcal{L}_F in Eq. (16) assumes the form

$$\begin{aligned} \mathcal{L}_F = & i\bar{\psi}_t^0 \not{\partial} \psi_t^0 - (m_t + \delta m_t) \bar{\psi}_t^0 \psi_t^0 - i\bar{\psi}_b^0 \not{\partial} \psi_b^0 - (m_b + \delta m_b) \bar{\psi}_b^0 \psi_b^0 \\ & - \frac{1}{\sqrt{2}} g_t^0 \bar{\psi}_t^0 H_0 \psi_t^0 + \frac{i}{\sqrt{2}} g_t^0 \bar{\psi}_t^0 \gamma^5 z_0 \psi_t^0 - \frac{1}{\sqrt{2}} g_b^0 \bar{\psi}_b^0 H_0 \psi_b^0 - \frac{i}{\sqrt{2}} g_b^0 \bar{\psi}_b^0 \gamma^5 z_0 \psi_b^0 \\ & + g_t^0 \bar{\psi}_{t,R}^0 w_0^+ \psi_{b,L}^0 + g_t^0 \bar{\psi}_{b,L}^0 w_0^- \psi_{t,R}^0 - g_b^0 \bar{\psi}_{b,R}^0 w_0^+ \psi_{t,L}^0 - g_b^0 \bar{\psi}_{t,L}^0 w_0^- \psi_{b,R}^0, \end{aligned} \quad (36)$$

where we have used the relations in Eqs. (11), (12), and (19). The parameters m_f and g_f are defined to be the physical masses and Yukawa couplings of the quarks while δm_f is the mass or coupling counterterm,

$$m_f = g_f \frac{v}{\sqrt{2}}, \quad \delta m_f = m_f \frac{\delta g_f}{g_f}. \quad (37)$$

Our calculations will be carried out using the bare fields and the standard free propagators for the quarks, and yield corrected fermion propagators iS_F^0 of the form

$$iS_F^0 = \frac{i}{\not{p} - m_f - \Sigma_f^0 - \delta m}. \quad (38)$$

The self-energy Σ_f^0 contains an axial vector component as well as the vector and scalar components familiar in parity-conserving theories. In particular, suppressing the quark label,

$$\Sigma^0 = \not{p} \Pi_V^0(p^2) + \not{p} \gamma^5 \Pi_A^0(p^2) + m \Pi_S^0(p^2). \quad (39)$$

The physical propagators iS_F are related to the bare propagators through the connection between the physical and bare fields,

$$\psi = \psi_R + \psi_L = \left(Z_R^{-1/2} P_R + Z_L^{-1/2} P_L \right) \psi^0, \quad (40)$$

where $P_R = \frac{1}{2}(1 + \gamma^5)$ and $P_L = \frac{1}{2}(1 - \gamma^5)$. Recalling the definition of the propagator, $iS_F(x - y) = \langle \Omega | T(\psi(x), \bar{\psi}(y)) | \Omega \rangle$, we find that

$$iS_F(p) = \left(Z_R^{-1/2} P_R + Z_L^{-1/2} P_L \right) iS_F^0(p) \left(Z_R^{-1/2} P_L + Z_L^{-1/2} P_R \right), \quad (41)$$

or, calculating the inverse of S_F^0 explicitly,

$$\begin{aligned} iS_F(p) &= \left\{ \not{p} \left[\frac{1}{2} (Z_L^{-1} + Z_R^{-1}) (1 - \Pi_V^0(p^2)) - \frac{1}{2} (Z_L^{-1} - Z_R^{-1}) \Pi_A^0(p^2) \right] \right. \\ &\quad \left. + \not{p} \gamma^5 \left[\frac{1}{2} (Z_L^{-1} - Z_R^{-1}) (1 - \Pi_V^0(p^2)) - \frac{1}{2} (Z_L^{-1} + Z_R^{-1}) \Pi_A^0(p^2) \right] \right. \\ &\quad \left. + Z_L^{-1/2} Z_R^{-1/2} (m + \delta m + m \Pi_S^0(p^2)) \right\} \\ &\quad \times \left\{ p^2 \left[(1 - \Pi_V^0(p^2))^2 - \Pi_A^0(p^2) \right] - (m + \delta m + m \Pi_S^0(p^2))^2 \right\}^{-1}. \end{aligned} \quad (42)$$

The mass or coupling counterterm $\delta m = m \delta g/g$ and the wave function renormalization constants Z_R and Z_L are determined by the condition that the physical propagator describe a freely propagating particle with a mass m and unit residue at the particle pole, $S_F \rightarrow 1/(\not{p} - m) = (\not{p} + m)/(p^2 - m^2)$ for $\not{p} \rightarrow m$. S_F has a simple pole at $p^2 = m^2$ provided that [25]

$$1 + \frac{\delta m}{m} = 1 + \frac{\delta g}{g} = \sqrt{[1 - \Pi_V^0(m^2)]^2 - \Pi_A^0(m^2)} - \Pi_S^0(m^2). \quad (43)$$

The coefficient of $\not{p} \gamma^5$ in Eq. (42) must vanish if S_F is to be a normal massive propagator with equal right- and left-handed residues at the particle pole. This requires that

$$(Z_R - Z_L) [1 - \Pi_V^0(m^2)] = (Z_R + Z_L) \Pi_A^0(m^2). \quad (44)$$

The coefficients of \not{p} and m in the numerator will be equal if, in addition,

$$\frac{1}{2} (Z_R + Z_L) [1 - \Pi_V^0(m^2)] - \frac{1}{2} (Z_R - Z_L) \Pi_A^0(m^2) = Z_R^{1/2} Z_L^{1/2} \left[1 + \frac{\delta g}{g} + \Pi_S^0(m^2) \right]. \quad (45)$$

Equations (44) and (45) are homogeneous of degree one in the Z 's, and only determine the ratio Z_R/Z_L . The two equations may also be combined to obtain the pole condition, Eq. (43). The magnitude of the Z 's is determined by the final condition, that S_F have unit residue at the pole. With the ratios Z_R/Z_L determined as above for both the quarks in a doublet, this final condition can only be enforced for one of the two quark propagators by adjusting the single remaining renormalization constant Z_L for the left handed fields Ψ_L , Eq. (17). The second quark field requires an additional wave function renormalization to reach the standard normalization. This extra renormalization constant is finite; the infinities in the wave function renormalizations can be absorbed entirely in Z_L and the two Z_R 's, which generate the only symmetric counterterms allowed in the kinetic part of \mathcal{L}_F , Eq. (16). Since our calculations are done entirely in terms of the bare fields, with the Z 's appearing only in the overall factors necessary for external particles, the choice of definition for the original Z_L is irrelevant, and we will give only the final renormalization constants. These are determined by setting the ratio of the coefficient of \not{p} in Eq. (42) to the derivative of the denominator equal to unity for $p^2 = m^2$. Using Eqs. (43-45), we find that

$$\begin{aligned} \frac{1}{Z_L} &= 1 - \Pi_V^0(m^2) + \Pi_A^0(m^2) - 2m^2\Pi_S^{\prime 0}(m^2) \sqrt{\frac{1 - \Pi_V^0(m^2) + \Pi_A^0(m^2)}{1 - \Pi_V^0(m^2) - \Pi_A^0(m^2)}} \\ &\quad - 2m^2 \frac{(1 - \Pi_V^0(m^2)) \Pi_V^{\prime 0}(m^2) - \Pi_A^0(m^2) \Pi_A^{\prime 0}(m^2)}{1 - \Pi_V^0(m^2) - \Pi_A^0(m^2)} \end{aligned} \quad (46)$$

and

$$\begin{aligned} \frac{1}{Z_R} &= 1 - \Pi_V^0(m^2) - \Pi_A^0(m^2) - 2m^2\Pi_S^{\prime 0}(m^2) \sqrt{\frac{1 - \Pi_V^0(m^2) - \Pi_A^0(m^2)}{1 - \Pi_V^0(m^2) + \Pi_A^0(m^2)}} \\ &\quad - 2m^2 \frac{(1 - \Pi_V^0(m^2)) \Pi_V^{\prime 0}(m^2) - \Pi_A^0(m^2) \Pi_A^{\prime 0}(m^2)}{1 - \Pi_V^0(m^2) - \Pi_A^0(m^2)}, \end{aligned} \quad (47)$$

where $\Pi'(p^2) = d\Pi(p^2)/dp^2$. These expressions hold to all orders in perturbation theory, with one set for each quark. At the one-loop level [16] of interest here, they simplify substantially to

$$Z_L = 1 + \Pi_V^0(m^2) - \Pi_A^0(m^2) + 2m^2\Pi_V^{\prime 0}(m^2) + 2m^2\Pi_S^{\prime 0}(m^2), \quad (48)$$

$$Z_R = 1 + \Pi_V^0(m^2) + \Pi_A^0(m^2) + 2m^2\Pi_V^{\prime 0}(m^2) + 2m^2\Pi_S^{\prime 0}(m^2), \quad (49)$$

$$\frac{\delta g}{g} = \frac{\delta m}{m} = -\Pi_V^0(m^2) - \Pi_S^0(m^2). \quad (50)$$

It is straightforward, finally, to establish the renormalization factors which must be used for the external fermions when scattering amplitudes are calculated using the bare fields. By using the standard reduction formulas [26], we can express the physical scattering amplitudes in terms of Fourier transforms of vacuum expectation values of time-ordered products of the physical fields. These vacuum expectation values appear multiplied by a factor $(\not{p} - m)$ and a spinor for each ingoing or outgoing fermion line. The physical fermion fields ψ , $\bar{\psi}$ in the time-ordered products can be replaced by the bare fields ψ^0 , $\bar{\psi}^0$ using the definition in Eq. (40). When the result is reexpressed in terms of the bare truncated Green's function Γ_n^0 , the

external factors are replaced by $S_F^0(p)(Z_R^{-1/2}P_L + Z_L^{-1/2}P_R)(\not{p} - m)$ and a spinor for each ingoing fermion line, and by $(\not{p} - m)(Z_R^{-1/2}P_R + Z_L^{-1/2}P_L)S_F^0(p)$ and a conjugate spinor for each outgoing line. However, it follows from the first line of Eq. (42) and the normalization of the physical propagator that

$$S_F^0(p)(Z_R^{-1/2}P_L + Z_L^{-1/2}P_R) = (Z_R^{1/2}P_R + Z_L^{1/2}P_L)S_F(p), \quad (51)$$

$$(Z_R^{-1/2}P_R + Z_L^{-1/2}P_L)S_F^0(p) = S_F(p)(Z_R^{1/2}P_L + Z_L^{1/2}P_R), \quad (52)$$

and that

$$S_F(p)(\not{p} - m) = (\not{p} - m)S_F(p) = 1, \quad \not{p} = m. \quad (53)$$

As a result, the physical scattering amplitudes are given by the truncated Green's functions Γ_n^0 calculated in terms of the bare fields, multiplied by a factor $(Z_R^{1/2}P_R + Z_L^{1/2}P_L)$ and the appropriate spinor for each incoming fermion line, and by a factor $(Z_R^{1/2}P_L + Z_L^{1/2}P_R)$ and a conjugate spinor for each outgoing fermion line. These factors generalize the usual factors of $Z^{1/2}$ for standard Dirac fields to the case of chiral interactions.

3 $H \rightarrow f\bar{f}$ to one loop

In the present section, we will sketch the calculation of the one-loop corrections to the matrix element for the decay $H \rightarrow f\bar{f}$ using the equivalence theorem, and compare this approximate result, valid for $M_H \gg M_W$ independently of the fermion masses, with the exact result obtained by other authors [27, 28, 29, 30].

3.1 Form of the decay matrix element

According to the discussion above, the matrix element for the decay $H \rightarrow f\bar{f}$ is given by the expression

$$-i\mathcal{M}_{H \rightarrow f\bar{f}} = Z_H^{1/2}\bar{u}(p_1 - p_2, m_f) \left(Z_R^{1/2}P_L + Z_L^{1/2}P_R \right) \Gamma_3^0 \left(Z_R^{1/2}P_R + Z_L^{1/2}P_L \right) v(p_2, m_f). \quad (54)$$

Here p_1 and p_2 are the momenta of the incoming Higgs boson and the outgoing antifermion, respectively, and Γ_3^0 is the truncated three-point vertex function calculated using the bare Lagrangian,

$$\Gamma_3^0 = -\frac{i}{\sqrt{2}}g_f^0 + \sum_{i=1}^6 L_i + \dots, \quad (55)$$

where the one-loop integrals L_i correspond to the triangle diagrams in Fig. 2.

To one-loop, the renormalization constants multiply only the leading term in Γ_3^0 . Using the definition of g_f^0 in Eq. (19) with $Z_\Phi = Z_w$ as established above, writing Z_i as $1 + \delta Z_i$, and expanding, we obtain the one-loop expression for $\mathcal{M}_{H \rightarrow f\bar{f}}$,

$$\begin{aligned} -i\mathcal{M}_{H \rightarrow f\bar{f}} &= \bar{u}(p_1 - p_2, m_f) \left[-\frac{i}{\sqrt{2}}g_f^0 \left(1 + \frac{1}{2}\delta Z_H - \frac{1}{2}\delta Z_w + \frac{1}{2}\delta Z_L^f + \frac{1}{2}\delta Z_R^f + \frac{\delta g_f}{g_f} \right) \right. \\ &\quad \left. + \sum_{i=1}^6 L_i \right] v(p_2, m_f). \end{aligned} \quad (56)$$

The “ δ ” contributions in this expression would appear as renormalization counterterms in a calculation based on physical rather than bare fields and, for convenience, we will refer to them as such.

It will be useful to split the counterterms into “bosonic” and “fermionic” parts, and rewrite the decay matrix element to one loop as

$$-i\mathcal{M}_{H\rightarrow f\bar{f}} = \bar{u}(p_1 - p_2, m_f) \left[-\frac{i}{\sqrt{2}}g_f(1 + L_{bos} + L_{fer}) + \sum_{i=1}^6 L_i \right] v(p_2, m_f). \quad (57)$$

where

$$L_{bos} = \frac{Z_H^{1/2}}{Z_w^{1/2}} - 1 = \frac{1}{2}\delta Z_H - \frac{1}{2}\delta Z_w + \dots, \quad (58)$$

and

$$L_{fer} = \left(1 + \frac{\delta g_f}{g_f}\right) \left(Z_R^f Z_L^f\right)^{1/2} - 1 = \frac{1}{2}\delta Z_R^f + \frac{1}{2}\delta Z_L^f + \frac{\delta g_f}{g_f} + \dots. \quad (59)$$

This grouping of terms has the advantage that L_{bos} , while dependent on the fermions through loop contributions, is independent of the flavor of the final fermion pair. It therefore gives a universal correction to all decays $H \rightarrow f\bar{f}$. The fermionic counterterm, in contrast, depends explicitly on the flavor of the final pair. This distinction will be important later.

3.2 The bosonic counterterm

The bosonic counterterm L_{bos} defined in Eq. (58) is determined by the derivatives of the self-energies Π_w^0 and Π_H^0 in Eqs. (31) and (33). Because the field renormalization constant Z_H differs from $Z_w = Z_\Phi$ only by a finite renormalization, the bosonic counterterm L_{bos} is finite to all orders. The tadpole contributions to the self energies are cancelled by the same counterterm $T_0 = v_0\delta m_0^2$ as cancels the apparent radiative changes in the physical vacuum expectation value v , and will be dropped. The remaining contributions to the boson self energies are given to one loop by the one-particle irreducible diagrams shown in Figs. 3 and 4. The presence of fermion loops in these diagrams leads to a breaking of the $SO(3)$ symmetry of the Higgs Lagrangian for unequal masses of the fermions in an $SU(2)_L$ doublet, with the result that $\Pi_z^0(p^2) \neq \Pi_w(p^2)$ for $p^2 \neq 0$, hence that $Z_z \neq Z_w$.

The boson-loop diagrams have been calculated by a number of authors [3, 4, 5, 6, 7, 31, 19, 32]. With the notation

$$\delta Z_i = \delta Z_{i,bos} + \delta Z_{i,fer}, \quad (60)$$

the results are

$$\delta Z_{H,bos} = \frac{\lambda}{16\pi^2} (12 - 2\pi\sqrt{3}), \quad (61)$$

$$\delta Z_{w,bos} = \frac{\lambda}{16\pi^2} (-1). \quad (62)$$

These one-loop quantities are separately finite. At two loops, the Z 's become singular [19, 33, 34], but the ratio $Z_{H,bos}/Z_{w,bos}$ and hence its contribution to L_{bos} remains finite.

The contributions to δZ_H and δZ_w from the fermionic loops are [35]

$$\delta Z_{H,fer} = \frac{1}{8\pi^2 v^2} \sum_f N_C^f m_f^2 \text{Re} \left[-B_0(M_H^2, m_f^2, m_f^2) - (M_H^2 - 4m_f^2) \partial B_0(M_H^2, m_f^2, m_f^2) \right], \quad (63)$$

$$\delta Z_{w,fer} = \frac{1}{8\pi^2 v^2} \sum_{(f,f')} N_C^f \text{Re} \left[- (m_f^2 + m_{f'}^2) B_0(0, m_f^2, m_{f'}^2) + (m_f^2 - m_{f'}^2)^2 \partial B_0(0, m_f^2, m_{f'}^2) \right], \quad (64)$$

where the standard scalar integral $B_0(p^2, m_0^2, m_1^2)$ and its derivative $\partial B_0(p^2, m_0^2, m_1^2)$ [32, 36] are defined in Appendix B. In these expressions, f' is the doublet partner of the fermion f . The color factor N_C^f is 3 for quarks and 1 for leptons. While the function $B_0(p^2, m_0^2, m_1^2)$ and the individual δZ 's are ultraviolet divergent, the divergencies cancel in the difference $\delta Z_{H,fer} - \delta Z_{w,fer}$, and the fermionic contribution to L_{bos} is also finite, as expected.

Because the overall coupling factors are proportional to m_f^2/v^2 , light or massless fermions do not contribute significantly to the radiative corrections. However, the contributions from top-quark loops can be significant, especially if $M_H \approx O(m_t)$. We will extract these corrections explicitly. Neglecting all fermion masses except for m_t , the fermionic contribution to L_{bos} is

$$\frac{1}{2} \delta Z_{H,fer} - \frac{1}{2} \delta Z_{w,fer} = \frac{1}{16\pi^2} N_C^f \frac{m_t^2}{v^2} \left[-\text{Re} \left[B_0(M_H^2, m_t^2, m_t^2) - B_0(0, 0, m_t^2) \right] - \left(1 - \frac{1}{a^2} \right) M_H^2 \text{Re} \left[\partial B_0(M_H^2, m_t^2, m_t^2) \right] - \frac{1}{2} \right] \quad (65)$$

The explicit expression for L_{bos} depends on the ratio of the Higgs-boson and top-quark masses. For Higgs-boson masses above the top-quark production threshold at $M_H = 2m_t$, we find that

$$L_{bos} = \frac{\lambda}{16\pi^2} \left(\frac{13}{2} - \pi\sqrt{3} \right) + \frac{3}{16\pi^2} \frac{m_t^2}{v^2} \left[b \left(2 + \frac{1}{a^2} \right) \left(\ln 2a + \ln \left(\frac{1+b}{2} \right) \right) - \frac{1}{2} - \frac{1}{a^2} \right], \quad M_H > 2m_t, \quad (66)$$

while for smaller masses, $M_H < 2m_t$,

$$L_{bos} = \frac{\lambda}{16\pi^2} \left(\frac{13}{2} - \pi\sqrt{3} \right) + \frac{3}{16\pi^2} \frac{m_t^2}{v^2} \left[\frac{1}{2a^4} \phi \sin(\phi) - \left(2a^2 - 3 + \frac{1}{a^2} \right) \frac{\phi}{\sin(\phi)} - \frac{1}{2} - \frac{1}{a^2} \right], \quad M_H < 2m_t, \quad (67)$$

where

$$a = M_H/2m_t, \quad b = \sqrt{1 - a^{-2}}, \quad \phi = \arccos(1 - 2a^2), \quad 0 < \phi < \pi. \quad (68)$$

We will henceforth refer to the cases $M_H > 2m_t$ ($M_H < 2m_t$) as those of a “heavy” (“light”) Higgs boson.

As noted earlier, the result for L_{bos} does not depend on the final state of the decay. For a fixed value of M_H , it is the same for all processes $H \rightarrow f\bar{f}$. As an useful example, we note that L_{bos} already gives the *complete* one-loop $\mathcal{O}(\lambda)$ and $\mathcal{O}(g_t^2)$ corrections to the dominant leptonic decay, $H \rightarrow \tau^+\tau^-$. The fermionic counterterm and the vertex corrections for this and other leptonic decays do not receive further corrections in these couplings.

It is worth mentioning that there are no threshold singularities in L_{bos} . The right-hand sides of Eqs. (66) and (67) are equal for $M_H = 2m_t$. In general, threshold singularities occur only when the gauge couplings of the Standard Model are included. We can consistently neglect the gauge couplings when using the equivalence theorem.

3.3 Fermionic counterterm

The fermionic counterterm L_{fer} defined in Eq. (59) can easily be reduced to the form

$$L_{fer} = \text{Re} \left\{ \Pi_{S,f}^0(m_f^2) - 2m_f^2 \left[\Pi_{V,f}^{0'}(m_f^2) + \Pi_{S,f}^{0'}(m_f^2) \right] \right\} \quad (69)$$

by using the renormalization conditions stated in Eqs. (48), (49), and (50). The potential tadpole contributions to the fermionic self energies and their derivatives are exactly cancelled by the counterterms. The remaining one-particle irreducible diagrams are shown in Fig. 5. All involve loops containing either a virtual Higgs boson or a massless Goldstone boson and the appropriate fermion line, f or f' , where f refers to the final fermion.

In contrast to the expressions for the bosonic self-energies, the expressions for the fermion self-energies depend on the Dirac matrices and, in particular, on γ^5 . This γ^5 dependence raises the possibility of problems with dimensional regularization related to the definition of such quantities as $\text{Tr}\gamma^5\gamma^{\mu_1}\dots\gamma^{\mu_D}$ and the antisymmetric tensor in D dimensions. However, as shown by Barbieri *et al.* [11], naive dimensional regularization with γ^5 and the remaining Dirac matrices treated as anticommuting is equivalent in the present context to the proper 't Hooft-Veltman scheme [36] to at least two loops for physical quantities. We have therefore calculated the fermion self energies and the triangle diagrams using naive dimensional regularization. Using the results in the appendices, we find that

$$\begin{aligned} L_{fer} = \frac{1}{16\pi^2} \frac{m_f^2}{v^2} \text{Re} \left\{ \left[B_0(m_f^2, m_f^2, M_H^2) - B_0(m_f^2, m_f^2, 0) - \frac{2m_{f'}^2}{m_f^2} B_0(m_f^2, m_{f'}^2, 0) \right] \right. \\ \left. - 2m_f^2 \left[-\partial B_1(m_f^2, m_f^2, M_H^2) - \partial B_1(m_f^2, m_f^2, 0) - \left(1 + \frac{m_{f'}^2}{m_f^2} \right) \partial B_1(m_f^2, m_{f'}^2, 0) \right. \right. \\ \left. \left. + \partial B_0(m_f^2, m_f^2, M_H^2) - \partial B_0(m_f^2, m_f^2, 0) - \frac{2m_{f'}^2}{m_f^2} \partial B_0(m_f^2, m_{f'}^2, 0) \right] \right\}. \quad (70) \end{aligned}$$

The explicit evaluation of this expression requires the specification of the flavor of the final state fermion. The results for $f = t$ and $f = b$ are given in Sect. (3.5) and (3.6), respectively.

3.4 The one-loop vertex diagrams

The six one-loop vertex diagrams L_i , $i=1-6$, are shown in Fig. 2. Their contributions are given by

$$L_1 = \left(\frac{m_f}{v}\right)^3 T_I(p_1^2, p_2^2, m_f^2, m_f^2, M_H^2), \quad (71)$$

$$L_2 = -\left(\frac{m_f}{v}\right)^3 \gamma^5 T_I(p_1^2, p_2^2, m_f^2, m_f^2, 0) \gamma^5, \quad (72)$$

$$L_3 = \frac{2m_{f'}}{v^3} (m_f P_L - m_{f'} P_R) T_I(p_1^2, p_2^2, m_f^2, m_{f'}^2, 0) (P_R m_f - P_L m_{f'}), \quad (73)$$

$$L_4 = 6\lambda v \left(\frac{m_f}{v}\right)^2 T_{II}(p_1^2, p_2^2, m_f^2, m_f^2, M_H^2), \quad (74)$$

$$L_5 = -2\lambda v \left(\frac{m_f}{v}\right)^2 \gamma^5 T_{II}(p_1^2, p_2^2, 0, 0, m_f^2) \gamma^5, \quad (75)$$

$$L_6 = 2\lambda v \frac{1}{v^2} (m_f P_L - m_{f'} P_R) T_{II}(p_1^2, p_2^2, 0, 0, m_{f'}^2) (P_R m_f - P_L m_{f'}). \quad (76)$$

The functions T_I and T_{II} are defined by the the integrals

$$T_I(p_1^2, p_2^2, m^2, m^2, M^2) = (2\pi\mu)^{4-D} \int \frac{d^D q}{(2\pi)^4} \frac{\not{p}_1 \not{q} + m \not{p}_1 + \not{q} \not{q} + 2m\not{q} + m^2}{[q^2 - m^2 + i\epsilon'][(q+p_1)^2 - m^2 + i\epsilon'][(q+p_2)^2 - M^2 + i\epsilon]}, \quad (77)$$

$$T_{II}(p_1^2, p_2^2, m^2, m^2, M^2) = (2\pi\mu)^{4-D} \int \frac{d^D q}{(2\pi)^4} \frac{-\not{q} + \not{p}_2 + M}{[q^2 - m^2 + i\epsilon'][(q+p_1)^2 - m^2 + i\epsilon'][(q+p_2)^2 - M^2 + i\epsilon]}. \quad (78)$$

These tensor integrals can be evaluated using standard techniques [32, 37].

While the triangle functions L_i , $i=2, 3, 5, 6$ depend individually on γ^5 , the γ^5 's drop out in the sum of these four one-loop contributions. The decay matrix element $\mathcal{M}_{H \rightarrow f\bar{f}}$, Eq. (56), has no γ^5 terms. The remaining Dirac matrices can be eliminated from the expression for $\mathcal{M}_{H \rightarrow f\bar{f}}$ by using the Dirac equations for the spinors $\bar{u}(p_1 - p_2, m_f)$ and $v(p_2, m_f)$. Extracting a factor $-im_f/v$ from the reduced expressions for the functions L_i and denoting the results by \widetilde{L}_i , we can then write Eq. (57) as

$$-i\mathcal{M}_{H \rightarrow f\bar{f}} = -i\frac{m_f}{v} \bar{u}(p_1 - p_2, m_f) v(p_2, m_f) (1 + \Delta\mathcal{T}), \quad (79)$$

where the spinor matrix element is purely scalar and the quantity $\Delta\mathcal{T}$ is defined as

$$\Delta\mathcal{T} = L_{bos} + L_{fer} + L_{tri}, \quad L_{tri} = \sum_{i=1}^6 \widetilde{L}_i. \quad (80)$$

The correction to the decay width can then be written as

$$\begin{aligned} \Gamma(H \rightarrow f\bar{f}) &= \Gamma_B(H \rightarrow f\bar{f}) |1 + \Delta\mathcal{T}|^2 \\ &= \Gamma_B(H \rightarrow f\bar{f}) [1 + 2\text{Re}\Delta\mathcal{T} + O(\lambda^2, \lambda g_f^2, g_f^4)], \end{aligned} \quad (81)$$

where [38]

$$\Gamma_B (H \rightarrow f\bar{f}) = \frac{N_c m_f^2 M_H}{8\pi v^2} \left(1 - \frac{4m_f^2}{M_H^2}\right)^{3/2} \quad (82)$$

is the Born result. The function $\sum_i \tilde{L}_i$ is rather complicated, and we will only give the results needed for the decays $H \rightarrow t\bar{t}$ and $H \rightarrow b\bar{b}$. They are listed in the following sections.

3.5 $H \rightarrow t\bar{t}$

In the mass region $M_H > 2m_t$, the only important fermionic decay of the Higgs particle is the decay into a pair of top quarks, with a branching ratio of approximately 10% [39]. Since $m_t \gg m_b$, we can calculate the decay rate for $H \rightarrow t\bar{t}$ using our general results for $H \rightarrow f\bar{f}$ evaluated in the limit $m_b=0$. To simplify the expressions, we again use the notation

$$a = \frac{M_H}{2m_t} \quad b = \sqrt{1 - \frac{1}{a^2}}. \quad (83)$$

Using the relation $\lambda = M_H^2/(2v^2)$, we add the contributions from the six triangular loops and obtain

$$\begin{aligned} L_{tri} = & -\frac{1}{16\pi^2} \left(\frac{m_t}{v}\right)^2 \frac{4a^4}{a^2-1} \left[\left(1 + \frac{2}{a^2} - \frac{1}{a^4}\right) m_t^2 C_0(M_H, m_t, m_t, m_t, M_H) \right. \\ & + \left(9 - \frac{6}{a^2}\right) m_t^2 C_0(M_H, m_t, M_H, M_H, m_t) - m_t^2 C_0(M_H, m_t, 0, 0, m_t) \\ & - \frac{1}{2a^2} m_t^2 C_0(M_H, m_t, 0, 0, 0) + \left(-6 + \frac{7}{a^2}\right) \ln 2a \\ & \left. - b \left(-6 + \frac{2}{a^2} + \frac{1}{a^4}\right) \left(\ln 2a + \ln \frac{1+b}{2}\right) + \frac{\pi\sqrt{3}}{2a^2} - \frac{i\pi}{2a^4} (a^2 - b) \right]. \quad (84) \end{aligned}$$

This result, based purely on the Lagrangian of the Higgs sector and the interaction of the Higgs and the Goldstone bosons with the fermion sector, is in complete agreement with the corresponding result¹ given by Kniehl [27], which was calculated using the *full* electroweak Lagrangian, assuming $M_H^2 > 4m_t^2 \gg M_W^2 \gg m_b^2$ and neglecting terms that are not enhanced by an inverse power of M_W^2 . The latter is equivalent to the limit $g_2 \rightarrow 0$, with $g_2/M_W = 2/v$. The agreement of the two results demonstrates the validity of the Goldstone boson equivalence theorem to one loop for $H \rightarrow t\bar{t}$.

To compare the decay matrix element of our equivalence-theorem calculation and the full electroweak calculation, we still need explicit expressions for the counterterms L_{bos} and L_{fer} for $M_H > 2m_t$. The expression for L_{bos} is given in Eq. (66). Evaluating the general expression for L_{fer} in Eq. (70) for $M_H > 2m_t$ using $m_b = 0$, we find that

$$\begin{aligned} L_{fer} = & \frac{1}{16\pi^2} \frac{m_t^2}{v^2} \left[(32a^4 - 36a^2 + 4) \ln 2a - 8a^2 + 7 \right. \\ & \left. - b (32a^4 - 20a^2) \left(\ln 2a + \ln \frac{1+b}{2}\right) \right]. \quad (85) \end{aligned}$$

¹The result given in Eq. (84) should be compared with the quantity $\frac{\alpha}{4\pi} \delta_{weak}$ as defined by Kniehl.

L_{fer} includes terms proportional to $a^4 \ln 2a$, a^4 , $a^2 \ln 2a$, and a^2 which involve powers of M_H^2/m_t^2 and are enhanced for large Higgs masses. However, these apparent enhancement terms cancel exactly when we expand the quantity b in terms of $1/a^2$, that is, L_{fer} is actually *not* enhanced in powers of M_H . This is a manifestation of the Veltman screening theorem [40]: at one loop, an internal Higgs-boson line with a large mass M_H leads only to a logarithmic enhancement. Since L_{fer} is derived from two-point functions with external fermion lines, no power-like enhancement is possible.

Assembling the partial results according to Eq. (80), we find the complete one-loop result for $\Delta\mathcal{T}$ in the limit $m_t \gg m_f$ for $f \neq t$,

$$\begin{aligned} \Delta\mathcal{T} = & -\frac{1}{16\pi^2} \left(\frac{m_t}{v}\right)^2 \left\{ \frac{a^2}{a^2-1} \left[\left(1 + \frac{2}{a^2} - \frac{1}{a^4}\right) M_H^2 C_0(M_H, m_t, m_t, m_t, M_H) \right. \right. \\ & + \left(9 - \frac{6}{a^2}\right) M_H^2 C_0(M_H, m_t, M_H, M_H, m_t) - M_H^2 C_0(M_H, m_t, 0, 0, m_t) \\ & \left. \left. - \frac{1}{2a^2} M_H^2 C_0(M_H, m_t, 0, 0, 0) + 2\pi\sqrt{3} - \frac{2i\pi}{a^2} (a^2 - b) \right] \right. \\ & + 8a^2 - \frac{11}{2} + \frac{3}{a^2} + \left(-32a^4 + 12a^2 + \frac{4}{a^2-1} \right) \ln(2a) \\ & + b \left(32a^4 + 4a^2 + 10 - \frac{3}{a^2} + \frac{12}{a^2-1} \right) \left(\ln(2a) + \ln \frac{1+b}{2} \right) \\ & \left. - 2a^2 \left(\frac{13}{2} - \pi\sqrt{3} \right) \right\}, \end{aligned} \quad (86)$$

where we have replaced an overall factor $4a^2 m_t^2$ by M_H^2 in the coefficients of the C_0 functions. In the context of the equivalence theorem this result is exact except for the approximation $m_f = 0$, $f \neq t$. The correction to the decay width is

$$\Gamma(H \rightarrow t\bar{t}) = \Gamma_B(H \rightarrow t\bar{t}) [1 + 2 \text{Re}\Delta\mathcal{T} + O(\lambda^2, \lambda g_t^2, g_t^4)]. \quad (87)$$

The functions $M_H^2 C_0$ in Eq. (86) are functions only of the ratio M_H^2/m_t^2 , i.e., of a^2 . Expanding these functions for $a^2 \gg 1$, we find that all of the contributions to $\Delta\mathcal{T}$ which are proportional to positive powers of a^2 cancel except for the last term in Eq. (86). That term arises from the boson-loop contribution to L_{bos} . The contributions from the fermion renormalization constants and the three-point functions are not power-enhanced, but grow only as $\ln(2a) = \ln(M_H/m_t)$. The purely bosonic corrections therefore give the dominant contribution to $\Delta\mathcal{T}$ for large Higgs-boson mass, a result that remains true to all orders in perturbation theory. In particular, at one loop,

$$\Delta\mathcal{T} \sim \frac{M_H^2}{16\pi^2 v^2} \left[\frac{1}{4} (13 - 2\pi\sqrt{3}) + O\left(\frac{m_t^2}{M_H^2} \ln \frac{M_H^2}{m_t^2}\right) + O\left(\frac{m_t^2}{M_H^2}\right) \right], \quad M_H^2 \gg m_t^2, \quad (88)$$

and the *fraction* of the total correction associated with the top-quark Yukawa coupling decreases rapidly for $M_H \gg m_t$. However, the actual correction associated with g_t may still be significant.

In Fig. 6 we show the equivalence theorem correction to the decay width, $\Gamma/\Gamma_B = 1 + 2 \text{Re}\Delta\mathcal{T}$, for $H \rightarrow t\bar{t}$ in the limit $m_b = 0$ (solid curve). This result is compared with the full

electroweak correction (short dashes) [41]. Away from the threshold, the equivalence theorem provides an excellent approximation. For $M_H = 500$ GeV (1 TeV), the EQT result is only 3.9% (1.8%) larger in magnitude than the full electroweak result. This is roughly the accuracy one would expect for a gauge contribution to $\Delta\mathcal{T}$ of order α/π .

We find that the $O(\lambda)$ correction (long dashes) represents the dominant contribution to the EQT correction for all values of M_H larger than $2m_t$, i.e., above the decay threshold. The one-loop Yukawa correction, $O(g_t^2)$, vanishes at about $M_H = 400$ GeV. For larger values of M_H , it is positive and adds to the $O(\lambda)$ contribution. While it initially grows more rapidly as a function of M_H than the $O(\lambda)$ correction, the M_H^2 behavior of the latter wins over the asymptotically logarithmic growth of the Yukawa correction for M_H greater than about 600 GeV. The Yukawa contribution represents a decreasing fraction of the total correction for higher masses, but is still significant numerically.

The very different behaviour of the EQT result and the full electroweak result at threshold is caused by a Coulomb singularity associated with the exchange of a virtual photon. Except for this QED effect, the EQT correction and the weak correction are in qualitative agreement at $M_H = 2m_t$: the EQT correction is about zero at threshold, and the weak correction is of the order of one percent. To have a better test of the validity of the equivalence theorem for values of $M_H \approx 2m_t$, we next consider the decay $H \rightarrow b\bar{b}$ which is free of the Coulomb singularity at $M_H = 2m_t$.

3.6 $H \rightarrow b\bar{b}$

In the mass region of a “light” Higgs boson, $M_H < 2m_t$, the dominant fermionic decay of the Higgs boson is the (much suppressed) decay $H \rightarrow b\bar{b}$. We can still treat this decay using the equivalence theorem provided that M_H is large compared to the masses of the W and Z bosons, and can use the comparison of the results with those of the full electroweak theory to test the limits of validity of the equivalence theorem for a “light” Higgs boson. For a “heavy” Higgs boson, $M_H > 2m_t$, the equivalence theorem result and the full electroweak result would be expected to agree to an accuracy similar to that in the previous section. However, the dependence of the one-loop correction to $H \rightarrow b\bar{b}$ on m_t is different from that encountered in top-quark production and therefore actually provides an independent check of the quality of the equivalence theorem for the case of a heavy Higgs boson.

We will again write the decay matrix element corrected to one loop in the form

$$-i\mathcal{M}_{H \rightarrow b\bar{b}} = -i\frac{m_b}{v}\bar{u}(p_1 - p_2, m_b)v(p_2, m_b)(1 + \Delta\mathcal{T}), \quad \Delta\mathcal{T} = L_{bos} + L_{fer} + L_{tri}. \quad (89)$$

The Born result is proportional to the ratio m_b/v . The one-loop equivalence-theorem corrections contributing to $\Delta\mathcal{T}$ are proportional to λ , m_t^2/v^2 , m_b^2/v^2 , and squares of lighter fermion masses. Evaluating these contributions to $\Delta\mathcal{T}$, we set $m_b = 0$ and also neglect other light fermions. In this limit, L_{fer} only receives contributions from the bottom-quark self-energy diagram which contains a (w, t) loop, Fig. 5, and L_{tri} reduces to a sum over only two triangle graphs, L_3 and L_6 in Fig. 2, with the tree-level factor $-im_b/v$ extracted.

The bosonic counterterm L_{bos} is given by Eq. (66) or (67), depending on the value of M_H . The fermionic counterterm, L_{fer} , and the triangle graphs, L_{tri} , depend on the flavor of the final fermion pair and need to be re-evaluated for the present decay into bottom-quarks, i.e., their results differ from the above results for the decay of the Higgs boson into top quarks.

The fermionic counterterm is evaluated using the general result in Eq. (70) with $m_b \rightarrow 0$,

$$L_{fer} = \frac{1}{16\pi^2} \frac{m_t^2}{v^2} [-2B_0(0, 0, m_t^2)] . \quad (90)$$

The triangle contribution is calculated from the expressions for L_3 and L_6 using the Dirac equation and setting $m_b = 0$:

$$\begin{aligned} L_{tri} = & \frac{1}{16\pi^2} \frac{m_t^2}{v^2} \left[2B_0(0, 0, m_t^2) + \left(2 - \frac{1}{a^2} \right) [B_0(M_H^2, m_t^2, m_t^2) - B_0(0, 0, m_t^2)] + 2 - 4 \ln 2a \right. \\ & \left. + \frac{1}{4a^4} M_H^2 C_0(M_H^2, 0, m_t^2, m_t^2, 0) + 2 \left(1 + \frac{1}{4a^2} \right) M_H^2 C_0(M_H^2, 0, 0, 0, m_t^2) + 2i\pi \right] . \quad (91) \end{aligned}$$

The divergencies related to the integrals B_0 cancel in the sum $\Delta\mathcal{T} = L_{bos} + L_{fer} + L_{tri}$. In case of a “light” Higgs, the complete one-loop electroweak radiative correction to the amplitude for the decay $H \rightarrow b\bar{b}$ is now found to be

$$\begin{aligned} \Delta\mathcal{T} = & \frac{1}{16\pi^2} \frac{m_t^2}{v^2} \left[\frac{5}{2} - \frac{4}{a^2} - 4 \ln 2a - \left(\frac{1}{a^2} - \frac{2}{a^4} \right) \phi \sin(\phi) - 3 \left(2a^2 - 3 + \frac{1}{a^2} \right) \frac{\phi}{\sin(\phi)} \right. \\ & \left. + \frac{1}{4a^4} M_H^2 C_0(M_H^2, 0, m_t, m_t, 0) + 2 \left(1 + \frac{1}{4a^2} \right) M_H^2 C_0(M_H^2, 0, 0, 0, m_t^2) \right. \\ & \left. + 2a^2 \left(\frac{13}{2} - \pi\sqrt{3} \right) + 2i\pi \right] , \quad M_H < 2m_t, \quad (92) \end{aligned}$$

and the heavy-Higgs result is

$$\begin{aligned} \Delta\mathcal{T} = & \frac{1}{16\pi^2} \frac{m_t^2}{v^2} \left[\frac{5}{2} - \frac{4}{a^2} - 4 \ln 2a + b \left(2 + \frac{5}{a^2} \right) \left[\ln(2a) + \ln \left(\frac{1+b}{2} \right) \right] \right. \\ & \left. + \frac{1}{4a^4} M_H^2 C_0(M_H^2, 0, m_t, m_t, 0) + 2 \left(1 + \frac{1}{4a^2} \right) M_H^2 C_0(M_H^2, 0, 0, 0, m_t^2) \right. \\ & \left. + 2a^2 \left(\frac{13}{2} - \pi\sqrt{3} \right) + i\pi \left(2 - b \left(1 + \frac{1}{a^2} \right) \right) \right] , \quad M_H > 2m_t. \quad (93) \end{aligned}$$

In Fig. 7 we show the equivalence theorem correction to the decay width, $\Gamma/\Gamma_B = 1 + 2 \text{Re}\Delta\mathcal{T}$, for $H \rightarrow b\bar{b}$ in the limit $m_b = 0$ (solid curve). This result is compared with the full electroweak correction (short dashes) [41]. For large values of M_H , the EQT result approximates the full correction very well, in agreement with our findings in the case of $H \rightarrow t\bar{t}$. The difference between the two corrections is again of the magnitude of the expected gauge corrections. The dominant contribution overall is the $O(\lambda)$ correction (long dashes), which grows more rapidly than the Yukawa correction for M_H larger than about 700 GeV.

For a Higgs mass of about 400 GeV, the top-quark Yukawa correction cancels the $O(\lambda)$ contribution, and the one-loop radiative corrections are actually determined mainly by the gauge corrections. However, the magnitude of the gauge correction is still small compared to the magnitude of the Yukawa correction at this point, with an absolute value less than 1%. The equivalence theorem breaks down for M_H less than about 200 GeV, where the condition $M_H \gg M_W, M_Z$ is of questionable validity. The gauge corrections are the dominant

corrections in this region, and are larger in magnitude than both the Yukawa and the $O(\lambda)$ corrections. In addition, the gauge correction displays Coulomb singularities at $M_H = 2M_Z$ and $2M_W$, a feature which is unique to gauge interactions and cannot be reproduced by the equivalence theorem. This threshold region requires special treatment.

Qualitatively, the EQT result approximates the full electroweak corrections rather well for Higgs masses larger than 200 GeV. The full electroweak correction remains finite at the threshold for top-quark production at $M_H = 2m_t$, and the expected threshold kink is reproduced by the equivalence-theorem calculation. We conclude that the equivalence theorem is a useful tool even in the case of a “light” Higgs boson with a mass in the range $2M_Z < M_H < 2m_t$, with the EQT correction to $\Gamma(H \rightarrow b\bar{b})$ giving a good estimate of the full electroweak correction.

4 Summary

The equivalence theorem is known to be an excellent tool in describing heavy Higgs physics. Calculations based on the equivalence theorem are usually carried out by using massless Goldstone bosons and a massive Higgs, neglecting all gauge and Yukawa couplings. While the original applications of the equivalence theorem were to tree-level processes [1, 2], it has since been shown that the equivalence theorem also provides a simple way of calculating the dominant contributions of internal W^\pm , Z , and H bosons [13, 14, 15].

In this paper, we have extended the equivalence theorem by systematically including the Yukawa interactions. This is possible because the approximations underlying the equivalence theorem are independent of the values of the Yukawa couplings. In particular, we have formulated a renormalization procedure which is simultaneously consistent with the requirements of the equivalence theorem, and has the correct relations to physical observables in the limit of vanishing gauge couplings. Because the top quark is quite massive, g_t is large, and it is generally necessary to include the top-quark Yukawa coupling in calculations of electroweak radiative corrections. The framework presented here allows these calculations to be done rather simply using the EQT, with the gauge couplings set to zero. One can hope to obtain good approximate results for radiative corrections to Higgs-sector processes in the case of a “light” Higgs, $2M_Z < M_H < 2m_t$, and excellent approximations for larger Higgs masses.

As an example, we calculated the one-loop corrections to fermionic Higgs decays using the equivalence theorem with fermions, and compared our results with the results obtained from a full electroweak calculation. Since the Yukawa interactions are negligible except for the top-quark coupling, we only included contributions from the latter. We find that the Higgs coupling λ and the Yukawa coupling g_t give the dominant corrections to the decay rates for $M_H > 2m_t$. The much smaller contributions of the transverse gauge couplings are only significant very close to 400 GeV in the case $H \rightarrow b\bar{b}$ where the dominant contributions cancel, and near the decay threshold for $H \rightarrow t\bar{t}$ where virtual photon exchange produces a Coulomb singularity.

In the mass range $2M_Z < M_H < 2m_t$, the process $H \rightarrow b\bar{b}$ is the only significant fermionic decay. The one-loop radiative corrections to this decay associated with the quartic Higgs-boson coupling, the top-quark Yukawa coupling, and the transverse electroweak gauge couplings are all similar in magnitude, but with differing signs. In particular, a partial cancellation of the Higgs and Yukawa contributions makes the gauge correction equally important. The total correction is very small, less than 2%. It seems plausible that the sum of the *magnitudes*

of Higgs and Yukawa contribution would give a good estimate for an upper bound on the magnitude of the complete electroweak radiative correction for a variety of electroweak processes. In absence of cancellations between the $O(g_t^2)$ and $O(\lambda)$ corrections, the equivalence theorem result is expected to be the dominant correction for Higgs masses larger than $2M_Z$. Threshold singularities arising from the gauge sector cannot, of course, be reproduced using the equivalence theorem, and require special treatment in any case.

According to Veltman's screening theorem [40], the $O(\lambda)$ corrections are the only one-loop corrections that grow proportional to M_H^2 . We find that this asymptotic growth of the correction is dominant only for Higgs-boson masses larger than 600 to 700 GeV, assuming a top-quark mass of 175 GeV. For smaller values of M_H , we find the $O(g_t^2)$ corrections to have the stronger dependence on M_H .

In conclusion we find that the calculation of radiative corrections using the equivalence theorem is greatly improved if the Yukawa interactions are included. The limit of $m_f = 0$, $f \neq t$ allows for a relatively simple calculation of the dominant radiative corrections, yielding an excellent approximation of the full electroweak corrections for the heavy Higgs-case, and order-of-magnitude estimates for $2M_Z < M_H < 2m_t$.

Acknowledgments

The authors would like to thank B. Kniehl for providing the data for the full electroweak corrections shown in Figs. 6 and 7. One of the authors (K.R.) would like to thank A. Dabelstein for useful conversations, and the Deutsche Forschungsgemeinschaft (DFG) for financial support under contract no. Li519/2-1. The work of the other author (L.D.) was supported in part by the U.S. Department of Energy under Contract No. DE-FG02-95ER40896.

A Tadpoles and self-energies of H , z , w and f .

The only neutral field that receives a shift in its vacuum expectation value because of tadpole contributions is the Higgs field h ; the z field does not. The z and w fields have couplings through \mathcal{L}_H , Eq. (15), that require an even number of fields to participate, and no purely bosonic tadpole graphs can be formed for the z . With the addition of the fermionic Lagrangian \mathcal{L}_F , Eq. (16), the z can form a tadpole with a fermion loop, but the presence of a factor γ^5 in the z -fermion coupling and a trace over the γ -matrices involved yields a vanishing result. However, fermion loops contribute to the Higgs tadpoles as shown in Fig. 1 since no γ^5 is involved in the H -fermion coupling. Taking into account both the bosonic and fermionic contributions, the Higgs one-point function (tadpole function) is

$$T = \frac{1}{16\pi^2} \left(-\frac{3M_H^2}{2v} A_0(M_H^2) + \sum_f N_C^f \frac{4m_f^2}{v} A_0(m_f^2) \right). \quad (94)$$

The graphs of the one-loop one-particle irreducible self-energy contributions to the bosonic and fermionic fields are shown in Figs. 3, 4, and 5. They can be evaluated as [32, 35]

$$\Pi_H^0(p^2) = -\frac{1}{16\pi^2} (3\lambda A_0(M_H^2) + 18\lambda^2 v^2 B_0(p^2, M_H^2, M_H^2) - 6\lambda^2 v^2 B_0(p^2, 0, 0))$$

$$+ \frac{1}{16\pi^2} \sum_f N_C^f \frac{m_f^2}{v^2} (4A_0(m_f^2) - (2p^2 - 8m_f^2)B_0(p^2, m_f^2, m_f^2)) , \quad (95)$$

$$\begin{aligned} \Pi_z^0(p^2) &= -\frac{1}{16\pi^2} (\lambda A_0(M_H^2) + 4\lambda^2 v^2 B_0(p^2, M_H^2, 0)) \\ &+ \frac{1}{16\pi^2} \sum_f N_C^f \frac{m_f^2}{v^2} (4A_0(m_f^2) - 2p^2 B_0(p^2, m_f^2, m_f^2)) , \end{aligned} \quad (96)$$

$$\begin{aligned} \Pi_w^0(p^2) &= -\frac{1}{16\pi^2} (\lambda A_0(M_H^2) + 4\lambda^2 v^2 B_0(p^2, M_H^2, 0)) \\ &- \frac{1}{16\pi^2} \sum_{(f,f')} N_C^f \frac{1}{v^2} [8m_f^2 m_{f'}^2 B_0(p^2, m_f^2, m_{f'}^2) \\ &\quad + 2(m_f^2 + m_{f'}^2) (-A_0(m_f^2) - A_0(m_{f'}^2) \\ &\quad + (p^2 - m_f^2 - m_{f'}^2) B_0(p^2, m_f^2, m_{f'}^2))] , \end{aligned} \quad (97)$$

$$\Pi_{V,f}^0(p^2) = \frac{1}{16\pi^2} \frac{m_f^2}{v^2} \left[-B_1(p^2, m_f^2, M_H^2) - B_1(p^2, m_f^2, 0) - \left(1 + \frac{m_{f'}^2}{m_f^2}\right) B_1(p^2, m_{f'}^2, 0) \right] \quad (98)$$

$$\Pi_{A,f}^0(p^2) = \frac{1}{16\pi^2} \frac{m_f^2}{v^2} \left(-1 + \frac{m_{f'}^2}{m_f^2} \right) B_1(p^2, m_{f'}^2, 0), \quad (99)$$

$$\Pi_{S,f}^0(p^2) = \frac{1}{16\pi^2} \frac{m_f^2}{v^2} \left(B_0(p^2, m_f^2, M_H^2) - B_0(p^2, m_f^2, 0) - 2 \frac{m_{f'}^2}{m_f^2} B_0(p^2, m_{f'}^2, 0) \right). \quad (100)$$

The z and w self-energies only differ in the fermionic part, and are equal for $m_f = m_{f'}$. It is easily shown using the explicit results given for A_0 and B_0 in Appendix B that the self energies Π_z^0 and Π_w^0 of the z and w bosons are equal for $p^2 = 0$, independently of the values of the masses m_f and $m_{f'}$. This equality does not extend to the derivatives of the self-energy functions. Finally, comparing the expressions in Eqs. (94) and (96), we find that

$$T_0/v_0 = \Pi_z^0(0) = \Pi_w^0(0) \quad (101)$$

as stated in Eq. (28).

B Loop integral expressions

B.1 Scalar integral expressions

Here we define the scalar integral expressions needed to calculate the two- and three-point functions [32, 36]. The more complicated vector and tensor loop integrals which arise from diagrams containing fermions can be reduced to sums of scalar integrals by standard techniques [32, 37]. We will follow the definitions in [32].

For the calculation of the one-loop self-energies in D dimensions we use

$$A_0(m_0^2) = \frac{(2\pi\mu)^{(4-D)}}{i\pi^2} \int d^D q \frac{1}{q^2 - m_0^2 + i\epsilon'}, \quad (102)$$

$$B_0(p^2, m_0^2, m_1^2) = \frac{(2\pi\mu)^{(4-D)}}{i\pi^2} \int d^D q \frac{1}{[q^2 - m_0^2 + i\epsilon'][(q+p)^2 - m_1^2 + i\epsilon']} . \quad (103)$$

The vertex corrections involve the integrals

$$C_0(p_1^2, p_2^2, m_0^2, m_1^2, m_2^2) = \frac{(2\pi\mu)^{(4-D)}}{i\pi^2} \int d^D q \frac{1}{[q^2 - m_0^2 + i\epsilon'][(q+p_1)^2 - m_1^2 + i\epsilon'][(q+p_2)^2 - m_2^2 + i\epsilon']} . \quad (104)$$

The arbitrary energy scale μ is introduced to fix the energy dimensions of the functions A_0 , B_0 , and C_0 independent of the value of D , and the infinitesimal quantity $i\epsilon'$ defines the integration path in the complex plane. Note that Böhm et al. [16] define A_0 with an overall minus sign, while Kniehl [27, 42] introduces a minus sign in the definition of C_0 .

These integrals can be evaluated in a straightforward manner using Feynman parameters. Complete analytic results are given by Denner [32], though it is useful to go back to the Feynman representation of the integrals for certain values of p_i^2 and m_i^2 to avoid problems with infrared singularities.

The integral B_1 that appears in Eq. (70) is given by

$$B_1(p^2, m_0^2, m_1^2) = \frac{1}{2p^2} [A_0(m_0^2) - A_0(m_1^2) + (m_1^2 - m_0^2 - p^2)B_0(p^2, m_0^2, m_1^2)] . \quad (105)$$

The derivatives of A_0 , B_0 , and B_1 with respect to p^2 , the square of the external momentum, are also needed in the calculation of the multiplicative wavefunction renormalization constants Z_i . The function A_0 actually does not depend on the external momentum, and its derivative with respect to p^2 therefore vanishes. For the remaining derivatives we introduce the usual notation

$$\partial B_i(M^2, m_0^2, m_1^2) \equiv \left. \frac{\partial}{\partial p^2} B_i(p^2, m_0^2, m_1^2) \right|_{p^2=M^2} , \quad i = 0, 1. \quad (106)$$

These derivatives are given, for example, by Denner [32].

The reduction of tensor integrals such as those which appear in Eqs. (77) and (78) to sums of scalar integrals is discussed in detail in [32]. The results given there agree with our calculations [35] except for a typographical error in Denner's expression for the function C_{00} in his Eq. (C.37). In our notation, the correct result for C_{00} is

$$C_{00} = \frac{1}{4} [B_0((p_1 - p_2)^2, m_2^2, m_1^2) + (m_0^2 - m_1^2 + p_1^2)C_1 + (m_0^2 - m_2^2 + p_2^2)C_2 + 2m_0^2 C_0] . \quad (107)$$

References

- [1] B.W. Lee, C. Quigg, and H.B. Thacker, Phys. Rev. Lett. **38**, 883 (1977); Phys. Rev. D **16**, 1519 (1977).
- [2] M.S. Chanowitz and M.K. Gaillard, Nucl. Phys. **B261**, 379 (1985).
- [3] G. Passarino, Phys. Lett. B **156**, 231 (1985).

- [4] W.J. Marciano and S.S.D. Willenbrock, Phys. Rev. D **37**, 2509 (1988).
- [5] L. Durand, J.M. Johnson, and J.L. Lopez, Phys. Rev. Lett. **64**, 1215 (1990).
- [6] G. Passarino, Nucl. Phys. **B343**, 31 (1990).
- [7] L. Durand, J.M. Johnson, and J.L. Lopez, Phys. Rev. D **45**, 3112 (1992).
- [8] L. Durand, J.M. Johnson, and P.N. Maher, Phys. Rev. D **44**, 127 (1991).
- [9] L. Durand, P.N. Maher, and K.R. Riesselmann, Phys. Rev. D **48**, 1084 (1993).
- [10] A. Ghinculov, Univ. Freiburg preprint THEP 95/11, hep-ph/9507240.
- [11] R. Barbieri, M. Beccaria, P. Ciafaloni, G. Curci, and A. Viceré, Phys. Lett. B **288**, 95 (1992); **312**, 511(E) (1993); Nucl. Phys. **B409**, 105 (1993).
- [12] J. Cornwall, D. Levin, and G. Tiktopoulos, Phys. Rev. D **10**, 1145 (1974), Appendix A; C.E. Vayonakis, Lett. Nuovo Cimento **17**, 383 (1976); G. Gounaris, R. Kögeler, and H. Neufeld, Phys. Rev. D **34**, 3257 (1986); Y. Yao and C. Yuan, *ibid.* **38**, 2237 (1988).
- [13] J. Bagger and C. Schmidt, Phys. Rev. D **41**, 264 (1990); A. Aoki, Kyoto University Report No. RIFP-705, 1987 (unpublished); in *Physics at TeV Scale*, Proceedings of the Meeting, Tsukuba, Japan, 1988, edited by K. Hidaka and K. Hikasa (KEK, Tsukuba, 1988).
- [14] H. Veltman, Phys. Rev. D **41**, 2294 (1990).
- [15] H.-J. He, Y.-P. Kuang, and X. Li, Phys. Rev. Lett. **69**, 2619 (1992); Phys. Rev. D **49**, 4842 (1994); H.-J. He, Y.-P. Kuang, and C.-P. Yuan, Phys. Rev. D **51**, 6463 (1995).
- [16] M. Böhm, H. Spiesberger, and W. Hollik, Fortschr. Phys. **34**, 687 (1986).
- [17] J. Goldstone, Nuovo Cimento **54**, 154 (1961); T.-P. Cheng and L.-F. Li, *Gauge Theory of Elementary Particle Physics* (Oxford University Press, Oxford, England, 1984), Chap. 5.
- [18] A. Sirlin and R. Zucchini, Nucl. Phys. **B266**, 389 (1986).
- [19] P.N. Maher, L. Durand, and K. Riesselmann, Phys. Rev. D **48**, 1061 (1993); **52**, 553(E) (1995).
- [20] R.S. Lytel, Phys. Rev. D **22**, 505 (1980).
- [21] We follow the convention for an unstable Higgs boson of equating M_H^2 with the value of p^2 at which the real part of the expression on the righthand side of Eq. (25) vanishes for $p^2 = p^2 + i\epsilon$, $\epsilon \rightarrow 0+$. δm_0^2 , v_0 , and λ_0 are taken as real. With this definition, $\Gamma_{H_0}^{(2)} \approx Z_H^{-1}(p^2 - M_H^2 + iM_H\Gamma_H)$ for $p^2 \approx M_H^2$, with $\Gamma_H = -Z_H \text{Im}\Pi_H^0(M_H^2)$. A definition of M_H^2 as the complex value of p^2 at which the expression in Eq. (25) vanishes exactly (i.e., a pole mass in terms of the propagator), requires some changes which we will not discuss here.

- [22] The two different approaches were used, for example, by Maher *et al.* [19] and by Ghinculov and van der Bij [34, 33], respectively, in their calculations of the two-loop self energies.
- [23] J.C. Taylor, *Gauge Theories of Weak Interactions* (Cambridge University Press, Cambridge, England, 1976), Chap. 14.
- [24] It is easy to see heuristically why the constant C has the form given. Most derivations of the equivalence theorem are based on variations of the Feynman-'t Hooft gauge-fixing condition, that $\langle ik_\mu \mathbf{W}_a^{0,\mu} + M_0 \mathbf{w}^0 \rangle = 0$ for any physical matrix element. With the observation that the ratio k^μ/M differs from the longitudinal polarization vector ξ for a massive particle only by terms of order M/k^0 , this becomes a relation between amplitudes which involve longitudinally polarized vector bosons, and amplitudes which involve the corresponding Goldstone scalars, provided $k^0 \gg M$. In particular, in the case of scattering processes, the amplitudes for the scattering of n longitudinally polarized vector bosons W_L^\pm, Z_L and any number of other external particles are related to the corresponding scattering amplitudes for the scalar bosons w^\pm, z to which W_L^\pm, Z_L reduce in the limit of vanishing gauge couplings, $g, g' \rightarrow 0$, by
- $$T(W_L^\pm, Z_L, H, \dots) = (iC)^n T(w^\pm, z, H, \dots) + \dots, \quad (108)$$
- where the corrections are of orders $M_W/\sqrt{s}, g^2, g'^2$, and M_W^2/M_H^2 [13, 14]. The constant of proportionality simply corrects for the ratios of the wave function renormalization constants $Z_W^{1/2}/Z_\Phi^{1/2} = Z_W^{1/2}/Z_w^{1/2}$ which appear in the scattering amplitudes, and for the ratio of the bare and physical masses. C reduces to unity for $g, g' \rightarrow 0$, the limit in which the two theories merge, provided that the renormalizations are defined at a mass scale $m \ll M_H, \sqrt{s}$ [13].
- [25] For an unstable fermion, we will use a definition analogous to that used for the Higgs boson [21], and define the square of the fermion mass as the value of p^2 at which the real part of the denominator in Eq. (42) vanishes. Other definitions are also possible.
- [26] See, for example, J.D. Bjorken and S.D. Drell, *Relativistic Quantum Fields* (McGraw-Hill Book Company, New York, 1965), Chap. 16.
- [27] B.A. Kniehl, Nucl. Phys. **B376**, 3 (1992).
- [28] A. Dabelstein and W. Hollik, Z. Phys. C **53**, 507 (1992).
- [29] D.Yu. Bardin, B.M. Vilenskii, and P.Kh. Khristov, Yad. Fiz. **53**, 240, (1991) [Sov. J. Nucl. Phys. **53**, 152 (1991)].
- [30] L. Durand, B.A. Kniehl, and K. Riesselmann, Phys. Rev. Lett. **72**, 2534 (1994); **75**, 1699(E) (1995).
- [31] S. Dawson and S. Willenbrock, Phys. Rev. D **62**, 1232 (1989).
- [32] A. Denner, Fortschr. Phys. **41**, 307 (1993).
- [33] A. Ghinculov and J.J. van der Bij, Nucl. Phys. **B436**, 30 (1995).

- [34] A. Ghinculov, Phys. Lett. B **337**, 137 (1994); **346**, 426(E) (1995).
- [35] K. Riesselmann, University of Wisconsin thesis, 1994 (unpublished).
- [36] G. 't Hooft and M. Veltman, Nucl. Phys. **B153**, 365 (1979).
- [37] G. Passarino and M. Veltman, Nucl. Phys. **B160**, 151 (1979), Appendices D, E.
- [38] L. Resnick, M.K. Sundareshan, and P.J.S. Watson, Phys. Rev. D **8**, 172 (1973).
- [39] A. Mendez, Mod. Phys. Lett. **A5** 2087 (1990); (E) **A6** (1991) 1533.
- [40] M. Veltman, Acta Phys. Pol. B **8**,475 (1977).
- [41] For a comparison of the electroweak corrections with the QCD corrections see: L. Durand, B.A. Kniehl, and K. Riesselmann, Phys. Rev. D **51**, 5007 (1995).
- [42] B.A. Kniehl, Nucl. Phys. **B352**, 1 (1991), Appendix A.




Figure 1: The one-loop tadpole diagrams which are cancelled by a counterterm to avoid a shift in the vacuum expectation value of the Higgs field. Thick lines correspond to the Higgs particle, dotted lines represent the massless Goldstone bosons, and solid lines with arrows refer to fermions. A summation over all Goldstone boson and fermion loops is implied.




Figure 2: The six triangle diagrams contributing to $H \rightarrow f\bar{f}$ at one loop within the framework of the equivalence theorem. The different lines have the same meaning as in Fig. 1. The fermion f' is the $SU(2)_L$ partner of f .




Figure 3: Higgs self-energy contributions at one loop. The different lines have the same meaning as in Fig. 1. A summation over all Goldstone boson and fermion loops is implied.




Figure 4: The Goldstone boson self energies at one loop. The different lines have the same meaning as in Fig. 1. The diagrams on the right need to be summed over the different Goldstone boson and fermion loop contributions, respectively. In case of an external w or z , the fermion loop consists of an $f f'$ or $f f$ pair, respectively, where f' is the $SU(2)_L$ partner of f .




Figure 5: The three diagrams contributing to the fermion self-energies at one loop. The different lines have the same meaning as in Fig. 1. The fermion f' is the $SU(2)_L$ partner of the external fermion f .




Figure 6: The one-loop correction factor to the decay width $H \rightarrow t\bar{t}$. The solid curve gives the equivalence theorem (EQT) result consisting of the sum of the $O(\lambda)$ and $O(g_t^2)$ corrections. Light fermion couplings are neglected. The result is compared with the full electroweak correction obtained in [27, 28, 29] and the EQT result without fermion corrections.




Figure 7: The one-loop correction factor to the decay width $H \rightarrow b\bar{b}$. The solid curve gives to the equivalence theorem result consisting of the sum of the $O(\lambda)$ and the $O(g_t^2)$ corrections. Light fermion couplings are neglected. The result is compared with the full electroweak correction obtained in [27, 28, 29] and the EQT result without fermion corrections.