

TESTS OF THE STANDARD MODEL

WOLFGANG HOLLIK

*Institut für Theoretische Physik, Universität Karlsruhe
D-76128 Karlsruhe, Germany*

In this conference report a summary is given on the recent theoretical work that has contributed to improve the theoretical predictions for testing the standard model in present and future experiments. Precision calculations for the Z resonance are reviewed and the status of the standard model is discussed in the light of the recent top discovery and of the results from precision experiments. Furthermore, theoretical progress for the Higgs search, for W physics at LEP II, and for the anomalous magnetic moment of the muon is summarized. New Physics beyond the minimal model is briefly discussed, in particular the minimal supersymmetric standard model in view of the recent electroweak precision data.

1 Introduction

The present generation of high precision experiments imposes stringent tests on the standard model of electroweak and strong interactions¹. Besides the impressive achievements in the determination of the Z boson parameters^{1,2} and the W mass³, the most important step has been the confirmation of the top quark at the Tevatron^{4,5,6}. Its mass determination by the CDF collaboration⁴ yields $m_t = 176 \pm 8 \pm 10$ GeV and by the D0 collaboration⁵: $m_t = 199_{-21}^{+19} \pm 22$ GeV, resulting in an weighted average of

$$m_t = 180 \pm 12 \text{ GeV}. \quad (1)$$

The high experimental sensitivity in the electroweak observables, at the level of quantum effects, requires the highest standards on the theoretical side as well. A sizeable amount of work has contributed over the last few years to a steadily rising improvement of the standard model predictions pinning down the theoretical uncertainties to a level sufficiently small for the current interpretation of the precision data, but still sizeable enough to provoke conflict with a further increase in the experimental accuracy.

In this report we review the recent theoretical work which has provided the current theoretical basis for tests of the standard model at present and future colliders. A detailed description of the standard model predictions for observables around the Z peak can be found in the Working Group report on 'Precision Calculations for the Z Resonance'⁷ (see also the talk by Bardin at this conference⁸). This is followed by a discussion of the present status of the standard model in the light of the recent experimental results. Afterwards we report on theoretical progress in the study of Higgs and WW physics, and for $g-2$ for muons. A short discussion of the status of the MSSM concludes this presentation.

2 Theory for precision tests

2.1 Radiative corrections

The possibility of performing precision tests is based on the formulation of the standard model as a renormalizable quantum field theory preserving its predictive power beyond tree level calculations. With the experimental accuracy in the investigation of the fermion-gauge boson interactions being sensitive to the loop induced quantum effects, also the more subtle parts of the standard model Lagrangian are probed. The higher order terms induce the sensitivity of electroweak observables to the top and Higgs mass m_t, M_H and to the strong coupling constant α_s .

Before one can make predictions from the theory, a set of independent parameters has to be determined from experiment. All the practical schemes make use of the same physical input quantities

$$\alpha, G_\mu, M_Z, m_f, M_H; \alpha_s \quad (2)$$

for fixing the free parameters of the standard model. Differences between various schemes are formally of higher order than the one under consideration. The study of the scheme dependence of the perturbative results, after improvement by resumming the leading terms, allows us to estimate the missing higher order contributions.

Two fermion induced large loop effects in electroweak observables deserve a special discussion:

- The light fermionic content of the subtracted photon vacuum polarization

$$\Delta\alpha = \Pi_{ferm}^\gamma(0) - \text{Re} \Pi_{ferm}^\gamma(M_Z^2)$$

corresponds to a QED induced shift in the electromagnetic fine structure constant. The recent update of the evaluation of the light quark content by Eidelman, Jegerlehner⁹ and Burkhardt, Pietrzyk¹⁰ both

yield the result

$$(\Delta\alpha)_{had} = 0.0280 \pm 0.0007$$

and thus confirm the previous value of ¹¹ with an improved accuracy. Other determinations ^{12,13} agree within one standard deviation. Together with the leptonic content, $\Delta\alpha$ can be resummed resulting in an effective fine structure constant at the Z mass scale:

$$\alpha(M_Z^2) = \frac{\alpha}{1 - \Delta\alpha} = \frac{1}{128.89 \pm 0.09}. \quad (3)$$

- For a general structure of the scalar sector, the electroweak mixing angle is related to the vector boson masses by

$$\begin{aligned} \sin^2 \theta &= 1 - \frac{M_W^2}{\rho M_Z^2} = 1 - \frac{M_W^2}{M_Z^2} + \frac{M_W^2}{M_Z^2} \Delta\rho \\ &\equiv s_W^2 + c_W^2 \Delta\rho \end{aligned} \quad (4)$$

where the ρ -parameter $\rho = (1 - \Delta\rho)^{-1}$ is an additional free parameter. In the standard model, one has the tree level relation $\rho = 1$. Loop effects, however, induce a deviation $\Delta\rho \neq 0$. The main contribution is from the (t, b) doublet ¹⁴, in 1-loop and neglecting m_b given by:

$$\Delta\rho^{(1)} = 3x_t, \quad x_t = \frac{G_\mu m_t^2}{8\pi^2 \sqrt{2}} \quad (5)$$

Higher order irreducible contributions have become available, modifying $\Delta\rho$ according to

$$\Delta\rho = 3x_t \cdot [1 + x_t \rho^{(2)} + \delta\rho_{QCD}] \quad (6)$$

The electroweak 2-loop part ^{15,16} is described by the function $\rho^{(2)}(M_H/m_t)$ derived in ¹⁶ for general Higgs masses. $\delta\rho_{QCD}$ is the QCD correction to the leading $G_\mu m_t^2$ term ^{17,18}

$$\delta\rho_{QCD} = -\frac{\alpha_s(\mu)}{\pi} c_1 + \left(\frac{\alpha_s(\mu)}{\pi}\right)^2 c_2(\mu) \quad (7)$$

with

$$c_1 = \frac{2}{3} \left(\frac{\pi^2}{3} + 1\right) \quad (8)$$

and the recently calculated 3-loop coefficient ¹⁸

$$c_2 = -14.59 \text{ for } \mu = m_t \text{ and 6 flavors} \quad (9)$$

with the on-shell top mass m_t . It reduces the scale dependence of $\delta\rho_{QCD}$ significantly and hence is an important entry to decrease the theoretical uncertainty of the standard model predictions for precision observables (see section 2.4).

2.2 The vector boson masses

The correlation between the masses M_W, M_Z of the vector bosons in terms of the Fermi constant G_μ , in 1-loop order given by ¹⁹:

$$\frac{G_\mu}{\sqrt{2}} = \frac{\pi\alpha}{2s_W^2 M_W^2} [1 + \Delta r(\alpha, M_W, M_Z, M_H, m_t)]. \quad (10)$$

The decomposition

$$\Delta r = \Delta\alpha - \frac{c_W^2}{s_W^2} \Delta\rho^{(1)} + (\Delta r)_{remainder}. \quad (11)$$

separates the leading fermionic contributions $\Delta\alpha$ and $\Delta\rho$. All other terms are collected in the $(\Delta r)_{remainder}$, the typical size of which is of the order ~ 0.01 .

The presence of large terms in Δr requires the consideration of higher than 1-loop effects. The modification of Eq. (10) according to

$$\begin{aligned} 1 + \Delta r &\rightarrow \frac{1}{(1 - \Delta\alpha) \cdot (1 + \frac{c_W^2}{s_W^2} \Delta\rho) - (\Delta r)_{remainder}} \\ &\equiv \frac{1}{1 - \Delta r} \end{aligned} \quad (12)$$

accommodates the following higher order terms (Δr in the denominator is an effective correction including higher orders):

- The leading log resummation ²⁰ of $\Delta\alpha$: $1 + \Delta\alpha \rightarrow (1 - \Delta\alpha)^{-1}$
- The resummation of the leading m_t^2 contribution ²¹ in terms of $\Delta\rho$ in Eq. (6). Beyond the $G_\mu m_t^2 \alpha_s$ approximation through the ρ -parameter, the complete $O(\alpha_s)$ corrections to the self energies are available from perturbative calculations ²² and by means of dispersion relations ²³. All the higher order terms contribute with the same positive sign to Δr , thus making the top mass dependence of Δr significantly flatter. This is of high importance for the determination of an upper bound on m_t from M_W measurements, which is affected by the order of 10 GeV. Quite recently, also non-leading terms to Δr of the type

$$\Delta r_{(bt)} = 3x_t \left(\frac{\alpha_s}{\pi}\right)^2 \left(a_1 \frac{M_Z^2}{m_t^2} + a_2 \frac{M_Z^4}{m_t^4}\right)$$

have been computed ²⁴. For $m_t = 180$ GeV they contribute an extra term of $+0.0001$ to Δr and thus are within the uncertainty from $\Delta\alpha$.

- With the quantity $(\Delta r)_{remainder}$ in the denominator non-leading higher order terms containing mass singularities of the type $\alpha^2 \log(M_Z/m_f)$ from light fermions are also incorporated ²⁵.

2.3 Z boson observables

Measurements of the Z line shape in $e^+e^- \rightarrow f\bar{f}$ yield the parameters M_Z , Γ_Z , and the partial widths Γ_f or the peak cross section

$$\sigma_0^f = \frac{12\pi}{M_Z^2} \cdot \frac{\Gamma_e \Gamma_f}{\Gamma_Z^2}. \quad (13)$$

Whereas M_Z is used as a precise input parameter, together with α and G_μ , the width, partial widths and asymmetries allow comparisons with the predictions of the standard model. The predictions for the partial widths as well as for the asymmetries can conveniently be calculated in terms of effective neutral current coupling constants for the various fermions.

Effective Z boson couplings: The effective couplings follow from the set of 1-loop diagrams without virtual photons, the non-QED or weak corrections. These weak corrections can conveniently be written in terms of fermion-dependent overall normalizations ρ_f and effective mixing angles s_f^2 in the NC vertices ²⁶:

$$\begin{aligned} J_\nu^{NC} &= \left(\sqrt{2}G_\mu M_Z^2\right)^{1/2} (g_V^f \gamma_\nu - g_A^f \gamma_\nu \gamma_5) \quad (14) \\ &= \left(\sqrt{2}G_\mu M_Z^2 \rho_f\right)^{1/2} \left((I_3^f - 2Q_f s_f^2)\gamma_\nu - I_3^f \gamma_\nu \gamma_5\right). \end{aligned}$$

ρ_f and s_f^2 contain universal parts (i.e. independent of the fermion species) and non-universal parts which explicitly depend on the type of the external fermions. In their leading terms, incorporating also the next order, the parameters are given by

$$\rho_f = \frac{1}{1 - \Delta\rho} + \dots, \quad s_f^2 = s_W^2 + c_W^2 \Delta\rho + \dots \quad (15)$$

with $\Delta\rho$ from Eq. (6).

For the b quark, also the non-universal parts have a strong dependence on m_t resulting from virtual top quarks in the vertex corrections. The difference between the d and b couplings can be parametrized in the following way

$$\rho_b = \rho_d(1 + \tau)^2, \quad s_b^2 = s_d^2(1 + \tau)^{-1} \quad (16)$$

with the quantity

$$\tau = \Delta\tau^{(1)} + \Delta\tau^{(2)} + \Delta\tau^{(\alpha_s)}$$

calculated perturbatively, at the present level comprising: the complete 1-loop order term ²⁷ with x_t from Eq. (5):

$$\Delta\tau^{(1)} = -2x_t - \frac{G_\mu M_Z^2}{6\pi^2 \sqrt{2}} (c_W^2 + 1) \log \frac{m_t}{M_W} + \dots, \quad (17)$$

the leading electroweak 2-loop contribution of $O(G_\mu^2 m_t^4)$ ^{16,28}

$$\Delta\tau^{(2)} = -2x_t^2 \tau^{(2)}, \quad (18)$$

where $\tau^{(2)}$ is a function of M_H/m_t with $\tau^{(2)} = 9 - \pi^2/3$ for $M_H \ll m_t$; the QCD corrections to the leading term of $O(\alpha_s G_\mu m_t^2)$ ²⁹

$$\Delta\tau^{(\alpha_s)} = 2x_t \cdot \frac{\alpha_s}{\pi} \cdot \frac{\pi^2}{3}, \quad (19)$$

and the $O(\alpha_s)$ correction to the $\log m_t/M_W$ term in (17), with a numerically very small coefficient ³⁰.

Asymmetries and mixing angles: The effective mixing angles are of particular interest since they determine the on-resonance asymmetries via the combinations

$$A_f = \frac{2g_V^f g_A^f}{(g_V^f)^2 + (g_A^f)^2}. \quad (20)$$

Measurements of the asymmetries hence are measurements of the ratios

$$g_V^f/g_A^f = 1 - 2Q_f s_f^2 \quad (21)$$

or the effective mixing angles, respectively.

Z width and partial widths: The total Z width Γ_Z can be calculated essentially as the sum over the fermionic partial decay widths

$$\Gamma_Z = \sum_f \Gamma_f + \dots, \quad \Gamma_f = \Gamma(Z \rightarrow f\bar{f}) \quad (22)$$

The dots indicate other decay channels which, however, are not significant. The fermionic partial widths, when expressed in terms of the effective coupling constants read up to 2nd order in the (light) fermion masses:

$$\begin{aligned} \Gamma_f &= \Gamma_0 \left((g_V^f)^2 + (g_A^f)^2 \left(1 - \frac{6m_f^2}{M_Z^2}\right) \right) \cdot \left(1 + Q_f^2 \frac{3\alpha}{4\pi}\right) \\ &+ \Delta\Gamma_{QCD}^f \end{aligned}$$

with

$$\Gamma_0 = N_C^f \frac{\sqrt{2}G_\mu M_Z^3}{12\pi}, \quad N_C^f = 1 \text{ (leptons)}, = 3 \text{ (quarks)}.$$

The QCD correction for the light quarks with $m_q \simeq 0$ is given by

$$\Delta\Gamma_{QCD}^f = \Gamma_0 \left((g_V^f)^2 + (g_A^f)^2 \right) \cdot K_{QCD} \quad (23)$$

with ³¹

$$K_{QCD} = \frac{\alpha_s}{\pi} + 1.41 \left(\frac{\alpha_s}{\pi}\right)^2 - 12.8 \left(\frac{\alpha_s}{\pi}\right)^3 - \frac{Q_f^2 \alpha \alpha_s}{4 \pi^2}.$$

For b quarks the QCD corrections are different due to finite b mass terms and to top quark dependent 2-loop diagrams for the axial part:

$$\Delta\Gamma_{QCD}^b = \Delta\Gamma_{QCD}^d + \Gamma_0 \left[(g_V^b)^2 R_V + (g_A^b)^2 R_A \right] \quad (24)$$

The coefficients in the perturbative expansions

$$R_V = c_1^V \frac{\alpha_s}{\pi} + c_2^V \left(\frac{\alpha_s}{\pi}\right)^2 + c_3^V \left(\frac{\alpha_s}{\pi}\right)^3 + \dots,$$

$$R_A = c_1^A \frac{\alpha_s}{\pi} + c_2^A \left(\frac{\alpha_s}{\pi}\right)^2 + \dots$$

depending on m_b and m_t , are calculated up to third order in the vector and up to second order in the axial part ³².

Radiation of secondary fermions through photons from the primary final state fermions can yield another sizeable contribution to the partial Z widths which, however, is compensated by the corresponding virtual contribution through the dressed photon propagator in the final state vertex correction. For this compensation it is essential that the analysis is inclusive enough, i.e. the cut to the invariant mass of the secondary fermions is sufficiently large ³³.

2.4 Status of the standard model predictions

For a discussion of the theoretical reliability of the standard model predictions one has to consider the various sources contributing to their uncertainties:

The experimental error propagating into the hadronic contribution of $\alpha(M_Z^2)$, Eq. (3), leads to $\delta M_W = 13$ MeV in the W mass prediction, and $\delta \sin^2 \theta = 0.00023$ common to all of the mixing angles, which matches with the future experimental precision.

The uncertainties from the QCD contributions, besides the 3 MeV in the hadronic Z width, can essentially be traced back to those in the top quark loops for the ρ -parameter. They can be combined into the following errors ³⁴, which have improved due to the recently available 3-loop result:

$$\delta(\Delta\rho) \simeq 1.5 \cdot 10^{-4}, \quad \delta s_\ell^2 \simeq 0.0001$$

for $m_t = 174$ GeV, and slightly larger for heavier top.

The size of unknown higher order contributions can be estimated by different treatments of non-leading terms of higher order in the implementation of radiative corrections in electroweak observables (‘options’) and by investigations of the scheme dependence. Explicit comparisons between the results of 5 different computer codes based on on-shell and \overline{MS} calculations for the Z resonance observables are documented in the ‘‘Electroweak Working Group Report’’ ³⁵ in ref. ⁷ (see also the talk by Bardin ⁸). The typical size of the genuine electroweak uncertainties is of the order 0.1%. The following table shows the uncertainty in a selected set of precision observables. In particular for the very precise s_e^2 the theoretical uncertainty is still remarkable. Improvements of the accuracy displayed in table 1 require systematic electroweak and QCD-electroweak 2-loop calculations. As an example for the importance of electroweak non-leading 2-loop effects, an explicit calculation of these terms has been performed

for ρ (the overall normalization) in neutrino scattering ³⁶: they are sizeable and comparable to the $O(G_\mu^2 m_t^4)$ term. Hence, one should take the registered uncertainties also for the Z region very seriously.

Table 1: Largest half-differences among central values (Δ_c) and among maximal and minimal predictions (Δ_g) for $m_t = 175$ GeV, $60 \text{ GeV} < M_H < 1 \text{ TeV}$ and $\alpha_s(M_Z^2) = 0.125$ (from ref. ³⁵)

Observable O	$\Delta_c O$	$\Delta_g O$
M_W (GeV)	4.5×10^{-3}	1.6×10^{-2}
Γ_e (MeV)	1.3×10^{-2}	3.1×10^{-2}
Γ_Z (MeV)	0.2	1.4
s_e^2	5.5×10^{-5}	1.4×10^{-4}
s_b^2	5.0×10^{-5}	1.5×10^{-4}
R_{had}	4.0×10^{-3}	9.0×10^{-3}
R_b	6.5×10^{-5}	1.7×10^{-4}
R_c	2.0×10^{-5}	4.5×10^{-5}
σ_0^{had} (nb)	7.0×10^{-3}	8.5×10^{-3}
A_{FB}^l	9.3×10^{-5}	2.2×10^{-4}
A_{FB}^b	3.0×10^{-4}	7.4×10^{-4}
A_{FB}^c	2.3×10^{-4}	5.7×10^{-4}
A_{LR}	4.2×10^{-4}	8.7×10^{-4}

Low angle Bhabha scattering for a luminosity measurement at 0.1% accuracy still requires more theoretical effort. For a description of the present status see the contributions by Jadach et al. and other authors in ⁷. Presently an accuracy of 0.16% has been claimed ³⁷.

3 Standard model predictions versus data

In table 2 the standard model predictions for Z pole observables and the W mass are put together. The first error corresponds to the variation of m_t in the observed range (1) and $60 < M_H < 1000$ GeV. The second error is the hadronic uncertainty from $\alpha_s = 0.123 \pm 0.006$, as measured by QCD observables at the Z ³⁹. The recent combined LEP results ¹ on the Z resonance parameters, under the assumption of lepton universality, are also shown in table 1, together with s_e^2 from the left-right asymmetry at the SLC ^{40,1}.

The value for the leptonic mixing angle from the left-right asymmetry A_{LR} has come closer to the LEP result, but due to its smaller error the deviation from the cumulative LEP average is still 3σ .

Significant deviations from the standard model predictions are observed in the ratios $R_b = \Gamma_b/\Gamma_{had}$ and $R_c = \Gamma_c/\Gamma_{had}$. The experimental values, together with the top mass (1) from the Tevatron, are compatible with the standard model at a confidence level of less than 1% (see ^{1,38}), enough to claim a deviation from the Standard Model. The other precision observables are in perfect

Table 2: Precision observables: experimental results (from refs. 1,2,3) and standard model predictions.

observable	exp. (1995)	standard model prediction
M_Z (GeV)	91.1884 ± 0.0022	input
Γ_Z (GeV)	2.4963 ± 0.0032	$2.4976 \pm 0.0077 \pm 0.0033$
σ_0^{had} (nb)	41.4882 ± 0.078	$41.457 \pm 0.011 \pm 0.076$
Γ_{had}/Γ_e	20.788 ± 0.032	$20.771 \pm 0.019 \pm 0.038$
Γ_{inv} (MeV)	499.9 ± 2.5	501.6 ± 1.1
$\Gamma_b/\Gamma_{had} = R_b$	0.2219 ± 0.0017	0.2155 ± 0.0004
$\Gamma_c/\Gamma_{had} = R_c$	0.1540 ± 0.0074	0.1723 ± 0.0002
A_b	0.841 ± 0.053	0.9346 ± 0.0006
ρ_ℓ	1.0044 ± 0.0016	1.0050 ± 0.0023
s_ℓ^2 (LEP)	0.23186 ± 0.00034	0.2317 ± 0.0012
$s_e^2(A_{LR})$	0.23049 ± 0.00050	0.2317 ± 0.0012
LEP+SLC	0.23143 ± 0.00028	
M_W (GeV)	80.26 ± 0.16	80.36 ± 0.18

agreement with the Standard Model. Note that the experimental value for ρ_ℓ exhibits the presence of genuine electroweak corrections by nearly 3 standard deviations.

The W mass prediction is obtained by Eqs. (10-12) from M_Z, G_μ, α and M_H, m_t . The quantity s_W^2 resp. the ratio M_W/M_Z is indirectly measured in deep-inelastic neutrino scattering, in particular in the NC/CC neutrino cross section ratio for isoscalar targets. The present world average from CCFR, CDHS and CHARM, including the new CCFR result ⁴⁴

$$s_W^2 = 1 - M_W^2/M_Z^2 = 0.2253 \pm 0.0047$$

is fully consistent with the direct vector boson mass measurements and with the standard theory.

Standard model fits: Assuming the validity of the standard model a global fit to all electroweak results from LEP, SLD, $p\bar{p}$ and νN constrains the parameters m_t, α_s as follows: ^{1,2,45}

$$m_t = 178 \pm 8_{-20}^{+17} \text{ GeV}, \quad \alpha_s = 0.123 \pm 0.004 \pm 0.002 \quad (25)$$

with $M_H = 300$ GeV for the central value. The second error is from the variation of M_H between 60 GeV and 1 TeV. The fit results include the uncertainties of the standard model calculations.

The indirect determination of the W mass from LEP/SLD data,

$$M_W = 80.359 \pm 0.055_{-0.024}^{+0.013} \text{ GeV},$$

is in best agreement with the direct measurement (see table 2). Moreover, the value obtained for α_s at M_Z coincides with the one measured from others than electroweak observables at the Z peak ³⁹.

The main Higgs dependence of the electroweak predictions is only logarithmic in the Higgs mass. Hence, the sensitivity of the data to M_H is not very pronounced. Using the Tevatron value for m_t as an additional experimental constraint, the electroweak fit to all data yields $M_H < 600$ GeV with approximately 95% C.L. ^{1,2}. Similar results with bounds on M_H which are 100-200 MeV higher, based on the electroweak data from the winter conferences, have been obtained in ^{42,43}.

Low energy results: The cross section for μ -neutrino electron scattering and the electroweak mixing angle measured by the CHARM II Collaboration ⁴⁶ agree with the standard model values:

$$\begin{aligned} \sigma(\nu e)/E_\nu &= 16.51 \pm 0.93) \cdot 10^{-42} \text{ cm}^2 \text{ GeV}^{-1} \\ (\text{SM} &: 17.23 \cdot 10^{-42}) \\ \sin^2 \theta_{\nu e} &= 0.2324 \pm 0.0083. \end{aligned} \quad (26)$$

The mixing angle coincides with the result on s_ℓ^2 from the Z , table 2, as expected by the theory. The major loop contributions in the difference, the different scales and the neutrino charge radius, largely cancel each other by numerical coincidence ⁴¹.

The recent results from the CLEO Collaboration ⁴⁷ on the flavor changing B decays, the branching ratio $BR(B \rightarrow X_s \gamma) = (2.32 \pm 0.67) \cdot 10^{-4}$ and the inclusive photon spectrum, are fully consistent with the standard model predictions based on the loop induced $b \rightarrow s \gamma$ transition ⁴⁸.

4 Future tests of the Standard Model

4.1 Higgs bosons

The minimal model with a single scalar doublet is the simplest way to implement the electroweak symmetry breaking. The experimental result that the ρ -parameter is very close to unity is a natural feature of models with doublets and singlets. In the standard model, the mass M_H of the Higgs boson appears as the only additional parameter beyond the vector boson and fermion masses. M_H cannot be predicted but has to be taken from experiment. The present lower limit (95% C.L.) from the search at LEP⁴⁹ is 65 GeV. Indirect determinations of M_H from precision data have already been discussed in section 3. The indirect mass bounds depend sensitively on small changes in the input data, and their reliability suffers at present from averaging data points which fluctuate by several standard deviations. As a general feature, it appears that the data prefer light Higgs bosons.

There are also theoretical constraints on the Higgs mass from vacuum stability and absence of a Landau pole⁵⁰, and from lattice calculations⁵¹. A recent calculation of the decay width for $H \rightarrow W^+W^-$ in the large M_H limit in 2-loop order⁵² has shown that the 2-loop contribution exceeds the 1-loop term in size (same sign) for $M_H > 930$ GeV. The requirement of applicability of perturbation theory therefore puts a stringent upper limit on the Higgs mass.

Higgs boson searches at LEP2 and future high energy hadron and e^+e^- colliders require precise predictions for the Higgs production and decay signatures together with detailed background studies. Improved calculations for the most relevant Higgs decays including higher order contributions have become available for

- $H \rightarrow b\bar{b}$ for $M_H < 2m_t$ in order $O(\alpha_s^2)$ ^{53,54}, and $O(\alpha_s G_\mu m_t^2)$ ⁵⁵ as well as $O(\alpha_s^2 G_\mu m_t^2)$ ⁵⁶;
- $H \rightarrow gg(g)$ up to order $O(\alpha_s^3)$ ⁵³;
- $H \rightarrow ZZ, WW$ for large M_H in electroweak 2-loop order⁵².

The QCD corrections for Higgs production at the LHC has been completed⁵⁷, with the result of a significant enhancement by the next order QCD contributions.

Higgs signal versus background studies were performed for the process $e^+e^- \rightarrow b\bar{b}l\bar{l}$ at tree level order, both (semi-)analytically⁵⁸ and by Monte Carlo methods⁵⁹. Current work is going on in the topical LEP200 Workshop. An overview with more details on the present theoretical status of Higgs physics can be found in these proceedings⁶⁰.

4.2 W bosons

W mass measurements at LEP2 with an error of about 40 MeV and tests of the trilinear vector boson self couplings⁶¹ require standard model calculations for the process $e^+e^- \rightarrow W^+W^- \rightarrow 4f$ and the corresponding 4-fermion

background processes at the accuracy level of 1%. A status report can be found in ref.⁶². One of the specific problems in the theoretical description of the production process for off-shell W bosons is the presence of the width term in the W propagator which violates gauge invariance, yielding gauge dependent amplitudes. As a solution it has been proposed⁶³ to take into account also the imaginary part in the $WW\gamma$ vertex from the light fermion triangle loops. This prescription is in accordance with gauge invariance and cures the Ward identities between 2- and 3-point functions involving W^\pm and γ .

4.3 g-2 for muons

The anomalous magnetic moment of the muon,

$$a_\mu = \frac{g_\mu - 2}{2} \quad (27)$$

provides a precision test of the standard model at low energies. Within the present experimental accuracy of $\Delta a_\mu = 840 \cdot 10^{-11}$, theory and experiment are in best agreement, but the electroweak loop corrections are still hidden in the noise. A new experiment, E 821 at Brookhaven National Laboratory⁶⁴, is being prepared for 1996 to reduce the experimental error down to 40 ± 10^{-11} and hence will become sensitive to the electroweak loop contribution, which at the 1-loop level⁶⁵ amounts to $195 \cdot 10^{-11}$.

For this reason the standard model prediction has to be known with comparable precision. Recent theoretical work has contributed to reduce the theoretical uncertainty by calculating the electroweak 2-loop terms^{66,67,68} and updating the contribution from the hadronic photonic vacuum polarization⁹

$$a_\mu^{had}(\text{vacuum pol.}) = (7024 \pm 153) \cdot 10^{-11}$$

which agrees within the error with the result of⁶⁹. The main sources for the theoretical error at present are the hadronic vacuum polarization and the light-by-light scattering mediated by quarks, as part of the 3-loop hadronic contribution^{70,71}. Table 3 shows the breakdown of a_μ . The hadronic part is supplemented by the higher order α^3 vacuum polarization effects⁷² but is without the light-by-light contribution, where the situation is unclear at present.

The 2-loop electroweak contribution turns out to be as big in size as the expected experimental error. The dominating theoretical uncertainty at present is the error in the hadronic vacuum polarization which can only be improved by new measurements of the cross section for $e^+e^- \rightarrow \text{hadrons}$ in the low energy range. But also the contribution involving light-by-light scattering needs clarification in order to reduce the theoretical error.

Table 3: Contributions Δa_μ to the muonic anomalous magnetic moment and their theoretical uncertainties, in units of 10^{-11} .

source	Δa_μ	error
QED ⁷³	116584708	5
hadronic ^{9,72}	6934	153
EW, 1-loop ⁶⁵	195	
EW, 2-loop ⁶⁷ (anomaly graphs)	-17	3
EW, 2-loop ⁶⁶ (all fermionic graphs)	-23	3
EW, 2-loop ⁶⁸ (bosonic graphs)	-17	2
light-by-light ⁷⁰	-36	16
light-by-light ⁷¹	-124	50
future experiment		40

5 Precision data and “New Physics”

New physics in R_b ? If the observed difference between the measured and calculated values of R_b is explained by a non-standard contribution $\Delta\Gamma_b$ to the partial width $\Gamma(Z \rightarrow b\bar{b})$, then also other hadronic quantities like Γ_Z, R_{had}, \dots are increased unless the value of α_s is reduced simultaneously. Including the new physics $\Delta\Gamma_b$ as an extra free parameter in the fit yields the values ^{1,2}:

$$\alpha_s = 0.102 \pm 0.008, \quad \Delta\Gamma_b = 11.7 \pm 3.8 \pm 1.4 \text{MeV}.$$

The top mass is affected only marginally, shifting the central value by +3 GeV, but the impact on α_s is remarkable.

Virtual New Physics: The generalization of the aforementioned method consists in the parametrization of the radiative corrections originating from the vector boson self-energies in terms of the static ρ -parameter $\Delta\rho(0) \equiv \epsilon_1$ and two other combinations of self-energies, ϵ_2 and ϵ_3 ^{74,75}. This allows a more general analysis of the electroweak data which accommodates extensions of the minimal model affecting only the vector boson self-energies. A further quantity ϵ_b has been introduced ⁷⁵ in order to parametrize specific non-universal left handed contributions to the $Zb\bar{b}$ vertex via

$$g_A^b = g_A^d(1+\epsilon_b), \quad g_V^b/g_A^b = (1 - \frac{4}{3}s_d^2 + \epsilon_b)(1+\epsilon_b)^{-1}. \quad (28)$$

There is a wide literature ⁷⁶ in this field with various conventions.

Phenomenologically, the ϵ_i are parameters which can be determined experimentally from the electroweak precision data. An updated analysis ⁷⁷ on the basis of the

recent electroweak results presented at this conference ¹ yields for ϵ_b the value

$$\epsilon_b = 9.9 \pm 4.5 \quad (\text{SM} : -6.6)$$

The large difference to the standard model value is an another way of visualizing the deviation between the measured and predicted number for the ratio R_b (table 2).

The minimal supersymmetric standard model (MSSM): The MSSM deserves a special discussion as the most predictive framework beyond the minimal model. Its structure allows a similarly complete calculation of the electroweak precision observables as in the standard model in terms of one Higgs mass (usually taken as M_A) and $\tan\beta = v_2/v_1$, together with the set of SUSY soft breaking parameters fixing the chargino/neutralino and scalar fermion sectors. It has been known since quite some time ⁷⁸ that light non-standard Higgs bosons as well as light stop and charginos predict larger values for the ratio R_b and thus diminish the observed difference ^{79,81,82,83}. Complete 1-loop calculations are meanwhile available for Δr ⁸⁰ and for the Z boson observables ^{81,82,83}.

The main results in view of the recent precision data are:

- R_c can hardly be moved towards the measured range.
- R_b can come closer to the measured value, in particular for light \tilde{t}_R and light charginos.
- α_s turns out to be smaller than in the minimal model because of the reasons explained in the beginning of this section.
- There are strong constraints from the other precision observables which forbid parameter configurations shifting R_b into the observed 1σ range.

For obtaining the optimized SUSY parameter set, therefore, a global fit to all the electroweak precision data (including the top mass measurements) has to be performed, as done in refs. ^{82,84}. Figure 1 displays the experimental data normalized to the best fit results in the SM and MSSM, with the data from this conference ⁸⁴. For the SM, α_s identified with the experimental number, therefore the corresponding result in Figure 1 is centered at 1. The most relevant conclusions are:

- The difference between the experimental and theoretical value of R_b is diminished by a factor $\simeq 1/2$,
- the central value for the strong coupling is $\alpha_s = 0.110$ and thus is very close to the value obtained from deep inelastic scattering,
- the other observables are practically unchanged,
- the χ^2 of the fit is slightly better than in the minimal model.

6 Conclusions

The experimental data for tests of the standard model have achieved an impressive accuracy. In the meantime,




Figure 1: Experimental data normalized to the best fit results in the SM and MSSM

many theoretical contributions have become available to improve and stabilize the standard model predictions. To reach, however, a theoretical accuracy at the level 0.1% or below, new experimental data on $\Delta\alpha$ and more complete electroweak 2-loop calculations are required. The observed deviations of several σ 's in R_b, R_c, A_{LR} reduce the quality of the standard model fits significantly, but the indirect determination of m_t is remarkably stable. Still impressive is the perfect agreement between theory and experiment for the whole set of the other precision observables. SUSY can improve the situation due to an enhancement of R_b by new particles in the range of 100 GeV or even below, but it is not possible to accommodate R_c . Within the MSSM analysis, the value for α_s is close to the one from deep-inelastic scattering. In the QCD sector, a deviation from the theoretical expectation in the inclusive jet cross section at the Tevatron has been reported⁸⁵ in terms of a significant excess of jets with $E_T > 200$ GeV compared to the NLO QCD prediction. It is, however, too early to draw conclusions about new physics from that.

Acknowledgments

I wish to thank D. Bardin, W. de Boer, A. Czarnecki, A. Dabelstein, R. Ehret, S. Meyer, A. Olshevsky, G. Passarino, D. Schaile for instructive discussions and for support in preparing this review.

References

1. A. Olshevsky, these proceedings (plenary talk)
2. The LEP experiments and the LEP Electroweak Working Group, internal note LEPEWWG/51-01 (1995), updated for this conference; The LEP Electroweak Heavy Flavour Working Group and the SLD Heavy Flavour Working Group, Internal Note LEPHF/95-02, SLD Physics Note 39
3. UA2 Collaboration, J. Alitti et al., Phys. Lett. B **276** (1992) 354; CDF Collaboration, F. Abe et al., Phys. Rev. D **43** (1991) 2070; D0 Collaboration, C.K. Jung, talk at the 27th International Conference on High Energy Physics, Glasgow 1994; CDF Collaboration, F. Abe et al., FERMILAB-PUB-95/033-E; FERMILAB-PUB-95/035-E (1995)
4. CDF Collaboration, F. Abe et al., FERMILAB-PUB-95/022-E (1995)
5. D0 Collaboration, S. Abachi et al., FERMILAB-PUB-95/028-E (1995)
6. A. Menzione, these proceedings (plenary talk)
7. Reports of the Working Group on Precision Calculations for the Z Resonance, CERN 95-03 (1995), Eds. D. Bardin, W. Hollik, G. Passarino
8. D. Bardin, these proceedings (talk at PA1)
9. S. Eidelman, F. Jegerlehner, Z. Phys. C **67** (1995) 585
10. H. Burkhardt, B. Pietrzyk, Phys. Lett. B **356** (1995) 398
11. F. Jegerlehner, Progress in Particle and Nuclear Physics **27** (1991) 1, updated from: H. Burkhardt, F. Jegerlehner, G. Penso, C. Verzegnassi, Z. Phys. C **43** (1989) 497;
12. M.L. Swartz, preprint SLAC-PUB-95-7001 (1995)
13. A.D. Martin, D. Zeppenfeld, Phys. Lett. B **345** (1995) 558
14. M. Veltman, Nucl. Phys. B **123** (1977) 89; M.S. Chanowitz, M.A. Furman, I. Hinchliffe, Phys. Lett. B **78** (1978) 285
15. J.J. van der Bij, F. Hoogeveen, Nucl. Phys. B **283** (1987) 477
16. R. Barbieri, M. Beccaria, P. Ciafaloni, G. Curci, A. Vicere, Phys. Lett. B **288** (1992) 95; Nucl. Phys. B **409** (1993) 105; J. Fleischer, F. Jegerlehner, O.V. Tarasov, Phys. Lett. B **319** (1993) 249
17. A. Djouadi, C. Verzegnassi, Phys. Lett. B **195** (1987) 265
18. L. Avdeev, J. Fleischer, S. M. Mikhailov, O.

- Tarasov, Phys. Lett. B **336** (1994) 560; E: Phys. Lett. B **349** (1995) 597; K.G. Chetyrkin, J.H. Kühn, M. Steinhauser, Phys. Lett. B **351** (1995) 331
19. A. Sirlin, Phys. Rev. **22** (1980) 971; W.J. Marciano, A. Sirlin, Phys. Rev. **22** (1980) 2695
 20. W.J. Marciano, Phys. Rev. D **20** (1979) 274
 21. M. Consoli, W. Hollik, F. Jegerlehner, Phys. Lett. B **227** (1989) 167
 22. A. Djouadi, Nuovo Cim. A **100** (1988) 357; D. Yu. Bardin, A.V. Chizhov, Dubna preprint E2-89-525 (1989); B.A. Kniehl, Nucl. Phys. B **347** (1990) 86; F. Halzen, B.A. Kniehl, Nucl. Phys. B **353** (1991) 567; A. Djouadi, P. Gambino, Phys. Rev. D **49** (1994) 3499
 23. B.A. Kniehl, J.H. Kühn, R.G. Stuart, Phys. Lett. B **214** (1988) 621; B.A. Kniehl, A. Sirlin, Nucl. Phys. B **371** (1992) 141; Phys. Rev. D **47** (1993) 883; S. Fanchiotti, B.A. Kniehl, A. Sirlin, Phys. Rev. D **48** (1993) 307
 24. K.G. Chetyrkin, J.H. Kühn, M. Steinhauser, preprint KA-TTP95-13 (1995)
 25. A. Sirlin, Phys. Rev. D **29** (1984) 89
 26. D. Bardin, M.S. Bilenky, G.V. Mithselmakher, T. Riemann, M. Sachwitz, Z. Phys. C **44** (1989) 493; W. Hollik, Munich preprints MPI-Ph/93-21, MPI-Ph/93-22 (1993), in: Precision Tests of the Standard Model, Ed. P. Langacker, World Scientific, Singapore 1995
 27. A.A. Akhundov, D.Yu. Bardin, T. Riemann, Nucl. Phys. B **276** (1986) 1; W. Beenakker, W. Hollik, Z. Phys. C **40** (1988) 141; J. Bernabeu, A. Pich, A. Santamaria, Phys. Lett. B **200** (1988) 569;
 28. A. Denner, W. Hollik, B. Lampe, Z. Phys. C **60** (1993) 193
 29. J. Fleischer, F. Jegerlehner, P. Rączka, O.V. Tarasov, Phys. Lett. B **293** (1992) 437; G. Buchalla, A.J. Buras, Nucl. Phys. B **398** (1993) 285; G. Degrossi, Nucl. Phys. B **407** (1993) 271; K.G. Chetyrkin, A. Kwiatkowski, M. Steinhauser, Mod. Phys. Lett. A **8** (1993) 2785
 30. A. Kwiatkowski, M. Steinhauser, Phys. Lett. B **344** (1995) 359; S. Peris, A. Santamaria, CERN-TH-95-21 (1995)
 31. K.G. Chetyrkin, A.L. Kataev, F.V. Tkachov, Phys. Lett. B **85** (1979) 277; M. Dine, J. Sapirstein, Phys. Rev. Lett. **43** (1979) 668; W. Celmaster, R. Goncalves, Phys. Rev. Lett. **44** (1980) 560; S.G. Gorishny, A.L. Kataev, S.A. Larin, Phys. Lett. B **259** (1991) 144; L.R. Surguladze, M.A. Samuel, Phys. Rev. Lett. **66** (1991) 560; A. Kataev, Phys. Lett. B **287** (1992) 209
 32. T.H. Chang, K.J.F. Gaemers, W.L. van Neerven, Nucl. Phys. B **202** (1982) 407; J.H. Kühn, B.A. Kniehl, Phys. Lett. B **224** (1990) 229; Nucl. Phys. B **329** (1990) 547; K.G. Chetyrkin, J.H. Kühn, Phys. Lett. B **248** (1992) 359; K.G. Chetyrkin, J.H. Kühn, A. Kwiatkowski, Phys. Lett. B **282** (1992) 221; K.G. Chetyrkin, A. Kwiatkowski, Phys. Lett. B **305** (1993) 285; Karlsruhe preprint TTP93-24 (1993); K.G. Chetyrkin, Karlsruhe preprint TTP93-5 (1993); K.G. Chetyrkin, J.H. Kühn, A. Kwiatkowski, in [7], p. 175; S. Larin, T. van Ritbergen, J.A.M. Vermaseren, ibidem, p. 265; Phys. Lett. B **320** (1994) 159
 33. A. Hoang, J.H. Kühn, T. Teubner, Nucl. Phys. B **455** (1995) 3; **452** (1995) 173
 34. B.A. Kniehl, in [7], p. 299
 35. D. Bardin et al., in [7], p. 7
 36. G. Degrossi, S. Fanchiotti, F. Feruglio, P. Gambino, A. Vicini, in [7], p. 163
 37. S. Jadach et al., Phys. Lett. B **353** (1995) 362
 38. T. Behnke, these proceedings (talk at PA8)
 39. S. Bethke, in: Proceedings of the Tennessee International Symposium on Radiative Corrections, Gatlinburg 1994, Ed. B.F.L. Ward, World Scientific 1995
 40. SLD Collaboration, K. Abe et al., Phys. Rev. Lett. **73** (1994) 25; M. Woods (SLD Collaboration), these proceedings (talk at PA1)
 41. S. Sarantakos, A. Sirlin, Nucl. Phys. B **217** (1983) 84; D.Yu. Bardin V.A. Dokuchaeva, Nucl. Phys. B **246** (1984) 221; M. Böhm, W. Hollik, H. Spiesberger, Z. Phys. C **27** (1985) 523
 42. P.Chankowski, S. Pokorski, Phys. Lett. B **356** (1995) 307
 43. J. Ellis, G.L. Fogli, E. Lisi, CERN-TH-95-202 (1995)
 44. CDHS Collaboration, H. Abramowicz et al., Phys. Rev. Lett. **57** (1986) 298; A. Blondel et al, Z. Phys. C **45** (1990) 361; CHARM Collaboration, J.V. Alaby et al., Phys. Lett. B **177** (1987) 446; Z. Phys. C **36** (1987) 611; CHARM-II Collaboration, D. Geiregat et al., Phys. Lett. B **247** (1990) 131; Phys. Lett. B **259** (1991) 499; CCFR Collaboration, C.G. Arroyo et al., Phys. Rev. Lett. **72** (1994) 3452; D. Harris (CCFR Coll.), these proceedings (talk at PA1)
 45. J. Busenitz, these proceedings (talk at PA1)
 46. CHARM II Collaboration, P. Vilain et al., Phys. Lett. B **335** (1994) 246; **345** (1995) 115
 47. CLEO Collaboration, M.S. Alam et al., Phys. Rev. Lett. **71** (1993) 674; **74** (1995) 2885
 48. A.J. Buras, M. Misiak, M. Münz, S. Pokorski, Nucl. Phys. B **424** (1994) 374; M. Ciuchini et al., Phys. Lett. B **334** (1994) 137; A. Ali, C. Greub, DESY 95-117/SLAC-PUB-95-6940
 49. J.-F. Grivaz, these proceedings (plenary talk)
 50. M. Lindner, M. Sher, H. Zaglauer, Phys. Lett. B **1989** (1989) 139
 51. Kuti et al., Phys. Rev. Lett. **61** (1988) 678; Hasenfratz et al., Nucl. Phys. B **317** (1989) 81; M. Lüscher,

- P. Weisz, Nucl. Phys. B **318** (1989) 705
52. A. Ghinculov, Nucl. Phys. B **455** (1995) 21
53. S. Larin, T. van Ritbergen, J. Vermaseren, NIKHEF-H/95-027 (1995)
54. K. Chetyrkin, A. Kwiatkowski, LBL-37269, TTP 95-23 (1995)
55. A. Kwiatkowski, M. Steinhauser, TTP 94-7 (1994); B. Kniehl, M. Spira, DESY 95-008 (1995)
56. B. Kniehl, M. Steinhauser, FERMILAB-PUB-95/197-T (1995)
57. M. Spira, A. Djouadi, D. Graudenz, P.M. Zerwas, DESY 94-123, CERN-TH/95-30 (1995)
58. D. Bardin, A. Leike, T. Riemann, DESY 95-057 (1995)
59. E. Boos, M. Sachwitz, H. Schreiber, S. Sichanin, Z. Phys. C **61** 1994) 675; **64** (1994) 391; M. Dubinin, V. Edneral, Y. Kurihara, Y. Shimizu, Phys. Lett. B **329** (1994) 379; G. Montagna, O. Nicosini, F. Piccinini, CERN-TH/95-08 (1995)
60. H. Haber, these proceedings (talk at PA1)
61. U. Baur, these proceedings (talk at PA 6)
62. W. Beenakker, F.A. Berends, et al., *WW* cross sections and distributions, to be published in the proceedings of the LEP2 Workshop, CERN Yellow Report, Eds. G. Altarelli et al.
63. E.N. Argyres et al., Phys. Lett. B **358** (1995) 339; U. Baur, D. Zeppenfeld, Phys. Rev. Lett. **75** (1995) 1002; C. Papadopoulos, Phys. Lett. B **352** (1995) 144
64. V.W. Hughes, in: *Frontiers of High Energy Spin Physics*, Universal Academy Press, Tokyo 1992, Ed. T Hasegawa (p. 717)
65. K. Fujikawa, B.W. Lee, A.I. Sanda, Phys. Rev. D **6** (1972) 2923; R. Jackiw, S. Weinberg, Phys. Rev. D **5** (1972) 2473; G. Altarelli, N. Cabbibo, L. Maiani, Phys. Lett. B **40** (1972) 415; I. Bars, M. Yoshimura, Phys. Rev. D **6** (1972) 374; W. Bardeen, R. Gastmans, B. Lautrup, Nucl. Phys. B **46** (1972) 315
66. A. Czarnecki, B. Krause, W. Marciano, Phys. Rev. D **52** (1995) 2619
67. S. Peris, M. Perrottet, E. de Rafael, Phys. Lett. B **355** (1995) 523
68. A. Czarnecki, B. Krause, W. Marciano, KA-TTP-34-1995 (1995)
69. S. Dubničká, A.Z. Dubničková, P. Striženeč, Bratislava preprint (contributed to this conference)
70. M. Hayakawa, T. Kinoshita, A.I. Sanda, Phys. Rev. Lett. **75** (1995) 790
71. J. Bijnens, E. Pallante, J. Prades, preprint NORDITA-95-75-N (1995)
72. T. Kinoshita, B. Nizic, Y. Okamoto, Phys. Rev. D **31** (1985) 2108; T. Kinoshita, W. Marciano, in: *Quantum Electrodynamics*, Ed. T. Kinoshita, World Scientific 1990 (p. 419)
73. T. Kinoshita, Phys. Rev. D **47** (1993) 5013; S. Laporta, E. Remiddi, Phys. Lett. B **301** (1992) 440; S. Laporta, Phys. Lett. B **312** (1993) 495;
74. G. Altarelli, R. Barbieri, Phys. Lett. B **253** (1991) 161; G. Altarelli, R. Barbieri, S. Jadach, Nucl. Phys. B **269** (1992) 3; E: Nucl. Phys. B **276** (1992) 444
75. G. Altarelli, R. Barbieri, F. Caravaglios, Nucl. Phys. B **405** (1993) 3; CERN-TH.6895/93 (1993)
76. M.E. Peskin, T. Takeuchi, Phys. Rev. Lett. **65** (1990) 964; W.J. Marciano, J.L. Rosner, Phys. Rev. Lett. **65** (1990) 2963; B.W. Lynn, M.E. Peskin, R.G. Stuart, in: *Physics with LEP*, eds. J. Ellis and R. Peccei, CERN 86-02 (1986); D.C. Kennedy, P. Langacker, Phys. Rev. Lett. **65** (1990) 2967; R. Barbieri, M.Frigeni, F. Caravaglios, Phys. Lett. B **279** (1992) 169; V.A. Novikov, L.B. Okun, M.I. Vysotsky, CERN-TH.6943/93 (1993), M. Bilenky, K. Kolodziej, M. Kuroda, D. Schildknecht, Phys. Lett. B **319** (1993) 319; S. Dittmaier, D. Schildknecht, M. Kuroda, Nucl. Phys. B **448** (1995) 3
77. F. Caravaglios, these proceedings (talk at PA1)
78. A. Denner, R. Guth, W. Hollik, J.H. Kühn, Z. Phys. C **51** (1991) 695; J. Rosiek, Phys. Lett. B **252** (1990) 135; F. Cornet, W. Hollik, W. Mösle, Nucl. Phys. B **428** (1994) 61; M. Bouleware, D. Finnell, Phys. Rev. D **44** (1991) 2054
79. G. Altarelli, R. Barbieri, F. Caravaglios, CERN-TH.7536/94 (1994); C.S. Lee, B.Q. Hu, J.H. Yang, Z.Y. Fang, J. Phys. G **19** (1993) 13; Q. Hu, J.M. Yang, C.S. Li, Comm. Theor. Phys. **20** (1993) 213; J.D. Wells, C. Kolda, G.L. Kane, Phys. Lett. B **338** (1994) 219; G.L. Kane, R.G. Stuart, J.D. Wells, Phys. Lett. B **354** (1995) 350
80. P. Chankowski, A. Dabelstein, W. Hollik, W. Mösle, S. Pokorski, J. Rosiek, Nucl. Phys. B **417** (1994) 101; D. Garcia, J. Solà, Mod. Phys. Lett. A **9** (1994) 211
81. D. Garcia, R. Jiménez, J. Solà, Phys. Lett. B **347** (1995) 309; **347** (1995) 321; D. Garcia, J. Solà, Phys. Lett. B **357** (1995) 349
82. P. Chankowski, S. Pokorski, preprint IFT-UW-95/5 (1995); P. Chankowski, these proceedings (talk at PA9)
83. A. Dabelstein, W. Hollik, W. Mösle, preprint KA-TP-5-1995 (1995), Proceedings of the Ringberg Workshop "Perspectives for Electroweak Interactions in e^+e^- Collisions", Ringberg Castle, February 1995, Ed. B.A. Kniehl, World Scientific 1995 (p. 345)
84. W. De Boer, S. Meyer, A. Dabelstein, W. Hollik, W. Mösle (to be published)
85. E. Buckley-Geer (CDF Collaboration), these proceedings (talk at PA10)

Tests of the Standard Model *

W. HOLLIK

Institut für Theoretische Physik

Universität Karlsruhe

D-76128 Karlsruhe, Germany

Abstract

In this conference report a summary is given on the recent theoretical work that has contributed to improve the theoretical predictions for testing the standard model in present and future experiments. Precision calculations for the Z resonance are reviewed and the status of the standard model is discussed in the light of the recent top discovery and of the results from precision experiments. Furthermore, theoretical progress for the Higgs search, for W physics at LEP II, and for the anomalous magnetic moment of the muon is summarized. New Physics beyond the minimal model is briefly discussed, in particular the minimal supersymmetric standard model in view of the recent electroweak precision data.

*plenary talk at the International Europhysics Conference on High Energy Physics, 27 July - 2 August 1995, Brussels, Belgium