

Hyperfine Splittings in Heavy Quarkonia From a New Improved Spin-Dependent Potential *

Robert J. Oakes
Department of Physics and Astronomy
Northwestern University
Evanston, IL 60208 USA
E-mail: oakes@fnalv.fnal.gov

ABSTRACT

A formula for an improved spin-dependent potential between a heavy quark and a heavy antiquark was developed using Heavy Quark Effective Theory techniques. The leading logarithmic quark mass terms emerging from loop contributions were explicitly extracted and summed up. There is no renormalization scale dependence in this new formula and it includes both the Eichten-Feinberg formula as well as the one-loop QCD result as special cases. The hyperfine splittings were calculated using the spin-spin part of this improved formula. For charmonium the $J/\psi-\eta_c$ S -wave splitting agrees well with the data while the energy difference ΔM_P between the center of gravity of the $1^3P_{0,1,2}$ states and the 1^1P_1 state has the correct sign but is somewhat larger than the experimental value; however, there are also several other contributions to ΔM_P of comparable magnitude (~ 1 MeV), which we discuss. The corresponding mass differences for the $b\bar{b}$ and $b\bar{c}$ mesons are also predicted.

Recently, significant progress has been made in the theoretical study of the spin-dependent potential between a heavy quark and a heavy antiquark. An improved formula for the spin-dependent potential was developed ¹ and used to calculate the hyperfine splittings in the $c\bar{c}$, $b\bar{b}$, and $b\bar{c}$ systems ². The improved spin dependent potentials were derived from QCD first principles using the techniques of the Heavy Quark Effective Theory (HQET) ³. The spin-dependent potential was separated into short distance parts involving Wilson coefficients and long distance parts which were expressed in terms of gauge invariant correlation functions of the color-electric and color-magnetic fields weighted by the Wilson loop path integral ⁴. If the tree level values for the Wilson coefficients are used the potential reduces to Eichten's and Feinberg's result ⁴. And using the one-loop values of the Wilson coefficients, also calculating the correlation functions to one-loop in perturbation theory, the spin-dependent potential at the one-loop level in perturbative QCD ^{5,6} is recovered. However, the leading logarithmic terms appearing in perturbative calculations were also summed

*Presented at the International Symposium on Heavy Flavor and Electroweak Theory, Beijing, 16-19 August 1995.

up in Ref. [1] using the Renormalization Group Equation (RGE) to obtain a scale independent result. Therefore, the spin-dependent heavy quark-antiquark potential derived in Ref. [1] is scale-independent and thus improves upon and generalizes both Eichten's and Feinberg's result ⁴ and the one-loop perturbative result ^{5,6}. In addition, this improved result ¹ satisfies all the general relations among the different parts of the spin-dependent potential ⁷.

The general formula for the spin-spin part of the renormalization-group-improved spin-dependent potential that was derived in Ref. [1] was used to calculate the hyperfine spin splittings in the $c\bar{c}$, $b\bar{b}$, and $c\bar{b}$ systems in Ref. [2]. Since the spin-spin potential is a short distance feature, perturbation theory can reliably be used in the calculation. The result for the $1^3S_1 - 1^1S_0$ splitting between the J/ψ and the η_c agrees well with the experimental value ⁸ and the predictions for the mass differences $\psi' - \eta'_c$, $\Upsilon(1S) - \eta_b$, $\Upsilon(2S) - \eta'_b$, and $B_c^* - B_c$ are reasonable. However, the contribution to the P -wave energy difference, ΔM_p , between the center of the gravity of $1^3P_{0,1,2}$ states and the 1^1P_1 state, while having the correct sign, is somewhat larger than the experimental data ⁸. That is, when the contributions of the leading logarithmic terms are summed up and included, the agreement with that data is not as good as when only the one-loop perturbative spin-spin potential, in which the leading logarithmic contributions are not summed up and included, is used. There are several other contributions to the rather small energy difference ΔM_p (~ 1 MeV) which estimates indicate are of the same order of magnitude as the spin-spin contribution. It therefore appears that the agreement of the one-loop perturbative result with the data is probably fortuitous.

In the derivation ¹ the renormalized two-particle effective Lagrangian was first calculated to order $1/m^2$. Then, treating the terms of higher order in $1/m$ in the effective Lagrangian as perturbations, the four point Green's function on the Wilson loop ⁹ with the time interval T was calculated in the limit where $m \rightarrow \infty$ first followed by $T \rightarrow \infty$ ^{4,10}. In this limit, using standard perturbative methods, the large T behavior of the Green's function is of the form

$$I \propto e^{-T\epsilon(m,r)}. \quad (1)$$

From Eq. (1) $\epsilon(m,r)$, the potential energy between the quark and the antiquark, can be extracted. Expanding $\epsilon(m,r)$ in powers of $1/m$ each of the spin-dependent potentials can be factorized into a short distance part, involving Wilson coefficients, and a long distance part, which can be expressed in terms of correlation functions of the color-electric and color-magnetic fields weighted by the Wilson-loop integral. Using the notation of Ref. [1], the resulting spin-spin potential is

$$\Delta H_{ss}(m_1, m_2, r) = \frac{\mathbf{S}_1 \cdot \mathbf{S}_2}{3m_1m_2} \left[c_3(\mu, m_1)c_3(\mu, m_2)V_4(\mu, r) - 6N_c g_s^2(\mu)d(\mu)\delta^3(\mathbf{r}) \right] \quad (2)$$

where m_1 , m_2 , and \mathbf{S}_1 , \mathbf{S}_2 are the masses and the spins of the heavy quark and the antiquark, respectively, μ is the renormalization subtraction point, N_c is the number

of colors, and $g_s(\mu)$ is the running coupling constant. The Wilson coefficients $c_3(\mu, m)$ and $d(\mu)$ were calculated in leading logarithmic approximation in Ref. [11] and Ref. [1], respectively, and are

$$c_3(\mu, m) = \left(\frac{\alpha_s(\mu)}{\alpha_s(m_1)} \right)^{-\frac{9}{25}}, \quad (3)$$

and

$$\begin{aligned} d(\mu) &= \frac{N_c^2 - 1}{8N_c^2} c_3(m_2, m_1) [1 - c_3^2(\mu, m_2)] \\ &= \frac{N_c^2 - 1}{8N_c^2} \left(\frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^{-\frac{9}{25}} \left[1 - \left(\frac{\alpha_s(\mu)}{\alpha_s(m_2)} \right)^{-\frac{18}{25}} \right]. \end{aligned} \quad (4)$$

In Eq. (2) $V_4(\mu, r)$ is the color magnetic–magnetic correlation function which can be expressed as

$$V_4(\mu, r) \equiv \lim_{T \rightarrow \infty} \int_{-T/2}^{T/2} dz \int_{-T/2}^{T/2} dz' \frac{g_s^2(\mu)}{T} \langle B^i(\mathbf{x}_1, z) B^i(\mathbf{x}_2, z') \rangle / \langle 1 \rangle, \quad (5)$$

where $\langle \dots \rangle$ is defined by

$$\langle \dots \rangle \equiv \int [dA^\mu] Tr \left\{ P \left[\exp \left(ig \oint_{C(r,T)} dz_\mu A^\mu(z) \right) \dots \right] \right\}_{x \in C} \exp(iS_{YM}(A)). \quad (6)$$

Here $C(r, T)$ represents the Wilson loop⁹, P denotes the path ordering, and $r \equiv |\mathbf{x}_1 - \mathbf{x}_2|$.

We emphasize that this is a general result for the hyperfine part of the spin–dependent potential to order $1/m^2$. It absorbs the short distance contributions to the potential into the coefficients $c_3(\mu, m)$ and $d(\mu)$ while the long distance contributions to the potential are contained in the correlation function $V_4(\mu, r)$. Moreover, the result is independent of the factorization scale since the μ –dependence in the coefficients cancels the μ –dependence in the correlation function. The first term in the bracket in Eq. (2) is a nonlocal term while the second term is a local one which is generated by mixing with the first (nonlocal) term under renormalization. We note that if the coefficients are evaluated at tree level; i.e., $c_3(\mu, m) = 1$ and $d(\mu) = 0$, the potential reduces to the Eichten–Feinberg result⁴. And if these coefficients are expanded to order $\alpha_s(\mu)$ and the correlation function is also evaluated only to one–loop, the logarithmic terms in Eq. (2) then reduce to the one–loop spin–spin potential^{5,6}. Therefore, this renormalization–group improved potential, Eq. (2), extends both Eichten’s and Feinberg’s result as well as the one–loop perturbative potential, containing each of these results as special cases.

To first order perturbation theory in ΔH_{ss} the energy shift caused by $\Delta H_{ss}(r)$ is

$$\Delta E = \int d^3\mathbf{r} \Psi_{l,l_z}^*(\mathbf{r}) \Delta H_{ss}(r) \Psi_{l,l_z}(\mathbf{r}) \quad (7)$$

where $\Psi_{l,l_z}(\mathbf{r})$ is the nonrelativistic wavefunction of the bound state with total angular momentum l and z -component l_z . For simplicity we suppress spin and color indices and retain only the space-dependent indices. Separating the radial part, $u(r)$, we write $\Psi_{l,l_z}(\mathbf{r})$ as

$$\Psi_{l,l_z}(\mathbf{r}) = u(r) Y_{l,l_z}(\theta, \phi), \quad (8)$$

where $Y_{l,m}(\theta, \phi)$ are the standard spherical harmonics.

The radial wavefunction was obtained by numerically solving the Schrödinger equation. For comparison, we used three popular potential models. One was the Cornell model¹², the second one was the logarithmic potential¹³, and the third one was the improved QCD-motivated potential¹⁴.

The calculation also required an expression for the running coupling constant $\alpha_s(q)$. The familiar RGE, one-loop result is

$$\alpha_s(q) = \frac{4\pi}{b_0 \ln \frac{q^2}{\Lambda_{\overline{\text{MS}}}^2}}, \quad (9)$$

where $b_0 = 11N_c - 2N_f$ and N_f is the number of quark flavors. It is clear from Eq. (9) that $\alpha_s(q)$ contains a Landau singularity in the nonperturbative region when $q^2 = \Lambda_{\overline{\text{MS}}}^2$ and becomes negative for $q^2 < \Lambda_{\overline{\text{MS}}}^2$. To avoid the resulting numerical ambiguities we first moved this singularity to $q^2 = 0$ and used a modified form of $\alpha_s(q)$ in the actual numerical calculations; namely,

$$\alpha_s(q) = \frac{4\pi}{b_0 \ln \left(\frac{q^2}{\Lambda_{\overline{\text{MS}}}^2} + 1 \right)}. \quad (10)$$

The value of $\Lambda_{\overline{\text{MS}}}$ was taken to be 200 MeV and 250 MeV in the numerical calculations, which is within the experimental range $\Lambda_{\overline{\text{MS}}} = 195 + 65 - 50$ MeV⁸.

To explore the sensitivity of our results to the location of the Landau singularity in $\alpha_s(q)$ we replaced the expression, Eq. (10), by

$$\alpha_s(q) = \frac{4\pi}{b_0 \ln \left(\frac{q^2}{\Lambda_{\overline{\text{MS}}}^2} + \lambda^2 \right)} \quad (11)$$

and varied λ^2 . The results for the S -wave hyperfine splitting were not sensitive to λ^2 and only for large λ^2 did ΔM_p significantly decrease. To fit ΔM_p to the measured value required λ^2 quite large, about 16, clearly out of the perturbative region.

Our numerical results for the ${}^3S_1 - {}^1S_0$ and ${}^3P_J - {}^1P_1$ splittings, where 3P_J denotes the center of gravity of the ${}^3P_{0,1,2}$ states, for the three potentials^{12,13,14} are presented in Tables I, II, III, respectively. For comparison we have also included the results for the $2S$ and $2P$ states.

Table 1. The hyperfine spin splittings in MeV predicted using the Cornell potential ¹²

$\Lambda_{\overline{\text{MS}}} \text{ (MeV)}$	$c\bar{c}$		$b\bar{b}$		$b\bar{c}$	
	200	250	200	250	200	250
$E(1^3S_1) - E(1^1S_0)$	117.1	128.3	97.7	104.3	67.0	71.4
$E(2^3S_1) - E(2^1S_0)$	75.3	82.5	39.6	42.2	37.7	40.3
$E(1^3P_J) - E(1^1P_1)$	-4.8	-7.0	-3.0	-4.1	-4.0	-5.6
$E(2^3P_J) - E(2^1P_1)$	-3.6	-5.2	-2.1	-2.9	-3.0	-4.2

 Table 2. The hyperfine spin splittings in MeV predicted using the Logarithmic potential ¹³

$\Lambda_{\overline{\text{MS}}} \text{ (MeV)}$	$c\bar{c}$		$b\bar{b}$		$b\bar{c}$	
	200	250	200	250	200	250
$E(1^3S_1) - E(1^1S_0)$	106.1	117.1	36.0	38.2	40.6	42.6
$E(2^3S_1) - E(2^1S_0)$	54.2	59.7	18.7	19.8	21.2	22.3
$E(1^3P_J) - E(1^1P_1)$	-5.4	-7.8	-3.4	-4.5	-4.5	-6.4
$E(2^3P_J) - E(2^1P_1)$	-2.6	-3.8	-2.1	-2.8	-2.8	-4.0

 Table 3. The hyperfine spin splittings in MeV predicted using the Improved-QCD motivated potential ¹⁴

$\Lambda_{\overline{\text{MS}}} \text{ (MeV)}$	$c\bar{c}$		$b\bar{b}$		$b\bar{c}$	
	200	250	200	250	200	250
$E(1^3S_1) - E(1^1S_0)$	107.9	119.1	44.6	47.6	43.4	45.7
$E(2^3S_1) - E(2^1S_0)$	68.5	75.6	20.9	22.4	25.2	26.7
$E(1^3P_J) - E(1^1P_1)$	-4.6	-6.7	-2.7	-3.7	-3.7	-5.3
$E(2^3P_J) - E(2^1P_1)$	-3.4	-5.0	-1.8	-2.5	-2.6	-3.8

The main features of these results can be summarized as follows:

- The results are μ -independent, as they must be.
- The calculated energy difference between the J/Ψ and the η_c mesons is quite close to the experimental value ⁸ for all three potentials.
- For each of these three potentials we predict the energy difference between Ψ' and η'_c to lie within the range 55 – 80 MeV.
- For the $b\bar{b}$ system there are significant discrepancies between the Cornell model and the other two models for the S-states. Since the Cornell model, with the parameters fit to the $c\bar{c}$ spectrum, does not predict the $b\bar{b}$ spectrum very well, the results calculated in the other two models are probably better predictions for the energy difference between the $\Upsilon(1S)$ and the η_b (35 – 50 MeV) and between the $\Upsilon(2S)$ and the η'_b (20 MeV).

- The predicted energy difference between B_c^* and B_c meson is in the range 40–70 MeV from all three of these models, which is consistent with previous results ¹⁵.
- The calculated value of $\Delta M_p \equiv E(1^3P_J) - E(1^1P_1)$ for the charmonium $1P$ states is in the range of -4 to -6 MeV, which has the same sign but is several times larger than the experimental value of -0.9 ± 0.2 MeV ⁸. This is not surprising since there are several other contributions to ΔM_p which estimates indicate are comparable in magnitude to the contribution coming from the hyperfine spin–spin interaction, H_{ss} . In fact, it is surprising that the prediction from only the one–loop spin–dependent potential is quite close to the experimental data.

To summarize, the hyperfine spin splittings in the $c\bar{c}$, $b\bar{b}$, and $b\bar{c}$ system were calculated ² using the RGE improved perturbative spin–spin potential ¹. The results for the hyperfine splittings of the S -wave states agree with the $J/\Psi - \eta_c$ measured splitting ⁸ and the prediction for $\Upsilon - \eta_b$ splitting is reasonable. However, the contribution to $\Delta M_p \equiv E(3P_J) - E(1P_1)$ for the charmonium P -wave states is somewhat larger than the experimental data ⁸, although it agrees in sign. That is, after summing up the leading logarithmic terms and including them in the perturbation calculations, the agreement with the data is not as good as the one–loop calculations ^{14,16,17,18}. But there are several additional contributions that are possibly comparable in magnitude. These include the following:

- The contributions of the spin–orbit and tensor potentials in the second order of perturbation theory: These contributions to ΔM_p only cancel to first order in perturbation theory. However, according to the power counting rules ¹⁹, the spin–orbit and tensor potential potentials shift the energies of the P -wave states by an amount of order mv^4 in first order, which indeed cancel in ΔM_p , but they do make a contribution to ΔM_p of order mv^6 in the second order of perturbation theory. This estimate is several MeV for the P -wave charmonium states, and therefore should not be ignored.
- Higher dimensional operators: Unlike the dimension–six operators, these give non-zero contributions to ΔM_p even at tree level. Compared to the one–loop contribution, these are suppressed by v^2 but enhanced by α_s^{-1} and $v^2/\alpha_s \sim 1$ in charmonium.
- The color-octet S -wave component in P -wave quarkonia states ¹⁹: This component of the wavefunction receives a tree–level contribution from the local term $\delta^3(\mathbf{r})$ in the spin–spin potential. This contribution too could be of order $v^2/\alpha_s \sim 1$ compared to what has been calculated.

- Next-to-leading order contributions from the two-loop potential: These are suppressed by order α_s , but since α_s is not a very small quantity in charmonium, one cannot dismiss the possibility that this contribution could be significant.

Before comparing with the experimental value of ΔM_p in charmonium, which is only about 1 MeV, all the above contributions should be included since they are possibly comparable in magnitude. In the $b\bar{b}$ case these effects are less important and one can expect the perturbative calculations the $b\bar{b}$ system to be more reliable, although ΔM_p is smaller, also.

This research was done in collaboration with Yu-Qi Chen and Yu-Ping Kuang and is presented in greater detail in references [1] and [2]. It was supported in part by the U.S. National Science Foundation under Grant No. PHYS89-04035, the National Science Foundation of China, the Fundamental Research Foundation of Tsinghua University, and the U.S. Department of Energy, Division of High Energy Physics, under Grant No. DE-FG02-91-ER40684.

1. Y.-Q. Chen, Y.-P. Kuang, and R.J. Oakes, *Phys. Rev.* **D52**, 264 (1995).
2. Y.-Q. Chen and R.J. Oakes, to appear in *Phys. Rev. D*.
3. H.D. Politzer and M.B. Wise, *Phys. Lett.* **206B**, 681 (1988); **208B**, 504 (1988); N. Isgur and M.B. Wise, *Phys. Lett.* **232B**, 113 (1989); *Phys. Lett.* **237B**, 527 (1990); H. Georgi, *Phys. Lett.* **264B**, 447 (1991); E. Eichten and B. Hill, *Phys. Lett.* **233B**, 427 (1990); *ibid* **234B** 511 (1990); B. Grinstein, *Nucl. Phys.* **B339**, 447 (1990); T. Mannel, W. Robert, and Z. Ryzak, *Nucl. Phys.* **B368**, 204 (1992); A.F. Falk, H. Georgi, B. Grinstein, and M.B. Wise, *Nucl. Phys.* **B343**, 1 (1990); H. Georgi, B. Grinstein, and M.B. Wise, *Phys. Lett.* **252B**, 456 (1990); M.E. Luke, *Phys. Lett.* **252B**, 447 (1990); A.F. Falk, M. Neubert, and M.E. Luke, *Nucl. Phys.* **B388**, 363 (1992); M.E. Luke, A. V. Manohar, *Phys. Lett.* **B286**, 348 (1992). Yu-Qi Chen, *Phys. Lett.* **B317**, 421 (1993).
4. E. Eichten and F. Feinberg, *Phys. Rev. Lett* **43**, 1205 (1979); *Phys. Rev.* **D23**, 2724 (1982).
5. W. Buchmüller, Y. J. Ng, and, S. -H. H. Tye, *Phys. Rev.* **D24**, 3003 (1981); S.N. Gupta, and S.F. Radford, *Phys. Rev.* **D24**, 2309 (1981). S.N. Gupta, S.F. Radford, and W.W. Repko, *Phys. Rev.* **D26**, 3305 (1982); S.N. Gupta, and S.F. Radford, *Phys. Rev.* **D25**, 3430 (1982).
6. J. Pantaleone, S. -H. H. Tye, and Y. J. Ng, *Phys. Rev.* **D33**, 777 (1986).
7. D. Gromes, *Z. Phys.* **C26**, 401 (1984); W. Buchmüller, *Phys. Lett.* **B112**, 479 (1982); Y.-Q. Chen and Y.-P. Kuang, *Z. Phys.* **C67**, 627 (1995).
8. L. Monlanet, *et al.* Particle Data Group, *Phys. Rev.* **D50**, 1173 (1994).
9. K. Wilson, *Phys. Rev.* **D10**, 2445 (1974).
10. L.S. Brown and W.I. Weisberg, *Phys. Rev.* **D20**, 3239 (1979);

11. G.L. Lepage and B.A. Thacker, in *field theory on the Lattice*, proceedings of the International Symposium, Seillac France, 1987, edited by A. Billoire *et al.* *Nucl. Phys. B* (Proc. Suppl.) **4**, 199 (1988); A.F. Falk, B. Grinstein, and M.E. Luke, *Nucl. Phys. B***357**, 185 (1991).
12. E. Eichten, K. Gottfried, T. Kinoshita, K.D. Lane, and T.-M. Yan, *Phys. Rev. D***17**, 3090 (1978); *ibid*, **21**, 313(E) (1980); *ibid*, **21**, 203 (1980).
13. C. Quigg and J.L. Rosner, *Phys. Lett.* **71B**, 153 (1977).
14. Y.-Q. Chen and Y.-P. Kuang, *Phys. Rev. D***46** 1165 (1992).
15. E.J. Eichten and C. Quigg, *Phys. Rev. D***49**, 5845 (1994).
16. D.B. Lichtenberg and R. Potting, *Phys. Rev. D***47**, 3013 (1993).
17. F. Halzen, C. Olson, M.G. Olsson, and M. L. Strong, *Phys. Lett.* **283B**, 379 (1992); *Phys. Rev. D***47**, 3013 (1993).
18. H. Grotch, K. J. Sebastian, and X. Zhang, *Phys. Rev. D***49**, 1643 (1994).
19. G.T. Bodwin, E. Braaten, T.C. Yuan, G.P. Lepage, *Phys. Rev. D***46**, R3707 (1992); G.T. Bodwin, E. Braaten, G.P. Lepage, *Phys. Rev. D***51**, 1125 (1995)