

BARI-TH/95-219  
 UTS-DFT-95-13  
 December 1995

## On the $b$ Quark Kinetic Energy in the $\Lambda_b$

P. Colangelo <sup>a</sup>, C.A. Dominguez <sup>b</sup>, G. Nardulli <sup>a,c</sup> and N. Paver <sup>d</sup>

<sup>a</sup> *Istituto Nazionale di Fisica Nucleare, Sezione di Bari, Italy*

<sup>b</sup> *Physics Department, University of Capetown, South Africa*

<sup>c</sup> *Dipartimento di Fisica, Università di Bari, Italy*

<sup>d</sup> *Dipartimento di Fisica Teorica, Università di Trieste, and  
 Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, Italy*

### Abstract

We consider the problem of determining the matrix element of the heavy quark kinetic energy operator  $\mathcal{K}$  between  $\Lambda_b$  baryonic states by three point function QCD sum rules in the Heavy Quark Effective Theory. We discuss the main features and the uncertainties associated to such an approach. Numerically, we find the result:  $\langle \mathcal{K} \rangle_{\Lambda_b} = -(0.6 \pm 0.1) \text{ GeV}^2$ . Within the uncertainty of the method, such result indicates a value of the heavy quark kinetic energy in the baryon similar in size to that in the meson.

PACS:12.39.Hg, 12.38.Lg

Typeset using REVTeX

## I. INTRODUCTION

Important progresses in the theoretical description of hadrons containing one heavy quark have been allowed by the development of the Heavy Quark Effective Theory (HQET) [1]. Based on the spin-flavor symmetry of QCD, valid in the limit of infinite heavy quark mass  $m_Q$ , this framework provides a systematic expansion of heavy hadron spectra and transition amplitudes in terms of the leading contribution (for  $m_Q \rightarrow \infty$ ), plus corrections decreasing as powers of  $1/m_Q$ . Being derived from ‘first principles’, i.e. from general properties of QCD, such an approach has the advantage of being model-independent to a large extent, and predictive at the same time. For this reason, it has been extensively applied to a variety of aspects concerning the dynamics of mesons and baryons made of both heavy and light quarks.

Due to the flavor-spin symmetry connecting matrix elements for different heavy flavors and angular momenta, the leading order description is simple and intuitive, and involves a few nonperturbative quantities in terms of which a large number of processes can be described. This leading order seems to be consistent with the main features of heavy hadron physics as they are presently observed. However, although being suppressed by the large heavy quark mass, the next-to-leading order corrections proportional to  $1/m_Q$  are also expected to play a significant role. In the framework of HQET, they are limited in number, and can be systematically identified and related to nonperturbative parameters having a clear physical meaning. A noticeable example can be found in the effective QCD heavy quark Lagrangian which, up to order  $1/m_Q$ , can be written as

$$\mathcal{L} = \bar{h}_v i(v \cdot D)h_v + \frac{1}{2m_Q}\mathcal{K} + \frac{1}{2m_Q}\mathcal{S} + O(\frac{1}{m_Q^2}). \quad (1.1)$$

Here  $h_v$  is the effective field for a heavy quark with velocity  $v$ , related to the Dirac field  $Q$  by

$$h_v(x) = e^{im_Q v \cdot x} \frac{1 + \not{v}}{2} Q(x), \quad (1.2)$$

$\mathcal{K}$  is the nonrelativistic kinetic energy operator defined by

$$\mathcal{K} = \bar{h}_v (iD^\perp)^2 h_v \quad (1.3)$$

where  $(D^\perp)^2 = D_\mu D^\mu - (v \cdot D)^2$ , with  $D^\mu$  the QCD covariant derivative, and  $\mathcal{S}$  is the Pauli term responsible of the spin splitting among the states belonging to the same heavy hadron multiplet in the  $m_Q \rightarrow \infty$  limit:

$$\mathcal{S} = \frac{1}{2} \left( \frac{\alpha_s(m_Q)}{\alpha_s(\mu)} \right)^{3/\beta_0} \bar{h}_v \sigma^{\mu\nu} g G_{\mu\nu} h_v. \quad (1.4)$$

Appropriately normalized matrix elements of the operators (1.3) and (1.4) account for the next-to-leading corrections to the heavy hadron mass spectra:

$$m_H = m_Q + \Delta - \frac{1}{2m_Q} \langle \mathcal{K} \rangle_H - \frac{c_H}{2m_Q} \langle \mathcal{S} \rangle_H, \quad (1.5)$$

where  $\Delta$  represents the binding energy of light quarks in the static approximation and  $c_H$  is a numerical coefficient. At the same time, they are involved as ‘preasymptotic’ corrections in the  $1/m_Q$  expansion of inclusive heavy hadron decay rates based on parton model ideas and on the operator product expansion [2,3]. For example, the inclusive  $H \rightarrow f$  weak decay rate can be written as:

$$\Gamma(H \rightarrow f) = \frac{G_F^2 m_Q^5}{192\pi^3} |V_{CKM}|^2 \left[ c_3^f \left( 1 + \frac{\langle \mathcal{K} \rangle_H}{2m_Q^2} \right) + c_5^f \frac{\langle \mathcal{S} \rangle_H}{2m_Q^2} + O\left(\frac{1}{m_Q^3}\right) \right], \quad (1.6)$$

where  $c_3^f$  and  $c_5^f$  are calculable short-distance coefficients that depend on the process under consideration.

The matrix elements of  $\mathcal{K}$  and  $\mathcal{S}$  are nonperturbative parameters that should be either determined phenomenologically from experimental data or estimated using a nonperturbative theoretical approach. In the case of heavy mesons ( $P_Q$ ), information on both  $\langle \mathcal{K} \rangle_{P_Q}$  and  $\langle \mathcal{S} \rangle_{P_Q}$  is available. Indeed, the chromomagnetic operator matrix element can be determined from the vector-pseudoscalar meson mass difference, indicating

$$\langle \mathcal{S} \rangle_{P_Q} = \frac{3}{4} (m_{V_Q}^2 - m_{P_Q}^2) \simeq 0.37 \text{ GeV}^2, \quad (1.7)$$

and it has also been theoretically estimated by QCD sum rules giving a result in agreement with this value [4]. Concerning the kinetic energy operator matrix element, general arguments imply the inequality [5]  $|\langle \mathcal{K} \rangle_{P_Q}| \geq \langle \mathcal{S} \rangle_{P_Q}$  giving  $|\langle \mathcal{K} \rangle_{P_Q}| \geq 0.37 \text{ GeV}^2$  using (1.7), and the most recent theoretical estimate using QCD sum rules indicates [4]<sup>1</sup>

$$\langle \mathcal{K} \rangle_{P_Q} = -(0.60 \pm 0.10) \text{ GeV}^2. \quad (1.8)$$

Finally, the first calculation with lattice QCD gives the inequality  $|\langle \mathcal{K} \rangle_{P_Q}| < 1 \text{ GeV}^2$ , in agreement with the above numbers [7].

On the other hand, no analogous information is available for the baryonic matrix elements of  $\mathcal{S}$  and  $\mathcal{K}$ . In fact, the heavy baryon mass splittings relevant to the determination of the chromomagnetic expectation value are not measured yet; moreover, general lower bounds on the kinetic energy have not been worked out in this case. To leading order in  $1/m_Q$ , the heavy quark kinetic energy inside the baryon can be connected to that in the meson and to the heavy quark masses by the relation (which assumes that the charm mass  $m_c$  is heavy enough for the expansion up to  $1/m_c$  to be meaningful) [8]:

$$\langle \mathcal{K} \rangle_{P_Q} - \langle \mathcal{K} \rangle_{\Lambda_Q} \simeq \frac{m_b m_c}{2(m_b - m_c)} [(m_B + 3m_{B^*} - 4m_{\Lambda_b}) - (m_D + 3m_{D^*} - 4m_{\Lambda_c})]. \quad (1.9)$$

Using present data, Eq. (1.9) would give  $\langle \mathcal{K} \rangle_{P_Q} - \langle \mathcal{K} \rangle_{\Lambda_Q} \simeq 0.07 \pm 0.20 \text{ GeV}^2$ , where the error mainly comes from the error on the measured  $m_{\Lambda_b}$ , that is not small enough:  $m_{\Lambda_b} = 5641 \pm 50 \text{ MeV}$  [9]. Arguments have been given in favour of the near equality of the kinetic energy operator matrix elements between meson and baryon states [10], which could

---

<sup>1</sup>For earlier estimates see [6].

also be expected on intuitive grounds, but some quantitative estimates are needed. Here, we would like to consider in this regard the simplest case, represented by the  $\Lambda_b$  baryon, where the chromomagnetic term vanishes in the infinite heavy quark mass limit  $\langle \mathcal{S} \rangle_{\Lambda_b} = 0$  as the light diquark system carries no spin (the same is true for the  $\Xi_Q$ ), and attempt an assessment of the heavy quark kinetic energy. Specifically, we discuss the possibility of such an estimate in the framework of QCD sum rules, where the calculation has also been done for the meson, so that it is of interest to compare the two cases.

## II. HEAVY BARYON CURRENT AND TWO-POINT FUNCTION

A basic element in the application of QCD sum rules to baryons is the definition of a suitable baryon interpolating effective field in terms of quark fields, and the corresponding vacuum-to-baryon matrix element. For the case of heavy baryons this problem has been discussed previously in Refs. [11]- [14], and we outline here this analysis for the sake of completeness.

The general expression of the interpolating field for a baryon state with one heavy quark and two light quarks can be written as

$$J_D^{(\ell)} = \epsilon^{\alpha\beta\gamma} (q_1^{T\alpha} C T^{(\ell)} \tau q_2^\beta) \tilde{\Gamma} (h_v^\gamma)_D \quad (2.1)$$

where  $T$  means transpose,  $\alpha$ ,  $\beta$  and  $\gamma$  are color indices,  $D$  is a Dirac index,  $C$  is the charge conjugation matrix,  $q_1$  and  $q_2$  are light quark field operators, and  $\tau$  is a flavor matrix. For the case of the spin  $1/2$   $\Lambda_b$  of interest here, the matrices  $\tau$  and  $\tilde{\Gamma}$  are given by:

$$\tau_{ij} = \frac{1}{\sqrt{2}} (\delta_{1i} \delta_{2j} - \delta_{2i} \delta_{1j}), \quad \tilde{\Gamma} = 1, \quad (2.2)$$

whereas two choices are possible for the matrix  $\Gamma^{(\ell)}$ :

$$\Gamma^{(1)} = \gamma_5, \quad \Gamma^{(2)} = \gamma_5 \gamma_0. \quad (2.3)$$

Such choices correspond to spin-parity  $J^P = 0^+$  for the light quark pair, as implied by the predicted spectroscopy of baryons containing one heavy quark in the infinite mass limit [15,16].

Consequently, the general  $\Lambda_b$  interpolating field is the linear combination

$$J = J^{(1)} + b J^{(2)}, \quad (2.4)$$

depending on the, a priori unspecified, parameter  $b$ .

Together with the baryonic interpolating currents, we need for our estimates the values of the vacuum-to-baryon couplings ( $\ell = 1, 2$ ):

$$\langle 0 | J_D^{(\ell)} | \Lambda_b(v) \rangle = f_{\Lambda_b}^{(\ell)} (h_v)_D. \quad (2.5)$$

In the framework of QCD sum rules these constants can be obtained from the two-point correlators:

$$\Pi_{CD}^{(\ell, \ell')}(k) = i \int dx \langle 0 | T(J_C^{(\ell)}(0) \bar{J}_D^{(\ell')}(x)) | 0 \rangle e^{ikx} \quad (2.6)$$

where, according to the familiar rules of the heavy quark theory,  $k$  is the residual momentum, corresponding to the decomposition  $p^\mu = m_b v^\mu + k^\mu$  of the heavy baryon momentum; we parametrize it as

$$k^\mu = \omega v^\mu. \quad (2.7)$$

Taking Eqs. (1.2) and (2.7) into account, we can write the LHS of Eq. (2.6) as

$$\Pi_{CD}^{(\ell, \ell')}(k) = (1 + \not{v})_{CD} \Pi^{(\ell, \ell')}(\omega). \quad (2.8)$$

The scalar functions  $\Pi^{(\ell, \ell')}$  satisfy dispersion relations of the form:

$$\Pi^{(\ell, \ell')}(\omega) = \frac{1}{\pi} \int d\Omega \frac{\rho^{(\ell, \ell')}(\Omega)}{\Omega - \omega} + \text{subtractions}, \quad (2.9)$$

where  $\rho$  represents the spectral function:  $\rho^{(\ell, \ell')}(\Omega) = \text{Im} \Pi^{(\ell, \ell')}(\Omega)$ . The explicit form of the polynomial representing the subtractions is not important because in the sequel we shall apply the borelized version of the sum rule, and the Borel transform of any polynomial vanishes.

Following the QCD sum rules method [17],  $\rho^{(\ell, \ell')}(\Omega)$  can be computed in two ways: either by saturating the dispersion relation (2.9) by physical hadronic states or by means of the operator product expansion (OPE), which is allowed insofar  $k^2$  in (2.6) is large and negative.

Performing the OPE on  $\Pi^{(\ell, \ell')}$  and taking into account the operators of dimension  $d = 0$  (the perturbative term),  $d = 3$  and  $d = 5$ , by a straightforward calculation in the heavy quark theory one finds the spectral functions:

$$\rho^{(\ell, \ell)}(\Omega) = \rho^{(0)}(\Omega) = \frac{\Omega^5}{20\pi^3} \quad (2.10)$$

for  $\ell = \ell'$  ('diagonal' sum rules), and

$$\rho^{(1,2)}(\Omega) + \rho^{(2,1)}(\Omega) = \rho^{(3)}(\Omega) + \rho^{(5)}(\Omega) = -\frac{2\langle\bar{q}q\rangle\Omega^2}{\pi} - \frac{2m_0^2\langle\bar{q}q\rangle}{\pi} \left(-\frac{1}{8} + \frac{1}{16}\right) \quad (2.11)$$

for  $\ell \neq \ell'$  ('nondiagonal' sum rules), as obtained previously [12–14]. In Eqs. (2.10) and (2.11)  $\rho^{(0,3,5)}$  denote, respectively, the leading perturbative term in the short distance expansion and the  $d = 3$  and  $d = 5$  contributions. Moreover,  $\langle\bar{q}q\rangle$  is the quark condensate ( $\langle\bar{q}q\rangle = -(230 \text{ MeV})^3$  at the scale of  $1 \text{ GeV}$ ) and  $m_0$  is connected to the quark-gluon condensate by the relation  $\langle\bar{q}g_s\sigma \cdot Gq\rangle = m_0^2\langle\bar{q}q\rangle$  (and  $m_0^2 = 0.8 \text{ GeV}^2$  [17]). The separation of the  $d = 5$  contribution in two contributions reflects the existence of two independent sources originating such a term, namely the nonlocal quark condensate [18]

$$\langle\bar{q}(x)q(0)\rangle = \Phi(x^2)\langle\bar{q}q\rangle \simeq \left(1 + \frac{x^2 m_0^2}{16}\right) \langle\bar{q}q\rangle \quad (2.12)$$

and the diagram where the gluon non perturbative field is emitted from the perturbative light quark line (in the adopted Fock-Schwinger gauge, no such gluon can be emitted from the heavy quark leg in the infinite heavy quark mass limit).

Actually, in previous calculations also the  $d = 4$  gluon condensate  $\langle \alpha_s G^2 \rangle$  and the higher dimension operators with  $d > 5$  have been included. The possibility of potentially non-negligible higher order terms is briefly discussed at the end of this Section. On the other hand, the gluon condensate contribution would be rather easy to include both in Eq. (2.10) and in the analogous three-point functions considered in the next Section to evaluate the kinetic energy. In principle, its presence accounts for the low momentum part of the gluon exchange diagram of order  $\alpha_s$ . In the baryon case the high momentum component, i.e. the  $\alpha_s$  correction to the perturbative diagram, requires a three-loop calculation, which has not been attempted yet. Consequently, we do not include the gluon condensate as it would only partially represent the gluon exchange, and we limit to the analysis of both two-point and three-point correlators with the operators having  $d = 0, 3$  and  $5$ .

Turning to the representation of spectral functions in (2.9) in terms of physical hadronic states, we write

$$\rho_{had}^{(\ell,\ell')}(\Omega) = \rho_{res}^{(\ell,\ell')}(\Omega) + \rho_{cont}^{(\ell,\ell')}(\Omega), \quad (2.13)$$

where  $\rho_{res}^{(\ell,\ell')}(\Omega)$  contains the contribution of the low-lying baryon state, in this case the  $\Lambda_b$ , and  $\rho_{cont}^{(\ell,\ell')}$  represents the contribution of higher mass states. Assuming, as usual, duality between partons and resonances,  $\rho_{cont}^{(\ell,\ell')}(\Omega)$  can be taken as equal to the operator product expansion  $\rho^{(\ell,\ell')}(\Omega)$  of Eqs. (2.10) and (2.11) for  $\Omega > \omega_c$ , with  $\omega_c$  a continuum threshold, and vanishing elsewhere. As for  $\rho_{res}^{(\ell,\ell')}(\Omega)$ , they are expressed as:

$$\rho_{res}^{(\ell,\ell)}(\Omega) = \frac{\pi [f_{\Lambda_b}^{(\ell)}]^2}{2} \delta(\Omega - \Delta) \quad (2.14)$$

$$\rho_{res}^{(1,2)}(\Omega) + \rho_{res}^{(2,1)}(\Omega) = \frac{\pi f_{\Lambda_b}^{(1)} f_{\Lambda_b}^{(2)}}{2} \delta(\Omega - \Delta) \quad (2.15)$$

where  $f_{\Lambda_b}^{(\ell)}$  have been defined in Eq. (2.5) and  $\Delta = m_{\Lambda_b} - m_b$  in the limit  $m_b \rightarrow \infty$  (see Eq. (1.5)). Equating the two representations of the two-point correlators, and taking the Borel transform in the variable  $\omega$  of both sides, one obtains the set of sum rules for the vacuum-to-baryon couplings of the interpolating fields:

$$[f_{\Lambda_b}^{(1)}]^2 = [f_{\Lambda_b}^{(2)}]^2 = \frac{2}{\pi} \int d\Omega e^{-\frac{\Omega-\Delta}{E}} \rho^{(0)}(\Omega) \quad (2.16)$$

$$f_{\Lambda_b}^{(1)} f_{\Lambda_b}^{(2)} = \frac{1}{\pi} \int d\Omega e^{-\frac{\Omega-\Delta}{E}} [\rho^{(3)}(\Omega) + \rho^{(5)}(\Omega)]. \quad (2.17)$$

One can notice that, due to the structure of Eq. (2.10), at the chosen order of approximation in the operator product expansion  $|f_{\Lambda_b}^{(1)}| \simeq |f_{\Lambda_b}^{(2)}| = f_{\Lambda_b}$  from the ‘diagonal’ sum rules.<sup>2</sup> This

---

<sup>2</sup>As it will be seen in the next Section, this does no longer occur in the case of the three-point function, in the sense that somehow different results for the kinetic energy are obtained from the two diagonal sum rules.

result should be confirmed by the non-diagonal sum rule, which can also be used to fix the relative sign between the two constants.

In order to extract useful information from the sum rules, we have to fix the parameters  $\Delta$ ,  $\omega_c$  and the allowed range for the Borel parameter  $E$  (the so-called ‘duality’ region). Indeed, in order to obtain a prediction for  $\Delta$ , we differentiate (2.16) ( or (2.17) ) in the variable  $1/E$  and take the ratio with (2.16) (or (2.17) ) itself, thus obtaining a new sum rule where only  $\Delta$  (and not  $f_{\Lambda_b}$ ) appears.

The results for  $f_{\Lambda_b}$  and  $\Delta$  as a function of the Borel variable  $E$ , and for  $\omega_c$  in the range  $1.1 - 1.3 \text{ GeV}$ , are shown in Fig. 1. In particular, in Fig. 1a we depict the result for the diagonal correlators (2.16) both concerning the coupling  $f_{\Lambda_b}$  and  $\Delta$ , in Fig. 1b the outcome of the nondiagonal correlator (2.17). In Fig. 1c we display the result from the sum rule for the correlator of the current  $J$  in Eq.(2.4):

$$\frac{1}{2}f_{\Lambda_b}^2(1+b)^2e^{-\Delta/E} = \int_0^{\omega_c} d\omega e^{-\omega/E} \left[ \frac{1+b^2}{20\pi^4}\omega^5 - \frac{2b\langle\bar{q}q\rangle}{\pi^2}(\omega^2 - \frac{m_0^2}{16}) \right] \quad (2.18)$$

with the choice  $b = 1$ , that we shall justify in the next Section.

As one can see, stability is always reached for values of  $E$  from  $0.3 \text{ GeV}$  on, which at first sight would be a welcomed property. However, the familiar criterion that the continuum should not be too large with respect to the ground state baryon contribution, favoring low values of  $E$ , rather strongly reduces the ‘working’ region to a range in the neighborhood of  $E = 0.4 - 0.5 \text{ GeV}$ . In the next Section this effect of the continuum will be found to be even more significant in the kinetic energy calculation using three-point sum rules. Clearly, it originates from the high power dependence of the spectral functions, that is expected on dimensional grounds. In this regard, considering the structure of the operator product expansion in Eqs. (2.10) and (2.11), it should be noticed that the suppression of dimensional condensates is provided by smaller powers of the integration variable  $\Omega$ , compared to the leading perturbative term, rather than by large mass denominators characterizing the applications of the QCD sum rule method with finite heavy quark mass.

Coming to the numerical results, it is worth observing that the diagonal and non-diagonal correlators, together with the correlator of the current  $J$  in (2.4), provide us with the same mass parameter  $\Delta_{\Lambda_b}$ :

$$\Delta_{\Lambda_b} = 0.9 \pm 0.1 \text{ GeV} . \quad (2.19)$$

The uncertainty on  $\Delta_{\Lambda_b}$  in (2.19) comes from the variation of the continuum threshold and from changing the Borel parameter  $E$  in the range  $E = 0.3 - 0.4 \text{ GeV}$  in Fig.1a and c, and  $E = 0.3 - 0.5 \text{ GeV}$  in Fig.1b. Using the result of the analysis for the mesonic system  $\Delta_B \simeq 0.5 \text{ GeV}$  [4] we get for  $\Lambda_b$  the mass  $M_{\Lambda_b} = M_B + \Delta_{\Lambda_b} - \Delta_B \simeq 5.7 \text{ GeV}$ , to be compared with the experimental measurement quoted above. The difference of  $200 \text{ MeV}$  between the mass of the  $\Lambda_b$  and the continuum threshold is of reasonable size as it nearly corresponds to the mass of a pair of pions.

As for the coupling  $f_{\Lambda_b}$ , the diagonal and non-diagonal sum rules in (2.16) and (2.17) give rather different results:

$$f_{\Lambda_b} = (2.0 \pm 0.5) \times 10^{-2} \text{ GeV}^3 \quad (2.20)$$

$$f_{\Lambda_b} = (3.5 \pm 0.5) \times 10^{-2} \text{ GeV}^3 \quad (2.21)$$

respectively. The different results can be put in better agreement using a smaller value for the quark condensate, as advocated in [12]. However, from the correlator of the current  $J$ , with  $b = 1$ , we obtain:

$$f_{\Lambda_b} = (2.9 \pm 0.5) \times 10^{-2} \text{ GeV}^3, \quad (2.22)$$

that is the result we shall use in the next Section.

As far as higher dimensional contributions to (2.20)-(2.22) are concerned, it was observed in [12,14] that the most important one should be the four-quark  $d = 6$  operator contributing to the diagonal sum rule (2.10) and to the sum rule (2.18). The evaluation of the relevant diagram is straightforward and leads to a term proportional to  $\langle \bar{q}(x)q(0)\bar{q}(x)q(0) \rangle$ . The numerical analysis can be done by making the assumption of factorization of the nonlocal four-quark condensate and, using this hypothesis, we would obtain a correction to (2.20) and (2.22) within the quoted uncertainty. However, the accuracy of this assumption is difficult to assess due to the fact that the numerical result would strongly depend on the explicit model for the nonlocal condensate and, as a matter of fact, the inclusion of this term has been questioned, e.g. in ref. [14]. For this reason, following [14], we prefer to omit this term, with the understanding that this neglect will be reflected in an additional uncertainty in the final result.

### III. THE THREE-POINT FUNCTION AND $\langle \mathcal{K} \rangle_{\Lambda_B}$

To apply the formalism of QCD sum rules to the kinetic energy operator  $\mathcal{K}$ , we consider the three-point correlators:

$$\Sigma_{CD}^{(\ell,\ell')}(k, k') = (1 + \not{x})_{CD} \Sigma^{(\ell,\ell')}(\omega, \omega') = i^2 \int dx dy \langle 0 | T(J_C^{(\ell)}(y) \mathcal{K}(0) \bar{J}_D^{(\ell')}(x) | 0 \rangle e^{-iky + ik'x}, \quad (3.1)$$

where  $(l, l') = 1, 2$  have the same meaning as in Eq. (2.6), and  $k, k'$  are the residual momenta, corresponding to the decomposition of the heavy baryon momentum  $p^\mu = m_b v^\mu + k^\mu$ ,  $p'^\mu = m_b v^\mu + k'^\mu$ . Analogously to Eq. (2.7), we have taken:

$$k^\mu = \omega v^\mu, \quad k'^\mu = \omega' v^\mu. \quad (3.2)$$

The scalar function  $\Sigma^{(\ell,\ell')}$  satisfies a double dispersion relation in the variables  $\omega, \omega'$ :

$$\Sigma^{(\ell,\ell')}(\omega, \omega') = \int d\Omega d\Omega' \frac{\sigma^{(\ell,\ell')}(\Omega, \Omega')}{(\Omega - \omega)(\Omega' - \omega')} + \text{subtractions} \quad (3.3)$$

where, as for the two-point function, the explicit form of the subtraction term is not important since it will be eliminated by the double Borel transform of the sum rule.

For both  $k^2$  and  $k'^2$  large and negative, so that short distances  $x$  and  $y$  should dominate, we perform the operator product expansion on the RHS of Eq. (3.1) taking into account only the operators of dimension  $d = 0$  (the perturbative term),  $d = 3$  and  $d = 5$ , analogously to Sec. 2, and neglecting the  $d = 4$  gluon condensate with the same motivation given there.



In this approximation we obtain for the various spectral functions the following expressions, applying the Cutkosky rules in momentum space:

$$\sigma^{(1,1)}(\Omega, \Omega') = -\frac{3\Omega^7}{140\pi^4}\delta(\Omega - \Omega') \quad (3.4)$$

$$\sigma^{(2,2)}(\Omega, \Omega') = -\frac{5\Omega^7}{140\pi^4}\delta(\Omega - \Omega') \quad (3.5)$$

$$\sigma^{(1,2)}(\Omega) + \sigma^{(2,1)}(\Omega) = \left[ \frac{2\langle \bar{q}q \rangle}{\pi^2}\Omega^4 + \frac{2m_0^2\langle \bar{q}q \rangle}{\pi^2}\Omega^2 \left( -\frac{3}{8} + \frac{3}{16} \right) \right] \delta(\Omega - \Omega'). \quad (3.6)$$

Thus,  $\sigma^{(1,1)}$  and  $\sigma^{(2,2)}$  only contain the leading perturbative term in the short distance expansion, and  $d = 3$  and  $d = 5$  condensates do not contribute to the operator product expansion of these ‘diagonal’ correlators. Also the  $d = 6$  four-quark operator does not contribute in this case. On the other hand, the ‘nondiagonal’ spectral function  $\sigma^{(1,2)}(\Omega) + \sigma^{(2,1)}(\Omega)$  is determined by the  $d = 3$   $\langle \bar{q}q \rangle$  vacuum condensate and by the  $d = 5$  quark-gluon condensate. Similar to Eq. (2.11), for the latter condensate the two separate contributions in (3.6) arise from the non local contribution of Eq. (2.12) and from the diagram where the non perturbative low-frequency gluon is emitted from the perturbatively propagating light quark.

The hadronic side of the sum rule is written as

$$\sigma_{had}^{(\ell,\ell')}(\Omega, \Omega') = \sigma_{res}^{(\ell,\ell')}(\Omega, \Omega') + \sigma_{cont}^{(\ell,\ell')}(\Omega, \Omega') , \quad (3.7)$$

where  $\sigma_{res}^{(\ell,\ell')}(\Omega, \Omega')$  contains the contribution of the  $\Lambda_b$  ground state and  $\sigma_{cont}^{(\ell,\ell')}$  contains the contribution of higher mass states. By applying the duality principle,  $\sigma_{cont}^{(\ell,\ell')}$  is assumed equal to the operator product expansion  $\sigma^{(\ell,\ell')}(\Omega, \Omega')$  of Eqs. (3.4)-(3.6) in the domain  $\Omega > \omega_c$  and  $\Omega' > \omega_c$ , and vanishing outside this domain. Owing to the  $\delta$ -function behavior, the details of the integration domain far from the diagonal  $\Omega = \Omega'$  should not be important.

The pole contribution is as follows:

$$\sigma_{res}^{(\ell,\ell)}(\Omega, \Omega') = \frac{[f_{\Lambda_b}^{(\ell)}]^2 \langle \mathcal{K} \rangle_{\Lambda_b}}{2(\Omega - \Delta)(\Omega' - \Delta)}, \quad (3.8)$$

$$\sigma_{res}^{(1,2)}(\Omega, \Omega') + \sigma_{res}^{(2,1)}(\Omega, \Omega') = \frac{f_{\Lambda_b}^{(1)} f_{\Lambda_b}^{(2)} \langle \mathcal{K} \rangle_{\Lambda_b}}{2(\Omega - \Delta)(\Omega' - \Delta)}, \quad (3.9)$$

where the matrix element  $\langle \mathcal{K} \rangle_{\Lambda_b}$  is defined as:

$$\langle \mathcal{K} \rangle_{\Lambda_b} = \langle \Lambda_b(v) | \mathcal{K} | \Lambda_b(v) \rangle = \langle \Lambda_b(v) | \bar{h}_v(iD^\perp)^2 h_v | \Lambda_b(v) \rangle . \quad (3.10)$$

Sum rules for  $\langle \mathcal{K} \rangle_{\Lambda_b}$  are obtained by taking double Borel transforms in the variables  $\Omega, \Omega'$  of both  $\Sigma_{OPE}$  and  $\Sigma_{res}$ , giving :

$$\mathcal{B}(E) \frac{1}{\omega - \Omega} = \frac{1}{E} e^{-\omega/E}, \quad \mathcal{B}(E) \frac{1}{\Delta - \Omega} = \frac{1}{E} e^{-\Delta/E}, \quad (3.11)$$

and similar for  $\Omega'$ , and by assuming that one can equate them for some range of the Borel parameters  $E_1, E_2$ . The symmetry of the spectral functions in  $\Omega, \Omega'$  suggests the choice  $E_1 = E_2$  and, moreover, we take  $E_1 = E_2 = 2E$  with  $E$  the Borel parameter of the sum rule for the two-point correlator, by analogy with the calculation of the Isgur-Wise form factor, both for baryons [13] and mesons [19], where the normalization at zero recoil requires such a relation. Finally, we assume the same continuum threshold  $\omega_c$  as in the two-point function sum rule, namely  $\omega_c = 1.1, 1.2$  and  $1.3 \text{ GeV}^2$ .

In this way, eliminating the couplings  $f_{\Lambda_b}^{(\ell)}$  by means of the two-point sum rules in Eqs. (2.16) and (2.17), we obtain for the particular choices of the baryonic interpolating field the following sum rules for the matrix element  $\langle \mathcal{K} \rangle_{\Lambda_b}$ :

$$\langle \mathcal{K} \rangle_{\Lambda_b} = -\frac{3}{7} \frac{I_7}{I_5} \quad (\ell = \ell' = 1) \quad (3.12)$$

$$\langle \mathcal{K} \rangle_{\Lambda_b} = -\frac{5}{7} \frac{I_7}{I_5} \quad (\ell = \ell' = 2) \quad (3.13)$$

$$\langle \mathcal{K} \rangle_{\Lambda_b} = -\frac{I_4 - \frac{3}{16} m_0^2 I_2}{I_2 - \frac{1}{16} m_0^2 I_0} \quad (\ell \neq \ell') \quad (3.14)$$

where

$$I_n = \int_0^{\omega_c} d\omega \omega^n e^{-\frac{\omega}{E}}. \quad (3.15)$$

In these relations, the dependence on  $\Delta$  has disappeared, and in principle the analysis could now follow the familiar procedure, checking the existence of a range in the Borel parameter  $E$  where the results for  $\langle \mathcal{K} \rangle_{\Lambda_b}$  are stable or at least smoothly dependent on  $E$ , and at the same time the operator product expansion displays a convergent structure with higher states giving smaller contributions. However, in the present case the two ‘diagonal’ sum rules (3.12) and (3.13) do not give the same result for  $\langle \mathcal{K} \rangle_{\Lambda_b}$ , at least in the approximation of the operator product expansion considered here, in contrast with the two-point function sum rules in (2.16). Thus, the separation of ‘diagonal’ and ‘nondiagonal’ sum rules is no longer useful, and we have to use the complete three-point correlator with the full  $\Lambda_b$  interpolating field of Eq. (2.4), where the *a priori* unknown parameter  $b$  appears. Clearly, the corresponding sum rule

$$\frac{1}{2} \langle \mathcal{K} \rangle_{\Lambda_b} f_{\Lambda_b}^2 (1+b)^2 e^{-\Delta/E} = \int_0^{\omega_c} d\omega e^{-\omega/E} \left[ -\frac{3+5b^2}{140\pi^4} \omega^7 + \frac{2b \langle \bar{q}q \rangle}{\pi^2} \omega^2 \left( \omega^2 - \frac{3m_0^2}{16} \right) \right] \quad (3.16)$$

would give results depending on the parameter  $b$ , whose value must be determined somehow.

In order to get a reasonable criterion to fix the value of  $b$ , we recall that, in principle, the three-point function sum rule can also be used to determine the parameter  $\Delta$ , by taking the derivative in  $1/E$  of (3.16). We can then compare such determination with that given by the two-point function sum rule (2.18) and assume, as an ‘optimal choice’, the value of  $b$  for which the two determinations are closer. In Fig. 2 we depict the ratio of the two determinations *versus*  $b$  and the Borel parameter  $E$ . As one can see, for the considered values of  $E$  such a ratio is close to unity for  $b = 1$ , and therefore we shall assume this value of  $b$  for the subsequent analysis of the sum rule for  $\langle \mathcal{K} \rangle_{\Lambda_b}$ .

In Fig. 3 we report the results of the analysis of the sum rule (3.16) for  $b = 1$ , corresponding to the chosen three values of the continuum threshold  $\omega_c$ . In this analysis we use (2.18) for  $f_{\Lambda_b}$  thus avoiding the assumption  $f_{\Lambda_b}^{(1)} = f_{\Lambda_b}^{(2)}$ . There is a remarkable plateau for  $E$  larger than  $0.25 \text{ GeV}$  indicating a value of  $\langle \mathcal{K} \rangle_{\Lambda_b}$  around  $-0.6 \text{ GeV}^2$ , with an uncertainty of  $0.1 \text{ GeV}^2$  coming from the variation of the continuum threshold in the range  $1.1 - 1.3 \text{ GeV}$ . However, to avoid upsetting the usual QCD sum rule criteria by an excessive contribution from the continuum, in practice the real ‘window’ in  $E$  is reduced to a small range around  $E \sim 0.3 \text{ GeV}$ . This feature, which is due to the highly divergent behavior in  $\Omega$  of the spectral functions, was already anticipated by the analysis of two-point function sum rules. Clearly, such a low value of the Borel parameter would call for the calculation of the order  $\alpha_s$  correction in both the three-point and in the two-point sum rules, a task that goes beyond the aim of the present paper. In the case of the beauty meson  $B$ , it is worth reminding that the radiative corrections turn out to be individually important in two and three point functions, but they compensate in the expression for  $\langle \mathcal{K} \rangle$  confirming the indication of the lowest order [4].

In conclusion, within the approximation of the operator product expansion adopted here, the result is

$$\langle \mathcal{K} \rangle_{\Lambda_b} = -(0.6 \pm 0.1) \text{ GeV}^2. \quad (3.17)$$

Concerning the value quoted in (3.17), we recall that the uncertainty there only accounts for the effect of the variation of the continuum threshold and the Borel parameter; to it one should add the theoretical uncertainty related to the neglect of the radiative corrections and the  $d = 6$  operator in the two-point function. We notice explicitly that the inclusion of the  $d = 6$  term, in the factorization hypothesis, could somehow enhance  $f_{\Lambda_b}$  and reduce  $|\langle \mathcal{K} \rangle_{\Lambda_b}|$ ; however, this result would be numerically reasonable only for values of the Borel parameter  $E$  larger than  $E = 0.4 \text{ GeV}$ , which would contradict our requirement on the continuum suppression. For these reasons, an additional uncertainty of the order of  $\pm 0.1$  in (3.17) cannot be excluded.

Let us finally observe that a gratifying feature of the sum rules is the consistency of the determinations of the mass parameter  $\Delta$  from two and three-point functions. We also notice that the result in (3.17) supports the conjecture that the value of the heavy quark kinetic energy in the baryon is similar to that in the meson.

## ACKNOWLEDGMENTS

We would like to thank P. Ball and V.M. Braun for interesting discussions.

## REFERENCES

- [1] For a review see: M.B. Wise, in Proceedings of the 6th Lake Louise Winter Institute, Lake Louise, 1991, edited by B.A. Campbell, A.N. Kamel, P. Kitching and F.C. Khanna (World Scientific, Singapore, 1991), p. 222; H. Georgi, in Proceedings of the TASI 1991, Boulder, 1991, edited by R.K. Ellis, C.T. Hill and J.D. Lykken (World Scientific, Singapore, 1992), p. 589.
- [2] For a review and references to earlier analyses see: B. Blok and M. Shifman, in Proceedings of the 3rd Workshop on the Tau-Charm Factory, Marbella, 1993, edited by J. Kirkby and R. Kirkby (Editions Frontieres, 1994), p. 247.
- [3] I.I. Bigi, N.G. Uraltsev, A. Vainshtein, Phys. Lett. B 293, 430 (1992); (E) B 297, 477 (1993); B. Blok and M. Shifman, Nucl. Phys. B 399, 441 (1993); B 399, 459 (1993).
- [4] P. Ball and V.M. Braun, Phys. Rev. D 49, 2472 (1994).
- [5] I.I. Bigi, M. A. Shifman, N.G. Uraltsev and A. Vainshtein, Phys. Rev. D 52, 196 (1995).
- [6] V. Eletski and E. Shuryak, Phys. Lett. B 276, 191 (1992); M. Neubert, Phys. Rev. D 46 (1992) 1076; Phys. Lett. B 322, 419 (1994).
- [7] M. Crisafulli, V. Gimenez, G. Martinelli and C.T. Sachrajda, report CERN-TH-7521-94, Rome 94/1071, SHEP 94/95-14.
- [8] I. Bigi, report UND-HEP-95-BIG02 (June 1995).
- [9] Review of Particle Properties, Phys. Rev. D 50 (1994).
- [10] A.V. Manohar and M.B. Wise, Phys. Rev. D 49, 1310 (1994).
- [11] E.V. Shuryak, Nucl. Phys. B 198, 83 (1982).
- [12] A.G. Grozin and O.I. Yakovlev, Phys. Lett. B 285, 254 (1992).
- [13] A.G. Grozin and O.I. Yakovlev, Phys. Lett. B 291, 441 (1992).
- [14] E. Bagan, M. Chabab, H.G. Dosch and S. Narison, Phys. Lett. B 278, 367 (1992); B 287, 176 (1992); B 301, 243 (1993).
- [15] T.-M. Yan, H.-Y. Cheng, C.-Y. Cheung, G.-L. Lin, Y.C. Lin and H.-L. Yu, Phys. Rev. D 46, 1148 (1992).
- [16] H. Georgi, Nucl. Phys. B 348, 293 (1991).
- [17] For a review see: *Vacuum structure and QCD sum rules*, edited by M.A. Shifman, North Holland, Amsterdam, 1992.
- [18] S.V. Mikhailov and A.V. Radyushkin, Phys. Rev. D 45, 1754 (1992).
- [19] A.V. Radyushkin, Phys. Lett. B 271, 218 (1991); M. Neubert, Phys. Rev. D 45, 2451 (1991).

## FIGURE CAPTIONS

### Fig. 1

Results from the two point sum rules Eq. (2.16) (a), Eq. (2.17) (b), and from the correlator of the current  $J$  in Eq. (2.4) with  $b = 1$  (c). The different curves correspond to different choices of the continuum threshold:  $\omega_c = 1.1 \text{ GeV}$  (continuous line),  $\omega_c = 1.2 \text{ GeV}$  (dashed line),  $\omega_c = 1.3 \text{ GeV}$  (dotted line).

### Fig. 2

The ratio of the mass parameters  $\Delta$  obtained from three point (3p) and two points (2p) correlators of the current  $J$  in Eq. (2.4):  $\mathcal{R} = \Delta(3p)/\Delta(2p)$ . The continuum threshold is fixed to  $\omega_c = 1.2 \text{ GeV}$ .

### Fig. 3

$\langle \mathcal{K} \rangle_{\Lambda_b}$  from the three point correlator with the current  $J$  in Eq. (2.4) with  $b = 1$ . The different curves correspond to the continuum thresholds:  $\omega_c = 1.1 \text{ GeV}$  (continuous line),  $\omega_c = 1.2 \text{ GeV}$  (dashed line),  $\omega_c = 1.3 \text{ GeV}$  (dotted line).




fig.1




fig.2




fig.3