

LIFETIME OF QUASIPARTICLES IN HOT QED PLASMAS

Jean-Paul BLAIZOT* and Edmond IANCU†

*Service de Physique Théorique‡, CE-Saclay**91191 Gif-sur-Yvette, France***Abstract**

The calculation of the lifetime of quasiparticles in a QED plasma at high temperature remains plagued with infrared divergences, even after one has taken into account the screening corrections. The physical processes responsible for these divergences are the collisions involving the exchange of very soft, unscreened, magnetic photons, whose contribution is enhanced by the thermal Bose-Einstein occupation factor. The self energy diagrams which diverge in perturbation theory contain no internal fermion loops, but an arbitrary number of internal magnetostatic photon lines. By generalizing the Bloch-Nordsieck model at finite temperature, we can resum all the singular contributions of such diagrams, and obtain the correct long time behaviour of the retarded fermion propagator in the hot QED plasma: $S_R(t) \sim \exp\{-\alpha T t \ln \omega_p t\}$, where $\omega_p = eT/3$ is the plasma frequency and $\alpha = e^2/4\pi$.

*CNRS

†CNRS

‡Laboratoire de la Direction des Sciences de la Matière du Commissariat à l'Énergie Atomique

It is commonly assumed that the notion of quasiparticles is a useful concept in the description of ultrarelativistic plasmas. This implies that the damping rate γ of quasiparticle excitations, obtained from the exponential decay of the retarded propagator, $S_R(t, \mathbf{p}) \sim e^{-i\omega(\mathbf{p})t} e^{-\gamma(\mathbf{p})t}$, is small compared to the quasiparticle energy $\omega(\mathbf{p})$. In hot gauge theories, the typical energy of the relevant quasiparticles is the temperature T , while one expects $\gamma \sim g^2 T$ [1, 2, 3], where g is the gauge coupling (in QED, $g = e$ is the electric charge). The same damping rate is expected for the collective excitations, whose typical energies are $\sim gT$ [2]. This suggests that, indeed, in the weak coupling regime, quasiparticles are well defined, and collective modes are weakly damped. However, the computation of γ in perturbation theory is plagued with infrared divergences, which casts doubt on the validity of these statements [1-7]. We shall show in this paper that the infrared divergences which remain after the screening corrections have been taken into account are due to collisions involving the exchange of very soft, unscreened, magnetic photons. Such divergences occur in all orders of perturbation theory and can be eliminated only by a non perturbative calculation of the quasiparticle propagator. We present in this letter such a non perturbative analysis, based on an extension of the Bloch-Nordsieck model at finite temperature. The final result is that quasiparticles do indeed exist, although they do not correspond to the usual exponential decay indicated above, but to a more complicated behaviour, $S_R(t, \mathbf{p}) \sim e^{-i\omega(\mathbf{p})t} e^{-\alpha T t \ln \omega_p t}$, where ω_p is the plasma frequency.

We shall consider in most of this letter the case of hard fermions, with momenta $p \sim T$, and restrict ourselves to QED. (The case of collective excitations will be discussed briefly at the end.) In perturbation theory, the damping rate is obtained from the imaginary part of the self-energy on the unperturbed mass-shell $\omega = p$:

$$\gamma = -\frac{1}{4p} \text{tr} (\not{p} \text{Im} \Sigma(\omega + i\eta, \mathbf{p})) \Big|_{\omega=p}. \quad (1)$$

To bare one-loop order, $\gamma = 0$ by kinematics : the on-shell fermion can emit (or absorb)

only a space-like photon ($q^2 \equiv q_0^2 - \mathbf{q}^2 \leq 0$), and there are no such photons at tree level (a more complete discussion of this point will be presented in a longer publication [8]). However, through the Landau damping mechanism, the spectral function of a photon in the plasma does acquire a non vanishing weight for space-like momenta [9]. One takes this into account by resumming the so-called hard thermal loops (HTL) in the internal photon line of the one loop self-energy, and obtains in this way the leading contribution to the damping rate [1, 2]. Physically, this contribution represents the effect of the collisions induced by one photon exchange [10]. Note that the corresponding collision rate would be quadratically divergent without the screening corrections [10, 11]. Such corrections are sufficient to render finite the transport coefficients [10], or the collisional energy loss [11], but not the total interaction rate [12], nor the damping rate [1-7]. These quantities, except for the damping rate of a quasiparticle with zero momentum [2, 13], remain logarithmically divergent, both in abelian and non-abelian gauge theories. In QCD, this problem is commonly bypassed by advocating the IR cut-off provided by a possible magnetic mass $\sim g^2 T$ (see e.g. [5]). But such a solution cannot apply for QED where one expects no magnetic screening [14].

In the imaginary time formalism, and in the Coulomb gauge (the one-loop result for γ is gauge independent[15]), the resummed one loop diagram is evaluated as [5]

$$\begin{aligned}
{}^*\Sigma(p) &= -g^2 T \sum_{q_0=i\omega_m} \int \frac{d^3q}{(2\pi)^3} \gamma_\mu S_0(p+q) \gamma_\nu {}^*D^{\mu\nu}(q) \\
&= -g^2 \int \frac{d^3q}{(2\pi)^3} \int_{-\infty}^{+\infty} \frac{dk_0}{2\pi} \int_{-\infty}^{+\infty} \frac{dq_0}{2\pi} \rho_0(k) \gamma_\mu k \gamma_\nu {}^*\rho^{\mu\nu}(q) \frac{N(q_0) + n(k_0)}{k_0 - q_0 - p_0}. \quad (2)
\end{aligned}$$

In this equation, $\mathbf{k} = \mathbf{p} + \mathbf{q}$, $p_0 = i\omega_n = i(2n+1)\pi T$, and $\omega_m = 2\pi mT$, with integers n and m . We denote by $\rho_0(k_0, k)$ the spectral density of the (hard) fermion, and by ${}^*\rho_{l,t}(q_0, q)$ those of the (soft) longitudinal (l) and transverse (t) photons. According to eq. (1), one obtains the damping rate γ from the imaginary part of ${}^*\Sigma$, after analytical continuation of p_0 to the real axis ($p_0 \rightarrow \omega + i\eta$). One thus gets $\gamma(\omega \simeq p) = g^2 p \Phi(\omega, p)$,

where $(\cos \theta = \hat{\mathbf{p}} \cdot \hat{\mathbf{q}})$

$$\begin{aligned} \Phi(\omega, p) \equiv & \int \frac{d^4 q}{(2\pi)^4} \int \frac{d^4 k}{(2\pi)^4} (2\pi)^4 \delta^{(4)}(p + q - k) \\ & [N(q_0) + n(k_0)] \rho_0(k) \left({}^* \rho_l(q) + (1 - \cos^2 \theta) {}^* \rho_t(q) \right), \end{aligned} \quad (3)$$

is the thermal phase-space for the decay of the fermion. The infrared singular region is that of small photon energy-momentum q_0 , $q \ll gT$. Since $\omega, p \sim T$, in studying this region, one can expand $|\mathbf{p} + \mathbf{q}| \simeq p + q \cos \theta$ and $N(q_0) + n(k_0) \simeq N(q_0) \simeq T/q_0 \gg 1$. One obtains then

$$\begin{aligned} \Phi(\omega, p) &= \frac{T}{2p} \int \frac{d^3 q}{(2\pi)^3} \int_{-\infty}^{\infty} \frac{dq_0}{2\pi q_0} 2\pi \delta(\omega - p + q_0 - q \cos \theta) \left({}^* \rho_l(q) + (1 - \cos^2 \theta) {}^* \rho_t(q) \right) \\ &= \frac{T}{4\pi p} \int_0^{\infty} dq q \theta(q - \delta E) \int_{-q}^q \frac{dq_0}{2\pi q_0} \left\{ {}^* \rho_l(q_0, q) + \left(1 - \frac{q_0^2}{q^2} \right) {}^* \rho_t(q_0, q) \right\}, \end{aligned} \quad (4)$$

where the δ -function has been used to perform the angular integration, and $\delta E \equiv |\omega - p| > 0$. Note the restriction to space-like momenta, $|q_0| \leq q$: only the off-shell photons contribute to the fermion damping, which is consistent with the physical interpretation of γ in terms of collisional processes, as alluded to earlier. The evaluation of the q_0 -integral can be done easily by using sum rules satisfied by the spectral function [5]. The singular contribution comes from the transverse component, for which we have

$$\int_{-q}^q \frac{dq_0}{2\pi q_0} {}^* \rho_t(q_0, q) = \frac{1}{q^2} - \frac{z_t(q)}{\omega_t^2(q)}, \quad (5)$$

where $\omega_t(q)$ and $z_t(q)$ are respectively the position and the residue of the transverse plasmon pole. As $q \rightarrow 0$, $z_t(q)/\omega_t^2(q) \rightarrow 1/\omega_p^2$, so that for $q \ll \omega_p \sim gT$ the integral is dominated by the term $1/q^2$. In fact, the function ${}^* \rho_t(q_0, q)/q_0$ is strongly peaked at $q_0 = 0$, and in the calculation of the integral (5) it can be replaced by the following approximate expression:

$$\frac{1}{q_0} {}^* \rho_t(q_0 \ll q) = \frac{3\pi}{2} \frac{\omega_p^2 q}{q^6 + (3\pi\omega_p^2 q_0/4)^2} \xrightarrow{q \rightarrow 0} \frac{2\pi}{q^2} \delta(q_0). \quad (6)$$

Keeping only this leading term, and ignoring the non-singular contribution of the electric

modes, one finds

$$\gamma(\delta E) = g^2 p \Phi(\delta E) \simeq \frac{g^2 T}{4\pi} \int_0^{\omega_p} dq q \theta(q - \delta E) \frac{1}{q^2} = \alpha T \ln \frac{\omega_p}{\delta E}, \quad (7)$$

with $\alpha = g^2/4\pi$. Thus, the singularity of γ when $\delta E = 0$, i.e., when the fermion is on-shell, can be attributed to the low frequency magnetic photons produced by Landau damping, whose contribution is enhanced by the Bose-Einstein occupation factor T/q_0 . Note that since the energy conservation requires $q_0 = q \cos \theta$, these photons are emitted, or absorbed, at nearly 90 degrees.

The upper cut-off $\omega_p \sim gT$ in eq. (7) accounts approximately for the terms which have been neglected when keeping only the $1/q^2$ contribution to the r.h.s of eq. (5): as $q \gg \omega_p$, the r.h.s. of eq. (5) is decreasing like ω_p^2/q^4 , so that the exact integrand in eq. (4) is indeed cut-off at $q \sim \omega_p$. Note that, as long as we are interested only in the coefficient of the logarithm, the precise value of this cut-off is unimportant. The scale ω_p however is uniquely determined by the physical process responsible for the existence of space like photons, i.e., the Landau damping. This is important since, as we shall see later, it is the scale which fixes the long time behavior of the retarded propagator.

In the forthcoming discussion, we shall take advantage of a major technical simplification which is based on the following observation: the logarithm in eq. (7) arises entirely from the magnetic contribution of the term $q_0 = i\omega_m = 0$ in the Matsubara sum of eq. (2) [4, 7]. The analytic continuation of this term to real energy ($p_0 \rightarrow \omega + i\eta$) is well defined, and it yields the following imaginary part

$$\begin{aligned} -\frac{1}{4p} \text{tr}(\not{p} \text{Im}^* \Sigma_0(p)) &\simeq -2g^2 \omega T \int \frac{d^3 q}{(2\pi)^3} \frac{1}{q^2} \text{Im} \frac{1}{(\omega + i\eta)^2 - (\mathbf{p} + \mathbf{q})^2} \\ &\simeq \frac{g^2 T}{4\pi} \int_0^{\omega_p} \frac{dq}{q} \theta(q - \delta E), \end{aligned} \quad (8)$$

where the approximate equality means that only regular terms have been dropped. As anticipated, this coincides with eq. (7). Note that, since the static transverse propagator is not modified by the HTL corrections, eq. (8) could have been obtained directly from

the term $q_0 = i\omega_m = 0$ in the *bare* one-loop self-energy, provided one introduces the same upper cut-off ω_p in the q -integral.

Some remarks are needed at this stage. At the bare one-loop level, $\gamma = 0$, as mentioned. Thus, in the bare one loop calculation, there is a cancellation between the logarithmic divergence of the static mode, and that of the *sum of all* the non-static ones [8]. The inclusion of the screening corrections on the photon line removes the divergence of the contribution of the nonstatic modes, leaving a divergence in the static contribution only. When $\omega \rightarrow p$, the integrand in eq. (8) becomes proportional to $\delta(\cos\theta)$, so that the singular contribution of the static mode originates from the emission or absorption of static magnetic photons at 90 degrees. As we have seen, these processes represent the dominant contribution of the resummed one loop calculation. That they can be taken into account by what looks like a bare one loop calculation is due to the fact that the integrated transverse photon spectral weight (5) is $1/q^2$ at small q , i.e., is indistinguishable from the propagator of a static photon. Finally, as already mentioned, the source of the space-like photons is Landau damping; this is the physical origin of the cut-off ω_p to be imposed on the one loop integral of eq. (8).

Power counting indicates that mass-shell divergences do occur in higher orders as well. The most singular diagrams are those where the fermion propagator is dressed by static magnetic photon lines. When evaluated on the tree-level mass-shell $\omega = p$, and in the presence of an IR regulator μ , such diagrams generate contributions of the type $(g^2 T/\mu)^{n-1}$, where n is the number of loops. Such power-like IR divergences are analogous to those identified in the analysis of the corrections to the screening mass in [16], and their presence signals a breakdown of perturbation theory. The calculation of the fermion propagator in the IR singular domain requires therefore a non perturbative approach, to which we now turn.

Since internal fermion loops are not essential anymore at this stage, we may restrict

ourselves to the “quenched approximation” involving only magnetostatic photons [16]. In this approximation, the imaginary-time fermion propagator can be written as the following functional integral

$$S(x, y) = Z_0^{-1} \int [d\mathbf{A}] G(x, y|\mathbf{A}) \exp \left\{ -\frac{1}{2} \left(\mathbf{A}, D_0^{-1} \mathbf{A} \right)_0 \right\}, \quad (9)$$

where $G(x, y|\mathbf{A})$ is the fermion propagator in the presence of a background gauge field, and

$$\left(\mathbf{A}, D_0^{-1} \mathbf{A} \right)_0 = \frac{1}{T} \int d^3x d^3y A^i(\mathbf{x}) D_{0ij}^{-1}(\mathbf{x} - \mathbf{y}) A^j(\mathbf{y}). \quad (10)$$

In this equation, the factor $1/T$ has its origin in the restriction to the zero Matsubara frequency. The propagator D_{0ij} is that of a free static photon,

$$D_0^{ij}(\mathbf{x}) \equiv \int \frac{d^3q}{(2\pi)^3} e^{i\mathbf{q}\cdot\mathbf{x}} D_0^{ij}(\mathbf{q}) \quad (11)$$

with

$$D_0^{ij}(\mathbf{q}) = \frac{\delta^{ij}}{q^2} + (\lambda - 1) \frac{q^i q^j}{q^4}, \quad (12)$$

in an arbitrary Coulomb or covariant gauge ($\lambda = 0$ corresponds to both the Landau and the strict Coulomb gauges).

The fermion propagator in a *static* background field depends only on the time difference $x_0 - y_0$, $G(x, y|\mathbf{A}) \equiv G(x_0 - y_0, \mathbf{x}, \mathbf{y}|\mathbf{A})$. Its Fourier transform can be analytically continued in the complex energy plane, and the resulting function coincides, in the upper half plane, with the retarded propagator. It is then convenient to take the Fourier transform of eq. (9). To this aim, we write ($p^\mu = (i\omega_n, \mathbf{p})$, $\omega_n = (2n + 1)\pi T$):

$$G(x, y|\mathbf{A}) = T \sum_n \int \frac{d^3p}{(2\pi)^3} e^{-ip \cdot (x-y)} G(p; \mathbf{x}|\mathbf{A}), \quad (13)$$

and similarly for S . We obtain

$$S(p) \equiv Z_0^{-1} \int [d\mathbf{A}] G(p; \mathbf{x}|\mathbf{A}) \exp \left\{ -\frac{1}{2} \left(\mathbf{A}, D_0^{-1} \mathbf{A} \right)_0 \right\}. \quad (14)$$

Since the energy p_0 enters eq. (14) as an external parameter, the continuation to real external energy $p_0 \rightarrow \omega + i\eta$, and the Fourier transform to real time, can both be performed *before* doing the functional integration [8]. This offers the possibility to calculate the retarded propagator $S_R(t, \mathbf{x})$ by directly inserting an approximate expression for $G_R(x, y|A)$ in the functional integral (9).

In the kinematical regime of interest, an approximate expression for $G_R(x, y|\mathbf{A})$ is obtained by neglecting the recoil of the fermion in the successive emissions or absorptions of very soft photons. More precisely, we note that inside the diagrams generated by eq. (9), we can approximate the fermion propagators by

$$S_0(\omega, \mathbf{p} + \mathbf{q}) = \frac{-\omega\gamma_0 + (\mathbf{p} + \mathbf{q}) \cdot \boldsymbol{\gamma}}{(\omega + i\eta)^2 - \epsilon_{p+q}^2} \simeq \frac{-1}{\omega - \epsilon_p - \mathbf{v} \cdot \mathbf{q} + i\eta} \frac{\gamma_0 - \hat{\mathbf{p}} \cdot \boldsymbol{\gamma}}{2}, \quad (15)$$

where \mathbf{q} is a linear combination of the internal photons momenta and $\mathbf{v} \equiv \partial\epsilon_p/\partial\mathbf{p}$ ($\mathbf{v} = \hat{\mathbf{p}}$ for the ultrarelativistic fermion). This is the familiar structure encountered in most treatments of IR divergences in QED (see, e.g., [17]) and which is economically exploited within the Bloch-Nordsieck model (see, e.g. [18]). In this model, the propagator with the desired IR structure is obtained in coordinate space as the solution to

$$-i(v \cdot \partial_x) G_0(x, y) = \delta(x - y), \quad (16)$$

where $v^\mu = (1, \mathbf{v})$. In the presence of an arbitrary background field, the corresponding propagator, $G(x, y|A)$, satisfies the following equation ($D_\mu = \partial_\mu + igA_\mu$)

$$-i(v \cdot D_x) G(x, y|A) = \delta(x - y), \quad (17)$$

which can be solved *exactly*. For retarded boundary conditions, and for static fields:

$$G_R(x, y|\mathbf{A}) = i\theta(x_0 - y_0) \delta^{(3)}(\mathbf{x} - \mathbf{y} - \mathbf{v}(x_0 - y_0)) U(x, y) \quad (18)$$

where $U(x, y)$ is the parallel transporter

$$U(x, x - vt) = \exp \left\{ ig \int_0^t ds \mathbf{v} \cdot \mathbf{A}(\mathbf{x} - \mathbf{v}s) \right\}. \quad (19)$$

The retarded propagator $S_R(x - y)$ is calculated by inserting the expressions (18), (19) of $G_R(x, y|\mathbf{A})$ in the functional integral (14). It can be written as

$$S_R(t, \mathbf{p}) = i\theta(t)e^{-it(\mathbf{v} \cdot \mathbf{p})} \Delta(t), \quad (20)$$

where

$$\Delta(t) \equiv Z_0^{-1} \int [d\mathbf{A}] U(x, x - vt) \exp \left\{ -\frac{1}{2} (\mathbf{A}, D_0^{-1} \mathbf{A})_0 \right\} \quad (21)$$

contains all the non-trivial time dependence. The functional integral is easily done:

$$\Delta(t) = \exp \left\{ \frac{g^2}{2} T \int_0^t ds_1 \int_0^t ds_2 v^i D_0^{ij}(\mathbf{v}(s_2 - s_1)) v^j \right\}. \quad (22)$$

In this equation, $D_0^{ij}(\mathbf{x})$ is the coordinate space representation of the magnetostatic photon propagator (see eq (11)).

The s_1 and s_2 integrations in eq. (22) can be done by going to the Fourier representation:

$$\Delta(t) = \exp \left\{ -g^2 T \int \frac{d^3 q}{(2\pi)^3} \frac{\tilde{D}(\mathbf{q})}{(\mathbf{v} \cdot \mathbf{q})^2} (1 - \cos t(\mathbf{v} \cdot \mathbf{q})) \right\}, \quad (23)$$

where $v^i D_0^{ij}(\mathbf{q}) v^j \equiv \tilde{D}(\mathbf{q})$. The integral in eq. (23) is identical to that one would get in the Bloch-Nordsieck model in 3 dimensions. It has no infrared divergence, but one can verify that the expansion of $\Delta(t)$ in powers of g^2 generates the most singular pieces of the usual perturbative expansion for the self-energy[8]. One can also verify that the integral in eq. (23) presents an ultraviolet logarithmic divergence. However, one should recall that the restriction to the static photon mode implies that such an integral is to be cut off at momenta $q \sim \omega_p$ (cf. the discussion after eq. (7)).

The calculation of $\Delta(t)$ is most simply done using the coordinate space representation (22). In the Feynman gauge $\lambda = 1$, we use $v^i D_0^{ij}(\mathbf{x}) v^j = 1/4\pi x$ and obtain

$$\begin{aligned} & \frac{1}{2} \int_0^t ds_1 \int_0^t ds_2 v^i D_0^{ij}(\mathbf{v}(s_2 - s_1)) v^j \\ &= \frac{1}{8\pi} \int_0^t ds_1 \int_0^t ds_2 \frac{\theta(|s_1 - s_2| - 1/\omega_p)}{|s_1 - s_2|} \simeq \frac{t}{4\pi} \ln \omega_p t. \end{aligned} \quad (24)$$

where the ultraviolet cut-off is introduced in the function $\theta(|s_1 - s_2| - 1/\omega_p)$. This expression is valid for times $t \gg 1/\omega_p$. It is insensitive to the specific procedure which is used to implement the cut-off [8], and independent of gauge fixing (the gauge-dependent contribution to the integral above, as given by the last term of the photon propagator (11), is $(\lambda - 1) \frac{t}{8\pi}$, and therefore subleading).

At times $t \gg 1/\omega_p$ the function $\Delta(t)$ is thus of the form [19]

$$\Delta(\omega_p t \gg 1) \simeq \exp(-\alpha T t \ln \omega_p t). \quad (25)$$

A measure of the decay time τ is given by

$$\frac{1}{\tau} = \alpha T \ln \omega_p \tau = \alpha T \left(\ln \frac{\omega_p}{\alpha T} - \ln \ln \frac{\omega_p}{\alpha T} + \dots \right). \quad (26)$$

Since $\alpha T \sim g\omega_p$, $\tau \sim 1/(g^2 T \ln(1/g))$. This corresponds to a damping rate $\gamma \sim 1/\tau \sim g^2 T \ln(1/g)$, similar to that obtained in a one loop calculation with an IR cut-off $\sim g^2 T$ (cf. eq. (7)).

However, contrary to what perturbation theory predicts, $\Delta(t)$ is decreasing faster than any exponential. It follows that the Fourier transform

$$S_R(\omega, \mathbf{p}) = \int_{-\infty}^{\infty} dt e^{-i\omega t} S_R(t, \mathbf{p}) = i \int_0^{\infty} dt e^{it(\omega - \mathbf{v} \cdot \mathbf{p} + i\eta)} \Delta(t), \quad (27)$$

exists for *any* complex (and finite) ω . Thus, the retarded propagator $S_R(\omega)$ is an entire function, with sole singularity at $\text{Im } \omega \rightarrow -\infty$.

The previous analysis can be extended [8] to quasiparticles with *soft* momenta, $p \sim gT$, for which the same infrared difficulty arises [5]. Because of the collective nature of the soft quasiparticles, the HTL resummations to be performed in this case are more elaborate [2]. It can be shown however that the leading divergences are again confined to the magnetostatic photon sector [7], and occur in diagrams which involve only the 3-point photon-fermion vertex [8] (to one-loop order, this has been noticed already in Refs.[5, 7]). In the kinematical regime of interest, one finds [8] that the approximate

fermion propagator and 3-point vertex are of the form:

$$S_{\pm}(\omega, \mathbf{p} + \mathbf{q}) = \frac{-z_{\pm}(p)}{\omega - \omega_{\pm}(p) - \mathbf{v}_{\pm}(p) \cdot \mathbf{q}} \frac{\gamma_0 \mp \hat{\mathbf{p}} \cdot \boldsymbol{\gamma}}{2}, \quad \Gamma_{\pm}^i = \frac{v_{\pm}^i}{z_{\pm}} \frac{\gamma_0 \pm \hat{\mathbf{p}} \cdot \boldsymbol{\gamma}}{2}. \quad (28)$$

In these equations, the subscripts \pm refer to the two positive-energy modes of the soft electron[20], with energies $\omega_{\pm}(p)$ and residues $z_{\pm}(p)$, and $\mathbf{v}_{\pm}(p) \equiv \partial\omega_{\pm}(p)/\partial\mathbf{p}$. The main approximation used in the present paper remains meaningful for a soft quasiparticle whose momentum $p \sim gT$ is much larger than the typical momenta $q \lesssim g^2T$ of the off-shell photons which are responsible for the leading IR singularity. By the same steps as before, we obtain the retarded propagator for the two fermionic modes \pm in the form

$$\begin{aligned} S_{\pm}(\omega, \mathbf{p}) &= i z_{\pm}(p) \int_0^{\infty} dt e^{it(\omega - \omega_{\pm}(p) + i\eta)} \Delta_{\pm}(t), \\ \Delta_{\pm}(t) &= \Delta(|v_{\pm}|t), \end{aligned} \quad (29)$$

with the function $\Delta(t)$ given by eq. (23).

To conclude, we have shown that, in the weak coupling regime, quasiparticles excitations in the ultrarelativistic QED plasma are slowly damped. Their spectral density retains the shape of a *resonance* strongly peaked around the perturbative mass-shell $\omega \sim \mathbf{v} \cdot \mathbf{p}$, with a typical width of order $\sim g^2T \ln(1/g)$. No singularity is associated to this resonance: the retarded propagator is an entire function in the complex energy plane.

References

- [1] R.D. Pisarski, Phys. Rev. Lett.**63** (1989) 1129.
- [2] E. Braaten and R.D. Pisarski, Phys. Rev. Lett.**64** (1990) 1338; Phys. Rev. **D42** (1990) 2156; Nucl. Phys. **B337** (1990) 569.
- [3] V.V Lebedev and A.V. Smilga, Phys. Lett. **B253** (1991) 231; Ann. Phys.**202** (1990) 229; Physica **A181** (1992) 187.

- [4] C.P. Burgess and A.L. Marini, Phys. Rev. **D45** (1992) R17; A.K. Rebhan, Phys. Rev. **D46** (1992) 482.
- [5] R.D. Pisarski, Phys. Rev. **D47** (1993) 5589.
- [6] T. Altherr, E. Petitgirard and T. del Rio Gaztelurrutia, Phys. Rev. **D47** (1993) 703; S. Peigné, E. Pilon and D. Schiff, Z. Phys. **C60** (1993) 455; A.V. Smilga, Phys. Atom. Nuclei **57** (1994) 519; R. Baier and R. Kobes, Phys. Rev. **D50** (1994) 5944; A. Niégawa, Phys. Rev. Lett. **73** (1994) 2023.
- [7] F. Flechsig, H. Schulz and A.K. Rebhan, Phys. Rev. **D52** (1995) 2994.
- [8] J.P. Blaizot and E. Iancu, *in preparation*.
- [9] H.A. Weldon, Phys. Rev. **D26** (1982) 1394.
- [10] G. Baym, H. Monien, C.J. Pethick, and D.G. Ravenhall, Phys. Rev. Lett. **64** (1990) 1867.
- [11] E. Braaten and M.H. Thoma, Phys. Rev. **D44** (1991) 1298.
- [12] H. Heiselberg and C.J. Pethick, Phys. Rev. **D47** (1993) R769.
- [13] R. Kobes, G. Kunstatter and K. Mak, Phys. Rev. **D45** (1992) 4632; E. Braaten and R. Pisarski, Phys. Rev. **D46** (1992) 1829.
- [14] E. Fradkin, Proc. Lebedev Phys. Inst. **29** (1965) 7; J.P. Blaizot, E. Iancu and R. Parwani, Phys. Rev. **D52** (1995) 2543.
- [15] A.K. Rebhan, Phys. Rev. **D46** (1992) 4779.
- [16] J.P. Blaizot and E. Iancu, “*Non perturbative aspects of screening phenomena in abelian and non abelian gauge theories*”, preprint Saclay T95/055, to appear in Nucl. Phys. **B**.

- [17] S. Weinberg, “The Quantum Theory of Fields” (Cambridge University Press, 1995).
- [18] N.N. Bogoliubov and D.V. Shirkov, “Introduction to the Theory of Quantized Fields” (Interscience Publishers, inc, New-York, 1959).
- [19] In a recent paper by K. Takashiba, *Thermal two point function of a heavy muon in hot QED plasma within Bloch Nordsieck Approximation*, hep-ph 9501223 (unpublished), a result similar to our eq. (25) has been reported. The derivations presented in this paper contain questionable conceptual and technical aspects, of which we shall mention just one. In that work, all the screening (HTL) corrections in the soft photon line are effectively ignored, except for the Debye mass $m_D \sim gT$ in the electrostatic mode. Thus the non-static sector is the same as in the *bare* theory. One expects therefore the ultraviolet behavior to be controlled by the thermal distribution functions, rather than by the Landau damping, i.e., the UV cut-off appearing in the calculation of the static mode contribution should be T , instead of gT . The fact that the large time behavior of the fermion propagator is found to be controlled by the soft scale gT (as in eq. (25)) is due to an accidental compensation at large momenta between the magnetostatic and the electrostatic propagators providing a cut-off at $q \sim m_D$. But this is not the correct physics. In the expression (4), the electric and magnetic sectors are independently cut off at momenta $\sim gT$ because of the Landau damping.
- [20] V.V. Klimov, Sov. J. Nucl. Phys. **33** (1981) 934; Sov. Phys. JETP **55** (1982) 199.