

RUB-TP II-28/95

January 11, 1996

Decay of high density matter in the electroweak theory

Jörg Schaldach, Peter Sieber, Dmitri Diakonov¹,
and Klaus Goeke²

Inst. für Theor. Physik II, Ruhr-Universität Bochum, D-44780 Bochum, Germany
St. Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg 188350, Russia

Abstract

High density fermion matter is meta-stable due to the anomalous non-conservation of baryon and lepton numbers in the electroweak theory. The meta-stable state decays by penetrating the sphaleron barrier separating topologically different vacua. The decay happens locally, and results in an annihilation of twelve fermions, accompanied by a production of gauge and Higgs bosons. We find numerically local bounce solutions determining the decay rate, which are classical paths in imaginary time, connecting two adjacent topological sectors. We also follow the real-time evolution of the bosonic fields after the tunneling and analyze the spectrum of the created bosons.

¹ diakonov@lnpi.spb.su

² goeke@hadron.tp2.ruhr-unibochum.de

1. It is well known that transitions between topologically different sectors of the electroweak theory are accompanied by a change of the fermion number [1]. The height of the separating sphaleron barrier is about $E_{\text{sphal}} \approx 10 \text{ TeV}$ [2] and accordingly the tunneling rate is extraordinarily small; it is exponentially suppressed by the factor $\exp(-2S_{\text{inst}}) \approx 10^{-153}$ with the instanton action $S_{\text{inst}} = 8\pi^2/g^2$; $g \approx 0.67$. This prevents baryon and lepton number violation from being an observable phenomenon under ordinary conditions.

This suppression is less strong, or the transition is even unsuppressed at all, if the temperature [3, 4, 5], the particle energy [6, 7, 8], or the fermion density [9, 10] is large. Hence, fermion number violating processes might have played a role in the earlier history of the Universe or could perhaps be realized in future supercolliders.

In this paper we assume to have zero (or very low) temperature but a macroscopic amount of fermions of a very high density in thermal equilibrium. We describe them by the chemical potential μ ; since we have zero temperature this is the energy up to which the fermionic levels are occupied. In a sphaleron transition one level crosses the gap, all others are shifted such that after the transition we have the same spectrum again. But now one more level is occupied or depleted, depending on whether the levels went up or down, hence the created or annihilated fermion has the energy μ .

The fermion number N_f of each doublet changes with the Chern-Simons number N_{CS} of the bosonic fields as $N_f = N_{CS}$. It means that the potential energy of the gauge and Higgs bosons is shifted [9, 10]:

$$V_{\text{pot}} = V_{\text{pot}} + N_{CS} \mu \quad (1)$$

Some remarks are necessary here: First, we have fixed the zero-point of the energy to the trivial vacuum with $N_{CS} = 0$. Second, we may well neglect the change of the Fermi energy as due to the creation or annihilation of a few fermions. Third, $N_f = N_{CS}$ actually applies to all fermion doublets, which also might have different chemical potentials. Therefore in eq. (1) we imply to be the sum of all these single chemical potentials.

The former degenerate ground states of the system with integer N_{CS} now have energy $N_{CS} \mu$ and hence are metastable (Fig. 1). They can decay by tunneling through the sphaleron barrier to the adjacent vacuum with lower energy. If μ exceeds a certain value μ_{crit} , the states become even unstable.

In [10], arguments were given that the tunneling transition rate corresponds




Fig. 1: Schematic plot of the tunneling process between two topological sectors.

to the baryon number violation rate at high accelerator energies. Thus the results of the present paper may also have some relevance for the high energy transition rate.

The decay of metastable "false" vacua was discussed by Coleman already in 1977 [11]. The transition rate per volume can be calculated in the semiclassical WKB approximation, it is of the form

$$\Gamma/V = A \exp(-B) : \quad (2)$$

Here B is twice the Euclidean action of the classical path that connects the decaying state and the escape point at the other side of the barrier and minimizes this action. It is a movement in the inverted potential; the system leaves the initial state, reaches the escape point with zero kinetic energy and would then move back the same way. Coleman called it "the bounce". The prefactor A can be found from the small oscillation determinant about a bounce, with one negative mode removed. In the following we present the first numerical determination of the bounce in electroweak theory.

2. We consider the minimal version of the standard electroweak theory with one Higgs doublet in the limit of vanishing Weinberg angle. In terms of dimen-

dimensionless rescaled quantities the bosonic part of the Lagrangian is

$$\mathcal{L} = m_W^4 \left[\frac{1}{g^2} \left(\frac{1}{4} F^a_{ij} F^{ij} + \frac{1}{2} (D_\mu \phi)^\dagger (D_\mu \phi) - \frac{1}{32} (Y - 4)^2 \right) \right] \quad (3)$$

with the covariant derivative $D_\mu = \partial_\mu - iA_\mu$; $A_\mu = \frac{1}{2} A^a_\mu \tau^a$, the field strength $F_{ij} = \frac{1}{2} F^a_{ij} \tau^a = i[D_i, D_j]$; $F^a_{ij} = \partial_i A^a_j - \partial_j A^a_i + \frac{1}{2} \epsilon^{abc} A^b_i A^c_j$, and the Higgs doublet $\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}$. $\kappa = m_H/m_W$ is the ratio of Higgs and gauge boson masses. The fermions are not considered explicitly here, but by the modification (1) of the potential energy of the bosonic fields. We work entirely in temporal gauge, $A_0 = 0$. The potential and kinetic energy and the Chern-Simons index are

$$\begin{aligned} V_{\text{pot}} &= \frac{m_W}{g^2} \int d^3r \left[\frac{1}{4} (F^a_{ij})^2 + \frac{1}{2} (D_i \phi)^\dagger (D_i \phi) + \frac{1}{32} (Y - 4)^2 \right]; \\ T_{\text{kin}} &= \frac{m_W}{g^2} \int d^3r \left[\frac{1}{2} (A^a_i)^2 + \frac{1}{2} Y^2 \right]; \\ N_{\text{CS}} &= \frac{1}{16\pi^2} \int d^3r \left[\epsilon_{ijk} A^a_i \partial_j A^a_k + \frac{1}{3} \epsilon_{abc} A^a_i A^b_j A^c_k \right]; \end{aligned} \quad (4)$$

N_{CS} is only well-defined if the configuration space can be identified with the sphere S_3 , which requires the fields to be continuous at infinity. We fix them to the trivial vacuum there ($A = 0$; $\phi = \begin{pmatrix} 0 \\ 2 \end{pmatrix}$). In general, we will look for fields that minimize the action and expect them to possess higher symmetries than completely arbitrary fields. Therefore we assume the fields to have the spherical symmetry of the sphaleron:

$$\begin{aligned} A^a_i(t; r) &= \epsilon_{aij} n_j \frac{1}{r} \frac{A(t; r)}{r} + (\epsilon_{aij} n_j n_i) \frac{B(t; r)}{r} + n_a n_i D(t; r); \\ (t; r) &= 2 H(t; r) + i G(t; r) n_i \frac{0}{1} : \end{aligned} \quad (5)$$

We assume the fields to be continuous and differentiable everywhere and to yield finite potential and kinetic energy. This poses some additional conditions on the radial functions at $r = 0$.

For negative chemical potential μ , the false vacuum with $N_{\text{CS}} = 0$ can decay to the adjacent topological sector with $N_{\text{CS}} = 1$, where the ground state has energy < 0 . The minimal energy barrier between the two sectors is given by the configurations of Akiba, Kikuchi, and Yanagida (AKY) [2] which are static fields that minimize the potential energy V_{pot} for given values of N_{CS} (Fig. 1).

The barrier vanishes completely if μ exceeds a critical value μ_{crit} . From the condition that for $\mu > \mu_{\text{crit}}$ the potential V_{pot} gets a negative mode around the vacuum, one finds [9, 10]

$$\mu_{\text{crit}} = \frac{16}{g^2} m_W^2 : \quad (6)$$

It is difficult to determine the bounce by solving numerically the equations of motion as an initial value problem. If we tried to find the propagation in time with a Runge-Kutta method or something similar, the system would always fall into some abyss of the potential V_{pot} . Instead, we rather prefer to find a stationary point of the Euclidean action

$$S_E = \int_1^{Z_{t_0}} dt T_{\text{kin}} + V_{\text{pot}} \quad (7)$$

directly, without using the equations of motion. t_0 is the time when the system reaches the escape point; since the lower bound is 1, we can choose an arbitrary value for t_0 , just as we fixed the origin for our spherical ansatz (5) somewhere. The corresponding translational invariance of the action does not influence the factor B in eq. (2), but contributes a space-time volume which allows to have a transition probability per time and volume.

S_E is a functional of the five functions $A; B; D; H; G$ of eq. (5), depending on radial distance r and time t . Since the bounce has infinite extension both in space and in time, we introduce new variables x and u which cover only finite intervals. In practice, we use for example

$$r(x) = r \arctan \frac{x}{2} \quad \text{and} \quad t(u) = t \arctan \frac{u}{2} : \quad (8)$$

Using ansatz (5) and the substitution (8) we get

$$\begin{aligned} S_E = & \frac{S_{\text{inst}}}{2} \int_1^{Z_{u_0}} du \int_0^{Z_1} dx \frac{1}{r^2} (A^2 + B^2 + 2r^2(H^2 + G^2 + \frac{1}{4}D^2) \\ & + (A^0 + BD)^2 + (B^0 - AD)^2 + 2r^2(H^0 + \frac{1}{2}GD)^2 \\ & + 2r^2(G^0 - \frac{1}{2}HD)^2 + \frac{1}{2r^2}(A^2 + B^2 - 1)^2 + (H^2 + G^2)(A^2 + B^2 + 1) \\ & + 2A(G^2 - H^2) - 4BGH + \frac{1}{2}r^2(H^2 + G^2 - 1)^2 \\ & + 2D(A^2 + B^2 - 1) + B A^0 - (A - 1)B^0 \end{aligned} \quad (9)$$

with

$$\frac{1}{r^2} = \frac{1}{r_{\text{crit}}^2}; \quad \dot{} = \dot{}(x) = \frac{dr}{dx} \frac{1}{r}; \quad \dot{} = \dot{}(u) = \frac{dt}{du} \frac{1}{t}; \quad (10)$$

and the dot and prime mean $\frac{d}{du}$ and $\frac{d}{dx}$, respectively. In order to find a stationary point of S_E numerically, we use a relaxation method which was discussed by Adler and Piran in great detail [12]. The functional is put on a grid, let us say of size $n_u \times n_x$, and the values of the five functions at the grid nodes are considered as $5n_u n_x$ independent variables. Of course there is no unique

way to discretize a functional, basically we followed the suggestions of [12]. We postpone the details to a more comprehensive publication. After putting some initial configuration on the grid, we sweep over it by changing the single variables one after another. Each variable is changed according to the first iteration step of a Newtonian algorithm, which would find the stationary point of S_E if it was considered as a function of the variable in question alone and all others were constants. This method is somewhat different from a procedure presented in [13] where a similar problem in the context of technibaryons in the Skyrme model is solved. As mentioned above, the fields at $x = 1$ ($r = 1$) are fixed to the trivial vacuum, and some conditions must be obeyed at the origin. We take care that the initial configuration fulfills these properties and that we do not lose them during the sweeps.

In principle, one has to perform some hundred or thousand sweeps until the situation is stable and the resulting configuration is a stationary point of S_E . But there are some difficulties which must be handled carefully to get proper results:

The main problem is that V_{pot} is not bound, it can be negative. Due to energy conservation the bounce itself cannot have positive potential energy V_{pot} at any time, its total energy $E_{\text{tot}} = T_{\text{kin}} + V_{\text{pot}}$ is constant and zero. But in its vicinity, one can construct configurations which have positive potential V_{pot} for some time and which give a lower action than the bounce. So actually the bounce is just a saddle point of S_E , it is a minimum only if we restrict the potential energy V_{pot} to be non-positive. So we enforce this restriction, simply by rejecting all changes of fields rendering V_{pot} to be positive when we sweep. There are more sophisticated ways to take into account invariances like energy conservation [14], but in our case the simple remedy proved to be most effective. Besides, we use energy conservation to check if a given configuration is already close to the bounce and if the numerical accuracy (e.g. grid size) is sufficient.

Another point is that the solution is not unique: We want to describe the bounce between the topological sectors with $N_{CS} = 0$ and 1; but this still allows so-called "time-independent" gauge transformations which do not change the Chern-Simons number. We choose the $N_{CS} = 0$ vacuum to be the trivial one, which fixes the gauge completely. Practically, we have a boundary condition for the $u = 1$ edge of the grid, but due to the large differences in the true time near this boundary, the field variables are only weakly coupled there. We often observe that the fields tend to another vacuum near the boundary,

influenced by the main configuration at larger t . In order to have a smooth behavior, we sometimes apply a gauge transformation to the whole grid except the close vicinity of the boundary which already is in the trivial vacuum.

Another degree of freedom is the choice of t_0 , the time when the bounce reaches the turning point, i.e. the potential energy becomes zero again. For the initial configuration this point lies on the right end of the grid, during the sweeps we observe that it moves slowly to the left. For $t > t_0$ the potential energy remains zero then (in the frame of our numerical accuracy). Actually, this shift to the left never stops completely, but finally the system moves as a whole, it just follows the zero-modes due to the free choice of t_0 .

A more detailed description of the numerical technique will be given in a subsequent paper.

In Table 1 we give our results for the bounce action and the Chern-Simons number of the escape point for $\beta = 1$ (i.e. $m_H = m_W$) and several values of α . The cases $\alpha = 0$ and 1 were not treated numerically, but clearly for $\alpha \neq 1$ the action vanishes since the barrier disappears. For $\alpha = 0$ it is known that we have the instanton action. Actually, the finite vacuum expectation value of the Higgs field destroys the scaling invariance of the instanton which we have in the case of pure gauge fields, so the bounce is an instanton in the limit of size zero here [15]. Only the introduction of the second massive parameter allows the bounce to have finite size again in the other cases.

	$S_E = S_{\text{inst}}$	N_{CS}^{esc}	N_W	N_H	E_W	E_H	$E_W + E_H$
0:0	1:00	1:00					
0:2	0:82	0:85	22:9	1:9	0:370	0:020	0:391
0:4	0:58	0:68	46:2	5:9	0:727	0:060	0:786
0:6	0:34	0:51	67:1	9:6	1:093	0:102	1:195
0:8	0:14	0:30	102:6	16:8	1:417	0:180	1:597
1:0	0:00	0:00					

Tab. 1: Bounce action, Chern-Simons number at the escape point, and the number of gauge and Higgs bosons and their energies after the tunneling for several chemical potentials $\alpha = \frac{m_H^2}{m_W^2}$; the energies are given in units of $\frac{8\pi^2}{g^2} m_W = \frac{4\pi}{\alpha}$. The mass of the Higgs boson is fixed to $m_H = m_W$.

One might expect that S_E could have many different stationary points,

in other words, there could be several bounce solutions with different escape points and values of the action. As a matter of fact, for our numerics we used at least two distinct initial configurations for each value of j , but always found the same solution.

Fig. 2 shows the time dependence of the potential energy V_{pot} and the Chern-Simons number N_{CS} , with t_0 fixed to zero. In Fig. 3 we plotted V_{pot} versus N_{CS} and can compare this to the minimal energy barrier. Obviously, the kinetic energy terms force the system to take a path which deviates quite a bit from a possible path through the A-K-Y configurations minimizing the potential energy only. As one may expect, the deviation grows with the height of the barrier.

3. Next we want to examine the behavior of the system after the tunneling. To this end, we take the (numerically found) escape point configuration and let it propagate in the real Minkowski time. We work on a two-dimensional grid again, but at this time we can solve the equations of motion using a kind of Runge-Kutta method. This was done in [16, 17] with a slightly disturbed sphaleron as initial configuration; we checked that our algorithm gives the same results in this case.

We have to distinguish two different kinds of behavior. The system can either stay in the topological sector where it came to after the tunneling or move classically over the barrier towards the next ground state with even lower energy. For $j < 0.2$, the system contains less energy than required to cross the next barrier, so that it will definitely stay in its topological sector. For values of j between about 0.2 and 0.9 the system has enough total energy so that in principle it could move over the barrier. The system, however, stays in its sector, because the energy is dissipated along the excitation modes in such a way that the system does not find a collective path towards the next vacuum. Finally for $j > 0.9$ the system actually moves classically into the next sector, and subsequently also crosses the following barriers, i.e. it keeps moving into the direction of growing N_{CS} .

In the case when the system stays in its topological sector, we see that after some short time the potential and kinetic energies become almost constant. For the transition described above, from $N_{\text{CS}} = 0$ to 1, we observe very small oscillations of T_{kin} around $j \neq 2$ and of V_{pot} around $j \neq 2$, which is in accordance with the virial theorem. The energy is concentrated near a sphere which




Fig. 2: Chern-Simons number N_{CS} and potential energy $-V_{\text{pot}}^\mu$ versus time t (in units of m_W^{-1}) for different values of the chemical potential $\rho = \mu = \mu_{\text{crit}}$.

expands with about 95% of the speed of light. N_{CS} becomes almost constant after a while, with small variations; but this constant is not the integer number which classifies the topological sector. Nevertheless, the fermion number is changed by one for each doublet: The equation $N_f = N_{CS}$ is not valid here because in the Minkowskian setup the non-vacuum fields cannot be restricted to a finite region [18].

Finally, we find that the time-averaged fields $\langle \frac{1}{b-a} \int_a^{R_b} dt \text{ with } b \neq 1 \rangle$ form a static vacuum configuration A_i^a ; with $V_{\text{pot}} = 0$ and integer N_{CS} , and that at larger times the fields can be considered as small fluctuations around that vacuum. This allows to analyze the particle contents of the bosonic fields,




Fig. 3: The potential energy $-V_{\text{pot}}^{\mu}$ versus the Chern-Simons number N_{CS} for various values of the chemical potential $\mu = \mu_{\text{crit}}$. The solid lines correspond to the configurations which minimize the action (7, 9) (bounce trajectories), the dashed lines are the minimal energy paths (A KY configurations).

we can perform a spectral decomposition in the basis of free particle modes [16]. We apply a gauge transformation to the fields which converts A_i^a to the trivial vacuum first, hence we work with fluctuations $a; b; d; h; g$ around the latter one. The basis is found by solving the linearized equations of motion; basically we follow [16] here, but use V_{pot}^{μ} instead of V_{pot} . The free particle modes in our spherical ansatz are

$$\begin{aligned}
 a_k(t; r) &= \frac{3}{2} \begin{pmatrix} k \\ 1 \end{pmatrix} \begin{pmatrix} k \\ 1 \end{pmatrix} (t) \begin{pmatrix} k \\ 2 \end{pmatrix} (t) r j_1(kr) \\
 b_k(t; r) &= \begin{pmatrix} k \\ 0 \end{pmatrix} (t) \begin{pmatrix} k \\ 1 \end{pmatrix} (t) \begin{pmatrix} k \\ 2 \end{pmatrix} (t) r j_2(kr) \\
 &+ \begin{pmatrix} k \\ 0 \end{pmatrix} (t) + \begin{pmatrix} k \\ 1 \end{pmatrix} (t) + \begin{pmatrix} k \\ 2 \end{pmatrix} (t) r j_0(kr) \\
 d_k(t; r) &= 2 \begin{pmatrix} k \\ 0 \end{pmatrix} (t) + \begin{pmatrix} k \\ 1 \end{pmatrix} (t) + \begin{pmatrix} k \\ 2 \end{pmatrix} (t) j_2(kr) \\
 &+ \begin{pmatrix} k \\ 0 \end{pmatrix} (t) + \begin{pmatrix} k \\ 1 \end{pmatrix} (t) + \begin{pmatrix} k \\ 2 \end{pmatrix} (t) j_0(kr) \\
 h_k(t; r) &= h^k(t) j_0(kr)
 \end{aligned} \tag{11}$$

$$g_k(t; r) = \frac{3}{2} \int_0^k dt k j_1(kr)$$

with

$$j_i^k(t) = j_i^k \sin(j_i t + \phi_i); \quad h^k(t) = h^k \sin(t + \phi); \quad (12)$$

$$j_0^2 = 1 + k^2; \quad j_{1/2}^2 = 1 - 2k + k^2; \quad j^2 = j^2 + k^2; \quad (13)$$

The j_s are the spherical Bessel functions. For each time t and momentum k we can get the functions $j_i^k(t)$ and $h^k(t)$ from radial "Fourier" integrals; a comparison with (12) yields the coefficients j_i^k and h^k .

In the linearized theory, the total energy is

$$E = E_W + E_H = m_W \int_0^{Z_1} dk n_W(k) j_0^2(k) + m_W \int_0^{Z_1} dk n_H(k) (h^k)^2 \quad (14)$$

with the spectral densities

$$\begin{aligned} n_0(k) &= \frac{9}{g^2 k^2} (k^2 + 1) j_0^2(k) \\ n_1(k) &= \frac{9}{g^2 k^2} j_1^2(k) \\ n_2(k) &= \frac{9}{g^2 k^2} j_2^2(k) \\ n_H(k) &= \frac{4}{g^2 k^2} (h^k)^2 : \end{aligned} \quad (15)$$

Fig. 4 shows the spectral densities for the gauge and Higgs particles, $n_W = n_0 + n_1 + n_2$ and n_H . The total particle numbers

$$N_W = \int_0^{Z_1} dk n_W(k) \quad \text{and} \quad N_H = \int_0^{Z_1} dk n_H(k) \quad (16)$$

and the energies E_W and E_H are listed in Table 1. The good agreement of $E_W + E_H$ with the energy gained from the fermions shows that the description in terms of the linearized theory is fully justified.

4. In summary, our work falls into two main parts, both of numerical character. First, we found the classical bounce trajectory between two topologically different sectors as a stationary point of the discretized Euclidean action on a grid in time and the radial space-coordinate. A careful treatment is required since this is a saddle point rather than a minimum, and numerical problems due to the gauge freedom must be handled properly. Second, we examine the real-time behavior of the fields after the tunneling, which gives some hints about the possible bosonic signature of baryon number violating processes. Our ansatz




Fig. 4: The spectral densities $n_W(k)$ and $n_H(k)$ (in units of m_W^{-1}) for various values of the chemical potential $\mu = \mu_{\text{crit}}$.

assumed a background of high fermionic density, but the results should also be interesting for the case of few particles with very high energy.

We are grateful to C Weiss for pointing our attention to the use of relaxation methods ([12]), and thank P Pobylitsa, M Polyakov, and V Petrov for numerous discussions. The work has been supported in part by the Deutsche Forschungsgemeinschaft, the RFBR grant 95-07-03662, and the joint Russian government-International Science Foundation grant R 2A 300.

References

- [1] G. 't Hooft, Phys. Rev. Lett. 37, 8 (1976).
- [2] T. Akiba, H. Kikuchi, and T. Yanagida, Phys. Rev. D 38, 1937 (1988).
- [3] V. Kuzmin, V. Rubakov, and M. Shaposhnikov, Phys. Lett. B 155, 36 (1985); B 191, 171 (1987).
- [4] P. Arnold and L. McLerran, Phys. Rev. D 36, 581 (1987); 37, 1020 (1988).
- [5] D. Diakonov, M. Polyakov, P. Sieber, J. Schaldach, and K. Goeke, Preprint RUB-TP II-25/95, hep-ph/9502245, submitted to Phys. Rev. D.
- [6] A. Ringwald, Nucl. Phys. B 330, 1 (1990).
- [7] O. Espinosa, Nucl. Phys. B 334, 310 (1990).
- [8] L. McLerran, A. Vainshtein, and M. B. Voloshin, Phys. Rev. D 42, 171 (1990).
- [9] V. Rubakov and A. Tavkhelidze, Phys. Lett. B 165, 109 (1985);
V. Rubakov, Prog. Theor. Phys. 75, 366 (1986);
V. Matveev, V. Rubakov, A. Tavkhelidze, and V. Tokarev, Nucl. Phys. B 282, 700 (1987).
- [10] D. Diakonov and V. Petrov, Phys. Lett. 275B, 459 (1992).
- [11] S. Coleman, Phys. Rev. D 15, 2929 (1977);
C. G. Callan, S. Coleman, Phys. Rev. D 16, 1762 (1977).
- [12] S. L. Adler and T. Piran, Rev. of Mod. Phys. 56, 1 (1984).
- [13] V. A. Rubakov, B. E. Stern, and P. G. Tinyakov, Phys. Lett. B 160, 292 (1985).
- [14] A. Kusenko, Phys. Lett. B 358, 51 (1995).
- [15] I. A. Aitchison, Nucl. Phys. B 191, 429 (1981).
- [16] M. Hellmuth and J. Kripfganz, Nucl. Phys. B 373, 749 (1991).
- [17] W. N. Cottingham and N. Hasan, Nucl. Phys. B 392, 39 (1993).
- [18] E. Farhi, J. Goldstone, S. Gutman, K. Rajagopal, and R. Singleton, Jr., Phys. Rev. D 51, 4561 (1995).