

CERN-TH/96-07
SCIPP 96/05
hep-ph/9601330

Recent Replacements in Higgs Physics

Howard E. Haber

CERN, TH-Division,
CH-1211 Geneva 23, Switzerland
and

Santa Cruz Institute for Particle Physics,
University of California, Santa Cruz, CA 95064 USA

Abstract

Recent replacements of the phenomenology of Higgs bosons in the Standard Model and the Minimal Supersymmetric Standard Model are reviewed.

Invited Talk at the
1995 International Europhysics Conference on High Energy Physics,
Brussels, Belgium, 27 July-2 August, 1995

CERN-TH/96-07
January, 1996

RECENT REFINEMENTS IN HIGGS PHYSICS

Howard E. Haber

CERN, TH-D ivision, CH-1211 Geneva 23, Switzerland
and

Santa Cruz Institute for Particle Physics, University of California, Santa Cruz, CA, 95064, USA

Recent refinements of the phenomenology of Higgs bosons in the Standard Model and the Minimal Supersymmetric Standard Model are reviewed.

1 Introduction

In this brief review, I will summarize a few of the advances in Higgs phenomenology during 1995. Space limitations allow me to present only a bare outline of the talk presented at the EPS meeting in Brussels, along with the attendant references. As a result, this review is necessarily incomplete, and I apologize in advance to authors of relevant work that has been omitted.

2 The Standard Model Higgs Boson is a Contradiction in Terms

Phenomenologists and experimentalists who plan the Higgs searches at future colliders spend much effort in designing a search for the Standard Model Higgs boson. However, the Standard Model Higgs boson is a meaningless term unless additional information is provided. This is because the Standard Model itself cannot be a fundamental theory of particle interactions. It must break down once the energy is raised beyond some critical scale. What is the value of ? Of course, this is unknown at present. can lie anywhere between a few hundred GeV and the Planck scale (M_{PL}).

Theorists who discuss the phenomenology of the Standard Model usually do not need to know the value of . At energy scales below , the physics beyond the Standard Model generally decouples, leaving a low-energy effective theory which looks almost exactly like the Standard Model. However, the Higgs boson presents a potential opportunity to probe . The stability of the Higgs potential places non-trivial constraints on the Higgs mass, due to the large value of the top quark mass. (More refined limits require only a metastable potential with a lifetime that is long compared to the age of the universe.) Recent computations of Refs. 1 and 2 show for example that if $m_t = 175 \text{ GeV}$, then for $m_t = 175 \text{ GeV}$ the Higgs mass must be larger than about 120 GeV.

Does this mean that if a Higgs boson mass of 100 GeV is discovered then the Standard Model Higgs boson is ruled out? The answer is yes, only if the phrase "the Standard Model Higgs boson" implies that

$= M_{\text{PL}}$. Formally, this is too narrow a definition. I would prefer to say that if a 100 GeV Higgs boson were discovered, then new physics beyond the Standard Model must enter at or below an energy scale of $\sim 1000 \text{ TeV}$ (based on the graphs presented in Ref. 2). Of course, in this case, if all the new physics were confined to lie in the vicinity of 1000 TeV, then LHC phenomenology would find no deviations from the Standard Model. Thus, physicists who plan searches for the Standard Model Higgs boson are not wasting their time. In particular, even if is rather close to the TeV scale, one would expect the lightest Higgs boson to retain all the properties of the so-called Standard Model Higgs boson.

To reiterate, the Standard Model Higgs boson is a sensible concept only if you specify the value of the energy scale at which the Standard Model breaks down. Now that we can all agree on the meaning of "the Standard Model Higgs boson", consider recent refinements in its phenomenology. Here, I would like to refer to two interesting directions.

First, if one assumes the validity of the Standard Model for low-energy physics (below the TeV scale), then one can test this theory by confronting it with the precision electroweak data. In addition to testing the Standard Model, one has the possibility of constraining the value of the Higgs mass, which enters through the radiative corrections to the Z and W boson self-energies. Combining them with the most recent LEP and SLC electroweak results³ with the recent top-quark mass measurement at the Tevatron,⁴ a weak preference is found^{3,5} for a light Higgs boson mass of order m_Z . One must take this result with a large grain of salt, since the overall chi-square of the Standard Model fit is not good (of order 2 per degree of freedom). Nevertheless, it does suggest the potential of future precision measurements for placing interesting constraints on the Standard Model Higgs mass.

Second, a number of two-loop computations of Higgs boson processes have recently been completed. Among them are an $O(\frac{3}{s})$ calculation of $h^0 \rightarrow gg$ and $O(\frac{2}{s}; \frac{2}{s} G_F m_t^2; G_F^2 m_t^4)$ terms in $h^0 \rightarrow b\bar{b}$. See Ref. 6 for details. Together with the recent computation of $K^0 \rightarrow \pi^0 \pi^0$ factors in $pp \rightarrow h^0 + X$,⁷ one now has both improved pro-

duction cross-section and branching ratio calculations, leading to more accurate Higgs boson phenomenology at both the LHC and future e^+e^- colliders.

3 The Radiatively-Corrected MSSM Higgs Mass

If the minimal supersymmetric extension of the Standard Model (MSSM) is correct, then we should identify the scale at which the Standard Model breaks down as the scale of low-energy supersymmetry breaking. In models of low-energy supersymmetry, it is presumed to lie between m_Z and about 1 TeV. The mass of the light CP-even neutral Higgs boson, h^0 , in the MSSM can be calculated to arbitrary accuracy in terms of two parameters of the Higgs sector, m_{A^0} and $\tan\beta$,⁸ and other MSSM soft-supersymmetry-breaking parameters that affect the Higgs mass through virtual loops.⁹ If the scale of supersymmetry breaking is much larger than m_Z , then large logarithmic terms arise in the perturbation expansion. These large logarithms can be resummed using renormalization group (RG) methods.

The formula for the full one-loop radiative corrected Higgs mass is very complicated.¹⁰ Moreover, the computation of the RG-improved one-loop corrections requires numerical integration of a coupled set of RG equations.¹¹ (The dominant two-loop next-to-leading logarithmic results are also known.¹²) Although this program has been carried out in the literature, the procedure is unwieldy and not easily amenable to large-scale Monte-Carlo analyses. Below, we summarize a very simple procedure for accurately approximating m_{h^0} . The method can be easily implemented, and incorporates both the leading one-loop and two-loop effects and the RG-improvement. Although the method is conceptually simple, complications arise when supersymmetric thresholds are fully taken into account. The details can be found in Ref. 13, along with other references to the original literature. Complementary work can be found in Ref. 14. In the limited space allotted here, only the simplest version of our method is outlined.

The dominant radiative corrections to m_{h^0} arise from an incomplete cancelation of the virtual top-quark and top-squark loops. The two top-squark masses ($M_{\tilde{e}_L}$ and $M_{\tilde{e}_R}$) are obtained by diagonalizing a 2×2 top-squark squared-mass matrix; the off-diagonal elements of this matrix are denoted by m_{tX_t} (where $X_t = A_t \cot\beta$). We also assume that $M_{\tilde{e}_L}, M_{\tilde{e}_R}, m_{A^0} \gg m_Z$. This case is particularly useful since the upper bound for m_{h^0} (at fixed $\tan\beta$) arises precisely in this limit. The leading terms in the one-loop approximation to $m_{h^0}^2$ are given by

$$m_{h^0}^2 = m_Z^2 \cos^2 2\beta + (m_{h^0}^2)_{1LL}(m_t) + (m_{h^0}^2)_{mix}(m_t); \quad (1)$$

where

$$(m_{h^0}^2)_{1LL} = \frac{3g_2^2 m_t^4}{8 m_W^2} \ln \frac{M_{\tilde{e}_L} M_{\tilde{e}_R}}{m_t^2} + O\left(\frac{m_W^2}{m_t^2}\right); \quad (2)$$

and

$$(m_{h^0}^2)_{mix} = \frac{3g_2^2 m_t^4 X_t^2}{16 m_W^2} 2h(M_{\tilde{e}_L}^2; M_{\tilde{e}_R}^2) + X_t^2 g(M_{\tilde{e}_L}^2; M_{\tilde{e}_R}^2) + O\left(\frac{m_W^2}{m_t^2}\right) \quad (3)$$

In eq. (3), the functions g and h are given by:

$$g(a;b) = \frac{1}{(a-b)^2} 2 \frac{a+b}{a-b} \ln \frac{a}{b}; \quad (4)$$

$$h(a;b) = \frac{1}{a-b} \ln \frac{a}{b}; \quad (5)$$

The notation used in eq. (1) emphasizes the m_t dependence of $(m_{h^0}^2)_{1LL}$ and $(m_{h^0}^2)_{mix}$. Subdominant terms not shown explicitly in eqs. (2) and (3) can be found in ref. 13. Among the next-to-leading two-loop corrections, the most important effect can be incorporated by replacing m_t in the formulae above by the running top quark mass evaluated at m_t . Explicitly, $m_t(m_t)$ is given in terms of the pole mass m_t^{pole} by

$$m_t(m_t) = m_t^{pole} \left[1 - \frac{4}{3} \frac{s}{t} + \frac{t}{2} \right] 0.966 m_t^{pole}; \quad (6)$$

where $t = h_t^2/4$ and the Higgs-top quark Yukawa coupling is $h_t = g m_t / 2m_W$. (Since by assumption $m_{A^0} \gg m_Z$, one should identify h_t as the Yukawa coupling of the low-energy effective one-doublet Higgs sector. For details appropriate to the case of $m_{A^0} \sim O(m_Z)$, see Ref. 13.) Due to the leading m_t^4 behavior of the one-loop corrections, the replacement of $m_t = m_t^{pole}$ in eqs. (2) and (3) by $m_t(m_t)$ is numerically important, leading to a significant reduction in the predicted value of m_{h^0} .

We now proceed to sum the leading logarithmic terms to all orders in perturbation theory via RG-improvement. As noted above, this requires extensive numerical analysis. Nevertheless, we have found a remarkably simple analytic formula that incorporates the dominant effects of the RG-improvement. We already noted that replacing the pole mass m_t^{pole} with the running mass $m_t(m_t)$ has the effect of including the dominant part of the $O(m_t^2/m_t^2)$ and $O(m_t^2/m_t^2 s)$ next-to-leading logarithmic contributions to $m_{h^0}^2$. We find that the two-loop leading logarithmic contributions to $m_{h^0}^2$ can be incorporated by replacing the running top quark mass with m_t evaluated at an appropriately chosen scale. For $M_{\tilde{e}_L} = M_{\tilde{e}_R} = M_{\tilde{t}}$, our result is:

$$m_{h^0}^2 = m_Z^2 \cos^2 2\beta + (m_{h^0}^2)_{1LL}(m_t(m_t)) + (m_{h^0}^2)_{mix}(m_t(M_{\tilde{t}})); \quad (7)$$




Figure 1: The radiatively corrected light CP-even Higgs mass is plotted as a function of A_t/M_S for $\tan \beta = 20$. The one-loop leading logarithmic computation is compared with the RG-improved result which was obtained by numerical analysis and by using the simple analytic result given in eq. (7). M_S characterizes the scale of supersymmetry breaking and can be regarded (approximately) as a common supersymmetric scalar mass.

where $m_t = \frac{P}{m_t M_t}$, and the running top-quark mass is given by

$$m_t(\mu) = m_t(m_t) \left[1 - \frac{s}{16} \ln \frac{\mu^2}{m_t^2} \right] : \quad (8)$$

All couplings on the right hand side of eq. (8) are evaluated at m_t . In our numerical work, we have verified that this prescription reproduces the full RG-improved Higgs mass to within 2 GeV for top-squark masses of 2 TeV or below. Figures 1 and 2 exhibit two graphs to support this claim. Further details can be found in Ref. 13.

4 Higgs Searches at Future Colliders

1995 was a year of re-assessment. In the United States, the Division of Particles and Fields published a Long Range Planning Study to assess the future of the US particle physics program in light of the demise of the SSC. One working group of this study focused on electroweak symmetry breaking and physics at the TeV scale. One of the tasks of this working group was to provide an in-depth survey of Higgs searches at future collider facilities. Their work can be found in Ref. 15. In Europe, 1995 was the year of the LEP-200 Study. The Higgs Boson Working group played a major role in making the case for upgrading the LEP collider to the highest possible energy. It now appears that LEP will be upgraded to a center-of-mass energy of 192 GeV during the next few years. This will permit the discovery of a Higgs boson up to around 95 GeV. This is very exciting for proponents




Figure 2: The radiatively corrected light CP-even Higgs mass is plotted as a function of M_S for $\tan \beta = 1.5$. For further details, see the caption to Figure 1.

of the MSSM if $\tan \beta$ is small [see Figure 2]. The LEP-200 Higgs Working group considered in detail the discovery reach of the upgraded LEP collider for the Standard Model Higgs boson and the MSSM Higgs boson (a few non-minimal approaches were also surveyed). The results of this work can be found in Ref. 16.

5 Conclusions

This past year has seen a number of theoretical and phenomenological refinements of Higgs boson properties. The most comprehensive assessments of the Higgs searches at future colliders have been presented. Meanwhile, we wait for experiments at LEP-2 and/or the LHC to shed light on the origin of the dynamics that is responsible for electroweak symmetry breaking.

Acknowledgments

The work of Section 3 is based on a collaboration with Andre Hoang and Ralf Hemping. This work was supported in part by the U.S. Department of Energy.

References

1. G. Altarelli and G. Isidori, Phys. Lett. B 337 (1994) 141.
2. J.A. Casas, J.R. Espinosa and M. Quiros, Phys. Lett. B 342 (1995) 171; J.R. Espinosa and M. Quiros, Phys. Lett. B 353 (1995) 257.

3. P. Antilogus et al. [LEP Electroweak Working Group], LEP EW G/95-02 (1995), contributions of the LEP Experiments to the 1995 International Eu-rophysics Conference on High Energy Physics, 27 July-2 August, 1995, Brussels, Belgium, and the 17th International Symposium on Lepton-Photon Interactions, 10-15 August, 1995, Beijing, China.
4. F. Abe et al. [CDF Collaboration], Phys. Rev. Lett. 74 (1995) 2626; S. Abachi et al. [D0 Collaboration], Phys. Rev. Lett. 74 (1995) 2632.
5. P. H. Chankowski and S. Pokorski, Phys. Lett. B 356 (1995) 307; MPI-Ph-95-39 [hep-ph/9509207]; J. Ellis, G. L. Fogli, and E. Lisi, CERN-TH/95-202 [hep-ph/9507424].
6. A. L. Kataev and V. T. Kim, Mod. Phys. Lett. A 9 (1994) 1309; B. Kniehl, Phys. Rep. 240 (1994) 211; S. A. Larin, T. van Ritbergen and J. A. M. Vermaseren, Phys. Lett. B 362 (1995) 134; B. Kniehl and M. Spira, DESY-95-041 [hep-ph/9505225]; L. Durand, B. A. Kniehl, K. R. Kieselmann, Phys. Rev. D 51 (1995) 5007; A. Djouadi, M. Spira, and P. Zerwas, DESY-95-210 [hep-ph/9511344].
7. M. Spira, A. Djouadi, D. Graudenz, and P. Zerwas, Nucl. Phys. B 453 (1995) 17.
8. See, e.g., chapter 4 of J. F. Gunion, H. E. Haber, G. L. Kane and S. Dawson, *The Higgs Hunter's Guide* (Addison-Wesley, Redwood City, CA, 1990).
9. H. E. Haber and R. Hemping, Phys. Rev. Lett. 66 (1991) 1815; Y. Okada, M. Yamaguchi and T. Yanagida, Prog. Theor. Phys. 85 (1991) 1; J. Ellis, G. Ridol and F. Zwimer, Phys. Lett. B 257 (1991) 83.
10. P. Chankowski, S. Pokorski, and J. Rosiek, Phys. Lett. B 274 (1992) 191; Nucl. Phys. B 423 (1994) 437; A. Dabelstein, Z. Phys. C 67 (1995) 495.
11. H. E. Haber and R. Hemping, Phys. Rev. D 48 (1993) 4280.
12. R. Hemping and A. H. Hoang, Phys. Lett. B 331 (1994) 99.
13. H. E. Haber, R. Hemping, and A. H. Hoang, CERN-TH/95-216 and TTP 95-09 (1995).
14. J. A. Casas, J. R. Espinosa, M. Quiros, A. Riotto, Nucl. Phys. B 436 (1995) 3 [E: B 439 (1995) 466]; M. Carena, J. R. Espinosa, M. Quiros, and C. E. M. Wagner, Phys. Lett. B 355 (1995) 209; M. Carena, M. Quiros, and C. E. M. Wagner, CERN-TH-95-157 (1995).
15. T. L. Barklow, S. Dawson, H. E. Haber and J. L. Siegrist, SLAC-PUB-95-6893 [hep-ph/9505296], in *Particle Physics (Perspectives and Opportunity, Report of the DPF Committee on Long Term Planning, edited by R. Peccei et al. (World Scientific, Singapore, 1995); J. F. Gunion, A. Stange, and S. Willenbrock, UCD-95-28 (1995), to appear in Electroweak Symmetry Breaking and New Physics at the TeV Scale, edited by T. Barklow, S. Dawson, H. E. Haber, and J. Siegrist (World Scientific, Singapore, to be published).*
16. G. Altarelli et al., "Interim Report on the Physics Motivations for an Energy Upgrade of LEP 2", CERN-TH/95-151 and CERN-PPE/95-78 (1995); M. Carena and P. Zerwas, conveners, *Higgs Physics, Report of the Higgs Boson Working Group of the LEP-200 Workshop.*