

OCHA-PP-73

January 1996

Electroweak baryogenesis with the vector-like quark model

Tomoko Uesugi

Department of Physics, Ochanomizu University, Tokyo 112, Japan

Abstract

We study on the electroweak baryogenesis problem in the vector-like quark model. This model can add to the minimal standard model the extra CP-violation source which breaks CP spontaneously. Using charge transport mechanism suggested by Nelson et al, we can get the net hypercharge flux : $f_Y \sim 10^{-5}$. This result could nicely explain the present baryon to entropy ratio : $\eta_B \sim 10^{-9}$.

1 Introduction

Recently, Nelson, Kaplan and Cohen suggested the electroweak baryogenesis scenario called the charge transport mechanism [1]. Using this mechanism, we study on the baryogenesis problem in the vector-like quark model. This model was first introduced by Bento, Branco and Parada [2] as one of the most simplest spontaneous CP-violating models. The vector-like quark model is attractive to the electroweak baryogenesis because it can not only add to the standard model an extra CP violation source, but also make the first order phase transition of the electroweak theory much stronger.

2 The vector-like quark model

We consider the extension of the standard model by adding extra SU(2) singlet quarks U as the fourth generation as well as an extra singlet Higgs S.

To appear in the Proceedings of the workshop on 'Electroweak Baryogenesis' held at Chikuhō on November 25 (1995). This talk is based on the collaboration with A. Sugamoto and A. Yamaguchi (Ochanomizu University).

The field content of this model is $(u d)_L^i, u_R^i, d_R^i, U_L, S, (i = 1; 2; 3; 4)$, where i and j are the generation indices, Φ denotes the standard Higgs doublet. We assume that the additional singlet Higgs S is a complex scalar and all the new fields introduced are odd under a Z_2 symmetry.

Under these conditions, the $SU(2) \times U(1) \times Z_2$ invariant Yukawa couplings are $L_Y = \frac{1}{2} (u d)_L^i (h_{ij} \tilde{\Phi}_R^j + f_{ij} u_R^j) + U \frac{1}{2} (f_{i4} S + f_{i4}^0 S^0) U_L u_R^i + h.c.$

During the phase transition, Φ and S are able to have vacuum expectation values as $\langle \Phi \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}$, $\langle S \rangle = \frac{v}{\sqrt{2}} e^{i\theta}$. We consider the most general renormalizable $SU(2) \times U(1) \times Z_2$ symmetric scalar potential. Which can be written as

$$V_S = \frac{1}{2} \mu^2 + \frac{1}{4} (\mu^2)^2 + S^\dagger S (a_1 + b_1 S^\dagger S) + (S^2 + S^{\dagger 2}) (a_2 + b_2 S^\dagger S) + b_3 (S^4 + S^{\dagger 4}) + [c_1 (S^2 + S^{\dagger 2}) + c_2 S^\dagger S].$$

In order to violate CP spontaneously, this potential must have the minimum value with the non-zero complex phase. The θ -dependent part of the potential has the minimum value at $\theta = \frac{1}{2} \cos^{-1} \frac{a_2 + \frac{b_2 v^2}{2} + \frac{c_1 v^2}{2}}{b_3 v^2}$, where v is space-dependent inside the bubble wall, so that θ must also have the space dependence inside the wall. The shape of the wall $v(z)$ can be obtained from the equation of motion of Φ , but here, we assume a wall profile simply as $v(z) = v_0 (1 + \tanh(z/z_w))$. We consider only the z -dependent solution, where the z axis is normal to the bubble wall, since we can get general solutions by boosting the wall profiles in the x - y plane. z_w is the parameter which determines the thickness of the wall.

3 Computation of the hypercharge flux

For simplicity, we treat only top and vector-like quarks to compute net hypercharge flux. What we have to do is only to solve the Dirac equation. The two-generations' up quark mass matrix is

$$M = \begin{pmatrix} f v & 0 \\ F V e^i & F^0 V e^i \end{pmatrix}$$

Using this mass matrix, we can write the Dirac equation in the chiral bases. For the stationary state $\Psi(t; z) = \Psi(z) e^{i E t}$, the Dirac equation can be written by

$$i \frac{\partial}{\partial z} \begin{pmatrix} \psi_{1R}(z) \\ \psi_{3L}(z) \end{pmatrix} = \begin{pmatrix} E & M^\dagger \\ M & E \end{pmatrix} \begin{pmatrix} \psi_{1R}(z) \\ \psi_{3L}(z) \end{pmatrix}$$

$$i \frac{\partial}{\partial z} \begin{pmatrix} \psi_{4L}(z) \\ \psi_{2R}(z) \end{pmatrix} = \begin{pmatrix} E & M^+ \\ M & E \end{pmatrix} \begin{pmatrix} \psi_{4L}(z) \\ \psi_{2R}(z) \end{pmatrix}$$

where $\psi_{L(R)}(z)$ are two component spinors, corresponding to the top and the vector-like quarks. We can solve this Dirac equation numerically as a simple system of ordinary differential equations with respect to z . Using the reflection coefficient from right-handed to left-handed, $R(E)$, and from left-handed to right-handed, $R(E)$, the hypercharge flux can be written by the integration over energy of the incident particles. For simplicity, we neglect the thermal distribution term, and the integral over transverse momentum k_T can be done analytically. Using this approximation for the hypercharge flux, and setting $\mu = 600$ [GeV], $m_t = 174$ [GeV], $m_H = 100$ [GeV] and $v_w = 0.5$ c, we can get the final result for the computation of the hypercharge flux, $f_Y \sim 10^{-5}$ [T³]. This value can be obtained by taking reasonable values for the various couplings in the scalar potential and for the mass of the singlet Higgs.

4 Summary

Using the charge transport mechanism, we could nicely explain present baryon to entropy ratio with the vector-like quark model. This model is able to add new source of CP violation as a spontaneous CP violation. In this case, the CP violating phase becomes space-dependent, and the hypercharge flux can be produced in the bubble wall. The produced hypercharge is finally converted to baryon number excess through the sphaleron transition as usual.

For details, see the forthcoming paper [3].

References

- [1] A. E. Nelson, D. B. Kaplan and A. G. Cohen, Nucl. Phys. B 373 (1992), 453; A. G. Cohen, D. B. Kaplan and A. E. Nelson, Ann. Rev. and Part. Sci., 43 (1993), 27.
- [2] L. Bento, G. Branco and P. A. Parada, Phys. Lett. B 267 (1991), 95.
- [3] T. Uesugi, A. Yamaguchi, A. Sugamoto
Electroweak baryogenesis with the vector-like quark model (in preparation).