

# Symmetry Breaking and Generational Mixing in Topcolor-Assisted Technicolor

Kenneth Lane<sup>Y</sup>

Department of Physics, Boston University  
590 Commonwealth Avenue, Boston, MA 02215

## Abstract

Topcolor-assisted technicolor provides a dynamical explanation for electroweak and flavor symmetry breaking and for the large mass of the top quark without unnatural fine tuning. A major challenge is to generate the observed mixing between heavy and light generations while breaking the strong topcolor interactions near 1 TeV. I argue that these phenomena, as well as electroweak symmetry breaking, are intimately connected and I present a scenario for them based on nontrivial patterns of technifermion condensation. I also exhibit a class of models realizing this scenario. This picture leads to a rich phenomenology, especially in hadron and lepton collider experiments in the few hundred GeV to few TeV region and in precision electroweak tests at the  $Z^0$ , atomic parity violation, and polarized Møller scattering.

## 1. Introduction

Topcolor-assisted technicolor (TC2) was proposed by Hill [1] to overcome major shortcomings of top-condensate models of electroweak symmetry breaking [2], [3] and of technicolor models of dynamical electroweak and flavor symmetry breaking [4], [5]. Technicolor and extended technicolor (ETC) have been unable to provide a natural and plausible understanding of why the top quark mass is so large [6]. On the other hand, models in which strong topcolor interactions drive top-quark condensation and electroweak symmetry breaking are unnatural. To reproduce the one-Higgs-doublet standard model consistent with precision electroweak measurements (especially of the parameter  $\rho = M_W^2/M_Z^2 \cos^2 \theta_W \approx 1$ ), the topcolor energy scale must be much greater than the electroweak scale of  $O(1 \text{ TeV})$ . This requires severe fine tuning of the topcolor coupling.

Hill's combination of topcolor and technicolor keeps the best of both schemes. In TC2, technicolor interactions at the scale  $\Lambda_{TC} \approx \Lambda_{EW} \approx 1 \text{ TeV}$  are mainly responsible for electroweak symmetry breaking. Extended technicolor is still required for the hard masses of all quarks and leptons except the top quark. Topcolor produces a large top condensate,  $\langle \bar{t}t \rangle$ , and all but a few GeV of  $m_t \approx 175 \text{ GeV}$ .<sup>1</sup> However, it contributes comparatively little to electroweak symmetry breaking. Thus, the topcolor scale can be lowered to near 1 TeV and the interaction requires little or no fine tuning.

In the simplest example of Hill's TC2, there are separate color and weak hypercharge gauge groups for the heavy third generation of quarks and leptons and for the two light generations. The third generation transforms under strongly-coupled  $SU(3)_1 \times U(1)_1$  with the usual charges, while the light generations transform conventionally under weakly-coupled  $SU(3)_2 \times U(1)_2$ . Near 1 TeV, these four groups are broken to the diagonal subgroup of ordinary color and hypercharge,  $SU(3)_C \times U(1)_Y$ . The desired pattern of condensation occurs because  $U(1)_1$  couplings are such that the spontaneously broken  $SU(3)_1 \times U(1)_1$  interactions are supercritical only for the top quark.

Two important constraints were imposed on TC2 soon after Hill's proposal was made. The first is due to Chivukula, Dobrescu and Terning (CDT) [7] who claimed that the technifermions required to break top and bottom quark chiral symmetries are likely to have custodial-isospin violating couplings to the strong  $U(1)_1$ . To keep  $\rho \approx 1$ , they

---

<sup>1</sup> A small part of  $m_t$  must be generated by ETC to give mass to the Goldstone bosons (top-pions) associated with top condensation. Hill has pointed out that some, perhaps all, of the bottom quark mass may arise from  $SU(3)_1$  instantons [1].

argued, the  $U(1)_1$  interaction must be so weak that it is necessary to tune the  $SU(3)_1$  coupling to within 1% of its critical value for top condensation and to increase the topcolor boson mass above 4.5 TeV. Thus, TC2 still seemed to be unnatural. CDT stated that their bounds could be relaxed if  $U(1)_1$  couplings did not violate isospin. However, they expected that this would be difficult to implement because of the requirements of canceling gauge anomalies and of allowing mixing between the third and first two generations.

The second constraint on TC2 is due to Komiris [8] who showed, presuming that the b{quark's topcolor interactions are not far from critical, the existence of relatively light scalar bound states of  $t_L b_R$  and  $b_L b_R$  that couple strongly ( $\sim m_t$ ) to third generation quarks. These scalars can induce excessive  $B_d - B_d$  mixing which is proportional to the product  $D_{Lbd}^d D_{Rbd}^d$  of the elements of the unitary matrices which diagonalize the (generally nonhermitian)  $Q = \frac{1}{3}$  quark mass matrix.

The question of isospin violation and naturalness raised by CDT was addressed in Ref. [9]. We proposed that different technifermion isodoublets,  $T^t$  and  $T^b$ , give ETC mass to the top and bottom quarks. These doublets then could have different  $U(1)_1$  charges which were, however, isospin conserving for the right as well as left-handed parts of each doublet.<sup>2</sup> In addition, we exhibited a TC2 prototype in which (i) all gauge anomalies cancel; (ii) there are no very light pseudo-Goldstone bosons (loosely speaking, "axions") because all spontaneously broken global technifermion symmetries are broken explicitly by ETC [10]; and (iii) a mechanism exists for mixing the heavy and light generations.

Although the problem of  $B_d - B_d$  mixing raised by Komiris was not considered in [9], the  $U(1)$  symmetries of the model presented there automatically allow just one of two ETC-induced transitions in the quark mass matrix:  $d_L; s_L \rightarrow b_R$  or  $d_R; s_R \rightarrow b_L$ . Thus, only  $D_{Lbd}^d$  or  $D_{Rbd}^d$ , respectively, can be sizable and the  $B_d - B_d$  constraint is satisfied. It is easy to see that the phenomenologically preferred transition is  $d_L; s_L \rightarrow b_R$ : The known mixings between the third and the first two generations are in the Kobayashi-Maskawa matrix for left-handed quarks,  $V = D_L^{uY} D_L^d$ . They are  $|V_{cb}| \sim |V_{ts}| \sim 0.03 \sim 0.05 \frac{m_s}{m_b}$  and  $|V_{ub}| \sim |V_{td}| \sim 0.002 \sim 0.015 \sin \theta_c \frac{m_s}{m_b}$  [11]. These elements must arise from  $D_L^d$ , hence from the  $d_L; s_L \rightarrow b_R$  transitions, because the corresponding elements in  $D_L^u$  are smaller by a factor of  $m_b/m_t \sim 0.03$ .

In the model of Ref. [9], the mechanism of topcolor breaking was left unspecified and all technifermions were taken to be  $SU(3)_1 - SU(3)_2$  singlets. Thus, the transition

---

<sup>2</sup> While this eliminates the large  $\theta$  discussed by CDT, there remain small,  $O(\theta)$ , contributions from the  $U(1)_1$  interaction.

$d_L; s_L \text{ \& } b_R$  had to be generated by an externally induced term  $M_{ETC}$  in the ETC mass matrix which transforms as  $(3;3)$  under the color groups. We then estimated

$$\mathcal{Y}_{cbj'} \mathcal{D}_{Lsbj'}^d \frac{m_{sb}}{m_b} < \frac{m_{sb}}{m_b^{ETC}} \cdot \frac{M_{ETC}^2}{M_s^2}; \quad (11)$$

where  $m_{sb}$  is the mixing term in the  $Q = \frac{1}{3}$  mass matrix,  $m_b$  is the mass of the  $b$  quark, and  $M_s$  is the mass of the ETC boson that generates the strange quark mass,  $m_s$ . In a walking technicolor theory [12],  $M_s > 100 \text{ TeV}$ . However, we expect  $M_{ETC} = O(1 \text{ TeV})$  because that is the scale at which topcolor breaking naturally occurs. This gives  $s$  mixing that is at least 300 times too small. We stated in [9] that providing mixing of the observed size between the heavy and light generations is one of the great challenges to topcolor-assisted technicolor.

This problem is addressed in the rest of this paper. I shall argue that generational mixing is intimately connected to topcolor and electroweak symmetry breaking and that all these phenomena occur through technifermion condensation. In Sections 2-4, I specify the gauge groups and describe the patterns of gauge symmetry breaking needed for standard model phenomenology. Nontrivial patterns of vacuum alignment play a central role in this. In Section 5, I present a class of models which illustrate this scenario. The phenomenology of these models is sketched in Section 6. Special attention is placed on the  $Z^0$  boson of the broken  $U(1)_1$  symmetry. Its effects may be noticeable in hadron collider production of jets and dileptons,  $e^+e^-$  collisions, atomic parity violation, polarized Moller scattering and other precision electroweak measurements. I also emphasize observational consequences of vacuum alignment, especially technirho vector mesons and their decay to pairs of technipions and, possibly, CP violation.

## 2. Gauge Groups

The gauge groups of immediate interest to us are

$$SU(N) \times SU(3)_1 \times SU(3)_2 \times U(1)_1 \times U(1)_2 \times SU(2); \quad (2.1)$$

where, for definiteness, I have assumed that the technicolor gauge group is  $SU(N)$ . To avoid light "axions", all of these groups (except for the electroweak  $SU(2)$  and, possibly, parts of the  $U(1)$ 's) must be embedded in an extended technicolor group,  $G_{ETC}$ . I will not specify  $G_{ETC}$ . This difficult problem is reserved for the future. However, as in Ref. [9],

I shall assume the existence of ETC induced four fermion operators which are needed to break quark, lepton and technifermion chiral symmetries. Of course, these operators must be invariant under the groups in Eq. (2.1).

The coupling constants of  $SU(3)_1 \times SU(3)_2 \times U(1)_1 \times U(1)_2$  are denoted by  $g_1, g_2, g_1^0, g_2^0$ , where  $g_1 = g_2$  and  $g_1^0 = g_2^0$ . When these gauge symmetries break,  $SU(3)_1 \times SU(3)_2 \rightarrow SU(3)_C$  and  $U(1)_1 \times U(1)_2 \rightarrow U(1)_Y$ . We shall see that the breaking to  $U(1)_Y$  must occur at an energy higher than the  $SU(2) \times U(1)_K$  breaking scale  $\Lambda_{EW}$ . Then, the usual color and weak hypercharge couplings are

$$g_s = \sqrt{\frac{g_1 g_2}{g_1^2 + g_2^2}}, \quad g_2; \quad g^0 = \sqrt{\frac{g_1^0 g_2^0}{g_1^{02} + g_2^{02}}}, \quad g_2^0: \quad (2.2)$$

These symmetry breakings give rise to eight color octet ( $V_8$ ) vector bosons and one neutral  $Z^0$ , all of which have mass of  $O(1 \text{ TeV})$  [13], [1].

Third generation quarks  $q^3 = (t; b)$  will transform as  $(3; 1)$  under  $SU(3)_1 \times SU(3)_2$ , while the first two generation quarks  $q^1 = (u; d), (c; s)$  transform as  $(1; 3)$ . Unlike the situation in the simple models of Refs. [1] and [9], we shall find it necessary to assume that all quarks and leptons carry both  $U(1)_1$  and  $U(1)_2$  charges. These hypercharge assignments must be such that the gauge interactions are supercritical only for the top quark. This new situation has important phenomenological consequences, outlined in Section 6.

### 3. $U(1)_1 \times U(1)_2$ Breaking

In the scenario I describe, the extra  $Z^0$  resulting from  $U(1)_1 \times U(1)_2$  breaking has a mass of at most a few TeV and couples strongly to light, as well as heavy, quarks and leptons. Then, two conditions are necessary to prevent conflict with neutral current experiments. First, there must be a  $Z^0$  boson with standard electroweak couplings to all quarks and leptons. To arrange this, there will be a hierarchy of symmetry breaking scales, with  $U(1)_1 \times U(1)_2 \rightarrow U(1)_Y$  at  $1\{2 \text{ TeV}$ , followed by  $SU(2) \times U(1)_K \rightarrow U(1)_{EM}$  at the lower scale  $\Lambda_{EW}$ . Assuming that technicolor interactions induce both symmetry breakdowns, the technifermions responsible for  $U(1)_1 \times U(1)_2 \rightarrow U(1)_Y$  call them  $L$  and  $R$  must belong to a vectorial representation of  $SU(2)$ . To simplify the analysis, I make the minimal assumption that the  $L, R$  are electrically neutral  $SU(2)$  singlets.

To produce this hierarchy of symmetry breaking scales, and yet maintain an asymptotically free technicolor, the  $L, R$  should belong to a higher dimensional representation of

$SU(N)$ , while the technifermions responsible for  $SU(2) \times U(1)_Y$  breaking must belong to fundamental representations. This is reminiscent of multiscale technicolor [14], but there both the higher and fundamental representations participate in electroweak symmetry breaking. In the present model, I shall assume that  $\psi_{L,R}$  belong to the  $\frac{1}{2}N(N-1)$  dimensional antisymmetric tensor representation. I assume that this set of technifermions is large enough to ensure that the technicolor coupling "walks" for a large range of momenta [12].

The second constraint is that the  $Z^0$  should not induce large flavor-changing interactions. This can be achieved if the  $U(1)_1$  couplings of the two light generations are GIM symmetric. Then flavor-changing effects will nominally be of order  $|V_{ub}|^2 \approx M_{Z^0}^{-2}$  for  $B_d = 2$  processes,  $|V_{cb}|^2 \approx M_{Z^0}^{-2}$  for  $B_s = 2$ , and negligibly small for  $S = 2$ . These should be within experimental limits.<sup>3</sup> Nevertheless, a variety of interesting, and potentially dangerous,  $Z^0$  phenomena are expected. These are discussed in Section 6.

#### 4. $SU(3)_1 \times SU(3)_2$ and Electroweak Breaking and Generational Mixing

Turn now to symmetry breaking at lower energy scales. I recounted above that sfb mixing is too small by a factor of 300 if  $SU(3)_1 \times SU(3)_2$  breaking is introduced to the quark sector only by a mixing term in the ETC boson mass matrix. Since  $b_R$  transforms as  $(3; 1; 1; \frac{1}{3})$  under  $SU(3)_1 \times SU(3)_2 \times SU(2) \times U(1)_Y$  and  $d_L; s_L$  as  $(1; 3; 2; \frac{1}{6})$ , it is tempting to suppose that the mechanism connecting  $d_L; s_L$  to  $b_R$  is at the same time responsible for breaking  $SU(3)_1 \times SU(3)_2 \rightarrow SU(3)_C$  and  $SU(2) \times U(1)_Y \rightarrow U(1)_{EM}$ . The generational mixing term transforms as  $(3; 3)$  under the color groups. Therefore, I introduce colored technifermion isodoublets transforming under  $SU(N) \times SU(3)_1 \times SU(3)_2 \times SU(2)$  as follows:

$$\begin{aligned} T_{L(R)}^1 &= \begin{pmatrix} U^1 \\ D^1 \end{pmatrix}_{L(R)} \quad 2(N; 3; 1; 2(1)) \\ T_{L(R)}^2 &= \begin{pmatrix} U^2 \\ D^2 \end{pmatrix}_{L(R)} \quad 2(N; 1; 3; 2(1)) : \end{aligned} \quad (4.1)$$

The transition  $d_L; s_L \rightarrow D_L^2 \rightarrow D_R^1 \rightarrow b_R$  occurs if the appropriate ETC operator exists and if the condensate  $\langle \bar{H}_L^1 T_R^2 \rangle$  forms.

---

<sup>3</sup> The most stringent constraint may come from  $M_{B_d} \approx M_{B_s}$ . In the model of Section 5, this ratio depends in a complicated way on the  $U(1)_1$  hypercharges  $b, b^0, d, d^0$  and the magnitudes and phases of  $V_{ub}$  and  $V_{td}$ .

The patterns of condensation,  $\langle \bar{H}_L^i T_R^j \rangle$ , that occur depend on the strength of the interactions driving them and on explicit chiral symmetry breaking (4T) interactions that determine the correct chiral (perturbative) ground state, i.e., "align the vacuum" [15]. The strong interactions driving technifermion condensation are  $SU(N)$ ,  $SU(3)_1$  and  $U(1)_1$ . The technicolor interactions do not prefer any particular form for  $\langle \bar{H}_L^i T_R^j \rangle$ ;  $SU(3)_1$  drives  $\langle \bar{H}_L^1 T_R^1 \rangle \neq 0$ ;  $U(1)_1$  drives  $\langle \bar{H}_L^1 T_R^1 \rangle$ ;  $\langle \bar{H}_L^2 T_R^2 \rangle \neq 0$  or  $\langle \bar{H}_L^1 T_R^2 \rangle \neq 0$ , depending on the strong hypercharge assignments.

In the approximation that technicolor interactions dominate condensate formation, so that

$$\langle \bar{H}_L^i T_R^j \rangle = \frac{1}{2} \text{Tr} U_{ij} \quad (i; j = 1; 2); \quad (4.2)$$

it is easy to prove the following: If  $T^1 \sim (3; 1)$  and  $T^2 \sim (1; 3)$  are the only technifermions and if the vacuum (aligning) interactions are  $SU(3)_1 \times SU(3)_2$  symmetric then, in each charge sector, the unitary matrix  $U_{ij} = \delta_{ij}$  or  $U_{ij} = (\delta_{ij})_{ij}$ , but not a nontrivial combination of the two. Therefore, in order that  $SU(3)_1 \times SU(3)_2$  invariant direct mass terms,  $d_L; s_L \times d_R; s_R$  and  $b_L \times b_R$ , occur as well as the mixing  $d_L; s_L \times b_R$ , it is necessary to introduce still other technifermions. The least number of additional technifermions involves  $SU(3)_1 \times SU(3)_2$  singlets. In the model described below, these will consist of three isodoublets:  $T^1$  giving direct mass terms to the light quarks and leptons;  $T^t$  giving the top quark its ETC mass; and  $T^b$  giving the bottom quark its ETC mass. These are the same technifermions used in the model of Ref. [9]. Introducing them enlarges the chiral symmetry and the number of Goldstone bosons of the model. Giving mass to all these bosons will require, among other things, a nontrivial pattern of  $T^1 \times T^2$  condensation,  $U = a_0 1 + ia_2 \cdot 2$ . This simultaneously breaks the color and electroweak symmetries to  $SU(3)_C \times U(1)_{EM}$  and provides large generational mixing, e.g.,  $m_{sb} = \langle \bar{H}_L^1 T^2 \rangle_{M_s} = M_s^2 m_s$ . The color singlet technifermions help align the vacuum in this nontrivial way as well as contribute to electroweak symmetry breaking.

## 5. A Model

In this section I follow the format of Ref. [9] to construct a TC2 model with the symmetry breaking just outlined. First, I list hypercharge assignments for all the fermions and explain certain general constraints on them. Then I derive a condition on the hypercharges that must be satisfied in order that colored technifermions condense to break topcolor  $SU(3)$ . I conclude by discussing other conditions available to fix the hypercharges.

Among these are the gauge anomaly constraints, given in the Appendix. The rest follow from specifying the ETC four-fermion operators necessary to give masses to quarks and leptons and to the Goldstone bosons associated with global symmetries. A family of solutions for the hypercharges satisfying all these constraints is obtained in the Appendix.

The fermions in the model, their color representations and  $U(1)$  charges are listed in Table 1. A number of choices have been made at the outset to limit and simplify the charges and to achieve the scenario's objectives:

1. In order that technifermion condensates conserve electric charge,  $u_1 + u_2 = v_1 + v_2$ ,  $x_1 + x_2 = x_1^0 + x_2^0$ ,  $y_1 + y_2 = y_1^0 + y_2^0$ , and  $z_1 + z_2 = z_1^0 + z_2^0$ .
2. The  $U(1)_1$  charges of technifermions respect custodial isospin.
3. The most important choice for our scenario is that of the  $U(1)_1$  charges of  $T^1$  and  $T^2$ . So long as  $u_1 \neq v_1$ , the broken  $U(1)_1$  interactions favor condensation of  $T^1$  with  $T^2$ . If this interaction is stronger than the  $SU(3)_1$  attraction for  $T^1$  with itself and if we neglect other vacuum-aligning ETC interactions, then  $h\Gamma_L^i T_R^j i / (i_2)_{ij}$  in each charge sector. This alignment is discussed below.
4. We shall see that  $u_1 \neq v_1$  implies  $Y_{1i} \neq Y_{1i}^0$  for the various fermions. Purely for simplicity, I have chosen  $Y_1 = b^0$  for all right-handed light quarks. I must choose  $Y_1(t_R) \neq Y_1(b_R)$  to prevent strong  $b$ -condensation. Again for simplicity, I put  $Y_1(t_R) = Y_1(b_R) = d^0$ . We shall see that  $dd^0$  is positive, as it must be for  $t$ -condensation.
5. For the  $SU(N)$  antisymmetric tensor  $\chi$ ,  $\chi^0 \neq 0$  guarantees  $U(1)_1 \times U(1)_2 \times U(1)_Y$  when  $h_L$  forms. Note that, if  $N = 4$ , a single real  $\chi_L$  is sufficient to break the  $U(1)$ 's. Otherwise, to limit the parameters,  $\chi^0 = 0$  may be assumed.

I now show that, in the absence of other ETC operators, the  $U(1)_1$  interactions can overwhelm  $SU(3)_1$  to produce the alignment pattern  $h\Gamma_L^i T_R^j i / (i_2)_{ij}$ . The coupling of the  $Z^0$  boson to a generic fermion with weak hypercharge  $Y = Y_1 + Y_2$  and electric charge  $Q = Y_1^0 + Y_2^0$  is

$$L_{Z^0} = g_{Z^0} Z^0 [(Y_1 - rY)_L - (Y_1^0 - rQ)_R]; \quad (5.1)$$

where  $g_{Z^0} = \frac{P}{g_1^2 + g_2^2}$ ,  $g_1^0$  and  $r = g_2^0/g_1^0 = 1$ . Small mixing terms induced by electroweak symmetry breaking have been neglected in this expression. A similar interaction can be written for the massive  $V_8$  bosons of broken  $SU(3)_1 \times SU(3)_2$ . Ignoring small terms

in the  $Z^0$  and  $V_8$  couplings, the four-fermion interaction these bosons generate for  $T^1$  and  $T^2$  is

$$L_{T^1 T^2} = \frac{2}{M_{Z^0}^2} u_1 T_L^1 T_L^1 + T_R^2 T_R^2 + v_1 T_R^1 T_R^1 + T_L^2 T_L^2 + \frac{v_8}{M_{V_8}^2} T_L^1 t_a T_L^1 + T_R^1 t_a T_R^1 \quad ; \quad (5.2)$$

where  $Z^0 = g_{Z^0}^2 = 4$  and the  $t_a$  are SU(3) matrices in the 3-representation. All the currents are SU(N) SU(2) singlets, and the current-current products are renormalized at the corresponding massive boson masses. Fierzing this interaction and retaining only the dominant SU(3) SU(N) SU(2) singlet operators involved in condensate formation gives

$$L_{T^1 T^2} = \frac{4}{3N M_{V_8}^2} \frac{u_1 v_1 Z^0 M_{V_8}^2}{M_{Z^0}^2} T_L^1 T_R^1 T_R^1 T_L^1 + T_L^2 T_R^2 T_R^2 T_L^2 + \frac{Z^0 M_{V_8}^2}{M_{Z^0}^2} u_1^2 T_L^1 T_R^2 T_R^2 T_L^1 + v_1^2 T_L^2 T_R^1 T_R^1 T_L^2 + \frac{4 v_8}{3} T_L^1 T_R^1 T_R^1 T_L^1 \quad ; \quad (5.3)$$

To determine which of the operators in Eq. (5.3) is dominant, I make the large(N) approximation that the anomalous dimensions of the 4T operators are given by the sum of the anomalous dimensions  $\gamma_{m_{ij}}$  of their constituent bilinears  $T^i T^j$ . Then, the condition that the vacuum energy  $E = \langle L_{T^1 T^2} \rangle$  is minimized by  $\langle T_L^i T_R^j \rangle / \langle (T^2)_{ij} \rangle$  is

$$\frac{Z^0 (u_1^2 + v_1^2) M_{V_8}^2}{M_{Z^0}^2} \frac{Z_{12}^2(M_{Z^0})}{Z_{11}^2(M_{V_8})} > \frac{4 v_8}{3} + \frac{u_1 v_1 Z^0 M_{V_8}^2}{M_{Z^0}^2} \frac{Z_{11}^2(M_{Z^0}) + Z_{22}^2(M_{Z^0})}{Z_{11}^2(M_{V_8})} \quad ; \quad (5.4)$$

where

$$Z_{ij}(M) = \exp \left[ \sum_{TC} \gamma_{m_{ij}} \right] \quad ; \quad (5.5)$$

Since the U(1) symmetries are broken at a higher scale than the SU(3) and electroweak symmetries,  $M_{Z^0}$  may be several times larger than  $M_{V_8}$ . However, the energy range from  $M_{V_8}$  to  $M_{Z^0}$  overlaps the region in which T condensates form. Thus, the anomalous dimensions  $\gamma_{m_{ij}} \neq 1$  there [12]. In this limit, the condition (5.4) becomes  $(u_1 + v_1)^2 > 4 v_8 = 3 Z^0$ .

The rest of my discussion of this model concerns how the  $U(1)_1$  and  $U(1)_2$  hypercharges are to be fixed. I start with the gauge anomaly conditions. The eight independent conditions are given in the Appendix. These constraints, together with the 4 equal{charge conditions, do not fix the 26 unknown  $U(1)_i$  charges. Further limitations on the  $Y_i$  follow from requiring the presence of ETC {generated four{ fermion operators breaking all but gauged symmetries. To give mass to quarks and leptons, I assume the following ETC operators:

$$\begin{aligned}
q_{iL}^1 & T_L^1 D_R^1 e_{jR} & =& a & a^0 = x_1 & x_1^0 \\
q_{iL}^1 & T_L^1 T_R^1 q_{jR}^1 & =& b & b^0 = x_1 & x_1^0 \\
q_L^h & T_L^1 D_R^1 & =& c & c^0 = x_1 & x_1^0 \\
q_L^h & T_L^t U_R^t t_R & =& d & d^0 = Y_1 & Y_1^0 \\
q_L^h & T_L^b D_R^b b_R & =& d + d^0 = z_1 & z_1^0 & :
\end{aligned} \tag{5:6}$$

To generate  $d_L ; s_L \text{ } b_R$ , I require the operator

$$q_{iL}^1 T_L^2 D_R^1 b_R =) \quad b + d^0 = 0 : \tag{5:7}$$

To forbid  $d_R ; s_R \text{ } b_L$ , ETC interactions must not generate the operator  $q_L^h T_L^1 D_R^2 d_{iR}$ . This gives the constraint

$$d \quad b^0 \notin 0 : \tag{5:8}$$

We shall see that this follows from requiring the existence of other four{ fermion operators and also the anomaly constraints. Thus, this operator does not appear without the intervention of  $U(1)_1$  breaking and so the transition  $d_R ; s_R \text{ } b_L$  is automatically suppressed relative to  $d_L ; s_L \text{ } b_R$  by a factor of  $M_{ETC}^2 / M_s^2 = O(10^{-4})$ .

Next, I enumerate the chiral symmetries and Goldstone bosons of the model, to determine what 4T operators are needed to give them mass. The simplest way to do this is to imagine that all gauge interactions, including  $SU(3)_1 \text{ } SU(3)_2 \text{ } U(1)_i$ , may be neglected compared to technicolor. Then, grouping the technifermions into three triplet{isodoublets,  $T^1 ; T^2$  and  $T^3 = T^1 ; T^t ; T^b$ , the chiral symmetry group of these technifermions plus  $L ; R$  is

$$G = SU(18)_L \text{ } SU(18)_R \text{ } U(1)_A : \tag{5:9}$$

The  $U(1)_A$  current involves all technifermions and has no technicolor anomaly. It is spontaneously broken principally by  $h_{LR}$ . A linear combination of this current and generators of  $SU(18)_A$  is exactly conserved and couples to the Goldstone boson eaten by the  $Z^0$ . The

orthogonal Goldstone boson gets mass from  $SU(3)_1$  instantons and broken ETC interactions. We need not be further concerned with  $U(1)_A$ .

When T condensates break  $SU(18)_L \times SU(18)_R$  to an  $SU(18)$  subgroup, there are 323 Goldstone bosons or technipions,  $\pi_T$ .<sup>4</sup> These may be conveniently classified according to the subgroup

$$H = SU(3)_1 \times SU(3)_2 \times SU(3)_3 \times SU(2) \times U(1)_3 \times U(1)_8; \quad (5:10)$$

where  $SU(3)_i$  acts on the triplet  $T^i$ ,  $SU(2)$  acts on the isodoublets within the triplets, and  $U(1)_{3,8}$  are generated by the diagonal charges of the  $SU(3)$  defined on the triplet  $T^1; T^2; T^3$ :

$$T^1 = \frac{1}{2} (3; 1; 1; 2; \frac{1}{2}; \frac{1}{12}); \quad T^2 = \frac{1}{2} (1; 3; 1; 2; \frac{1}{2}; \frac{1}{12}); \quad T^3 = \frac{1}{3} (1; 1; 3; 2; 0; \frac{1}{3}); \quad (5:11)$$

The 323 Goldstone bosons consist of: three  $SU(3)$  singlet isotriplets,  $(1; 1; 1; 3)$ ; three octet isotriplets plus three octet isosinglets; two singlets,  $(1; 1; 1; 1)$ ; and three sets of  $(3; 3) \times (3; 3)$  isotriplets and isosinglets.

The diagonal linear combination of the three  $(1; 1; 1; 3)$ 's become  $W_L$  and  $Z_L^0$ . Thus, ignoring the effects of color interactions, the decay constant of the technipions is  $F_T = 246 \text{ GeV} = \frac{P}{9} = 82 \text{ GeV}$ .<sup>5</sup> A linear combination of the  $(8; 1; 1; 1)$  and  $(1; 8; 1; 1)$  are absorbed in  $SU(3)_1 \times SU(3)_2 \times SU(3)_3$ , driven by  $\text{tr}(T_L^1 T_R^2)$ . Of the remaining 312 Goldstone bosons, all those which are  $SU(3)_1 \times SU(3)_2$  nonsinglets (there are 272 of these) acquire mass of at least  $\frac{P}{(s_T^4 = F_T^2)}$ ,  $250 \text{ GeV}$  from color interactions (see the papers by Peskin and Preskill in Ref. [15]).

<sup>4</sup> I do not know whether this is a record number of Goldstone bosons, as has been speculated. It certainly is a matter of concern whether they may make a large positive contribution to the S parameter. This is the case if they may be approximated as pseudo-Goldstone bosons [16]. As I have discussed elsewhere [17], this may be a poor approximation for the technipions in a walking technicolor model with its large anomalous dimensions. Furthermore, in such a model, there are additional, possibly negative, contributions to S which cannot be evaluated simply by scaling from QCD (see also Ref. [18]).

<sup>5</sup> I am suppressing the role of the  $SU(2) \times U(1)$  chiral symmetry of  $(t; b)_L$  and  $t_R$  in this discussion. The three Goldstone top-pions,  $\pi_t$ , arising from its breakdown combine with the  $(1; 1; 1; 3)$ 's to form the longitudinal weak bosons. In our normalization, Hill's estimate of the top-pion decay constant is  $F_t \approx 35 \text{ GeV}$  [1]. The uneaten component of the top-pions acquires its mass from the ETC part of the top quark mass:  $M_t^2 \approx m_t^{ETC} \text{tr}(t) = F_t^2$ .

This leaves 40 technipions whose mass must arise from ETC (generated 4T interactions). They transform as  $(1;1;8;3)$   $(1;1;8;1)$   $(1;1;1;3)$   $(1;1;1;3)$   $(1;1;1;1)$   $(1;1;1;1)$ . Consider the two isotriplets  $(1;1;1;3)$  orthogonal to the longitudinal weak bosons. It is possible to form one linear combination of these states that contains no  $T^i T^i$  component for one of the values of  $i = 1;2;3$ . Therefore, there must be a 4T term involving two technifermions of the form  $T_L^i T_L^j T_R^j T_R^i$ , with  $i \neq j$ , to insure that both isotriplets get mass. The only operators consistent with  $u_1 = v_1 \neq 0$  have  $i = 1$  or  $2$  and  $j = 3$ , with  $T^3 = T^1$  or  $T^t$  or  $T^b$ . Finally, in order that such an interaction contribute to  $(1;1;1;3)$  technipion masses, it is necessary that the condensates  $\langle T_L^1 T_R^1 \rangle$  and  $\langle T_L^2 T_R^2 \rangle$  form, i.e., that the matrix  $U$  in Eq. (4.2) is a nontrivial combination of  $1$  and  $i_2$ . Any of these 4T operators, in concert with  $SU(N) \times SU(3) \times U(1)$  interactions, can lead to such a pattern of vacuum alignment. As a specific choice consistent with Eqs. (5.6) and (5.7), I assume the existence of the operator

$$T_L^1 T_L^t T_R^t (a + b_3) T_R^1 =) \quad y_1 \quad Y_1^0 = u_1 \quad v_1 : \quad (5.12)$$

Equations (5.6), (5.7), (5.12) and the anomaly conditions for  $U(1)_{1;2} [SU(N)]^2$  and  $U(1)_{1;2} [SU(3)_{1;2}]^2$  lead to the relations:

$$\begin{aligned} a \quad d^0 &= b \quad b^0 = c \quad c^0 = x_1 \quad x_1^0 = \frac{1}{2}N (u_1 \quad v_1) \\ d \quad d^0 &= y_1 \quad Y_1^0 = u_1 \quad v_1 \\ d + d^0 &= z_1 \quad Z_1^0 = (2N + 1)(u_1 \quad v_1) \\ d &= N (u_1 \quad v_1) \\ d^0 &= b = (N + 1)(u_1 \quad v_1) \\ b^0 &= \frac{1}{2}(N + 2)(u_1 \quad v_1) \\ (N - 2)(\quad^0) &= 3N (u_1 \quad v_1) : \end{aligned} \quad (5.13)$$

We see that the constraint  $d = b \neq 0$  forbidding  $d_R; s_R \neq s_L$  is satisfied. Also,  $dd^0 > 0$ , just what is needed for top, but not bottom, quarks to condense. The condition  $\quad^0 \neq 0$  that  $\langle T_L T_R \rangle$  breaks  $U(1)_1 \times U(1)_2 \rightarrow U(1)_Y$  is equivalent to  $u_1 = v_1 \neq 0$ , necessary for  $\langle T_L^1 T_R^2 \rangle \neq 0$ .

Finally, there are  $(1;1;8;3)$ ,  $(1;1;8;1)$  and  $(1;1;1;1)$  technipions composed of  $T^1$  and  $T^b$  that do not acquire mass from the operator in Eq. (5.12). Combinations of spontaneously broken currents such as  $T^2 - T^2$   $3T^b - 5T^b$  are also left conserved by this

operator. Thus, we need a 4T operator involving both  $T^1$  and  $T^b$ . One choice (of several) that is consistent with all the operators assumed so far is

$$T_L^1 \quad T_L^t \quad T_R^b \quad (a + b_3) T_R^1 \quad (=) \quad Y_1 \quad Z_1^0 = Z_2^0 \quad Y_2 = X_1 \quad X_1^0 = \frac{1}{2} N (u_1 \quad v_1) : \quad (5.14)$$

Note that  $T^t$  and  $T^b$  must have the same electric charges, i.e.,  $Y_1 + Y_2 = Z_1 + Z_2$ .

We now have 18 linear plus 3 nonlinear conditions on the 26 hypercharges. In the Appendix, I exhibit solutions to these equations for which  $\sum_j u_j = \sum_j v_j = 0 \pmod{1}$ . The vacuum alignment program, including determination of the eigenvalues and eigenstates of the technipion mass matrix, is outlined in Section 6 and then deferred to a later paper.

## 6. The Phenomenology of Topcolor-Assisted Technicolor

The picture of topcolor-assisted technicolor I have drawn in this paper leads to a wide variety of phenomena in the TeV energy region, many of which are likely to be accessible in Tevatron collider experiments and, possibly, in LEP 2 experiments. Here is a list of the more obvious issues:

1. The  $Z^0$  boson, with  $M_{Z^0} = 1.3 \text{ TeV}$ .
2. The  $V_8$  colorons, with mass  $M_{V_8} < 1 \text{ TeV}$ . Their phenomenology was discussed in Refs. [13] and [19].
3. The quantum numbers, masses, and production and decay modes of techninos, technipions and top-pions.
4. A possible outcome of vacuum alignment is the appearance of CP violating phases in the unitary matrices defining mass eigenstate quarks (see Eichten, Lane and Preskill in Ref. [15]).
5. Cosmological consequences of the fermion which, apparently, must have a component that is stable against weak decay.
6. Since  $\sum_j u_j = \sum_j v_j$  must be 0 mod 1, some of the hypercharges in Eq. (5.13) are 0 mod N. This raises the question of the triviality of the  $U(1)_1$  interaction: does it set in at an energy much lower than the one at which we can envisage  $U(1)_1$  being unified into an asymptotically free ETC group?

Each of these topics requires extensive study. Here, I briefly discuss only the  $Z^0$  and the aspects of vacuum alignment. Details are under investigation by others or postponed to later papers

## Z<sup>0</sup> Physics

The mass of the Z<sup>0</sup> arises mainly from (condensation,

$$M_{Z^0} \approx g_{Z^0} j^0 F ; \quad (6:1)$$

where  $j^0 = 3N(u_1 - v_1) = N(2) = O(1)$ , and  $F = O(1 \text{ TeV})$  is the decay constant. This is the basis of my estimate of  $M_{Z^0}$ . The Z<sup>0</sup> decays into technifermion, quark and lepton pairs, with large couplings to all. Thus, its width is large, probably several hundred GeV [13]. I emphasize that in this scenario the Z<sup>0</sup> necessarily couples strongly to the first two generations of quarks and leptons.

There are several precision electroweak studies that probe for the Z<sup>0</sup> [20]. Mixing of the Z<sup>0</sup> and Z<sup>0</sup> affects the latter's couplings to quark and lepton pairs. If the Z<sup>0</sup> width is not an issue, the magnitude of these mixing effects is

$$\frac{\delta g_{Z^0}}{g_{Z^0}} \approx \frac{g_{Z^0} M_Z^2}{g_Z M_{Z^0}^2} ; \quad (6:2)$$

where  $g_Z = \frac{g}{\sqrt{g^2 + g'^2}}$ . This mixing also affects the S parameter [16].

Mixing and direct Z<sup>0</sup> interactions together influence other, very low energy measurements. For example, in the class of models outlined above, the electron has an axial-vector coupling to the Z<sup>0</sup>. This is probed in atomic parity violation experiments, which are especially sensitive to the product of this coupling with the vector part of the isoscalar nuclear current [21]. The effective interaction is

$$L_{APV} = \frac{g_{Z^0}^2 (a^0 - a)(b + b)}{4M_{Z^0}^2} e^{-1} (u - u + d - d) ; \quad (6:3)$$

The product  $(a^0 - a)(b + b) = N(3N + 4)(u_1 - v_1)^2 = 16$  can be large in this model. Out of concern for this, I have tried to construct models within the present framework in which the electron's coupling to Z<sup>0</sup> is purely vectorial. So far, I have not found one that has a nontrivial  $(u_1 - v_1 \neq 0)$  solution to the anomaly conditions.

As a second example, the polarized Moller scattering experiment recently proposed by Kumar and his collaborators [22] is sensitive to the combination  $a^2 - a'^2$  of electron couplings to the Z<sup>0</sup>. The effective interaction is (apart from mixing effects)

$$L_{\text{Moller}} = \frac{g_{Z^0}^2}{2M_{Z^0}^2} a^2 (e_L - e_L)^2 + a'^2 (e_R - e_R)^2 ; \quad (6:4)$$

The  $Z^0$  will also be visible in current and planned high energy collider experiments. At subprocess energies well below the  $Z^0$  mass, its effects are still well approximated by four fermion "contact" interactions, similar to those expected for composite quarks and leptons [23]. Thus, at the Tevatron collider, the  $Z^0$ 's strong couplings to quarks produce an excess of high  $E_T$  jets<sup>6;7</sup> and high mass dileptons. The effective interactions are

$$L_{qq} = \frac{g_{Z^0}^2}{2M_{Z^0}^2} \sum_{q=u;d;c;s} (b_{qL} \bar{q}_L + b^0_{qR} \bar{q}_R) (a_{qL} \bar{q}_L + a^0_{qR} \bar{q}_R) + d (t_L \bar{t}_L + b_L \bar{b}_L) + d^0 (t_R \bar{t}_R + b_R \bar{b}_R) ; \quad (6:5)$$

$$L_{q\ell} = \frac{g_{Z^0}^2}{M_{Z^0}^2} \sum_{q=u;d;c;s} (b_{qL} \bar{q}_L + b^0_{qR} \bar{q}_R) \sum_{\ell=e;\mu;\tau} (a_{\ell L} \bar{\ell}_L + a^0_{\ell R} \bar{\ell}_R) ;$$

In these expressions, we have ignored all effects of mixing among quark generations. Note that there are simplifications of the couplings such as  $g_{Z^0}^2 b^2 = M_{Z^0}^2 (N+1)(N-2) = 3N F^2$ . The  $Z^0$  interaction affecting Bhabha scattering and muon pair production in  $e^+e^-$  collisions is

$$L_{\ell\ell} = \frac{g_{Z^0}^2}{2M_{Z^0}^2} \sum_{\ell=e;\mu;\tau} (a_{\ell L} \bar{\ell}_L + a^0_{\ell R} \bar{\ell}_R) (a_{\ell L} \bar{\ell}_L + a^0_{\ell R} \bar{\ell}_R) ; \quad (6:6)$$

Jet production in  $e^+e^-$  collisions is modified by  $L_{q\ell}$ . Corresponding interactions influence tau pair production. At the LHC, the excess of high  $E_T$  jets will be enormous and the  $Z^0$  shape should be observable as a resonance in dileptons if not in dijets. A high luminosity  $e^+e^-$  collider with  $\sqrt{s} \sim M_{Z^0}$  can make detailed studies of the  $Z^0$  couplings. One with  $\sqrt{s} \sim 500$  GeV may be able to detect signs of  $\gamma Z$  interference.

### Vacuum Alignment and Technihadron Physics

The spectrum of techninos  $T$  in this model is the same as that given above for the technipions. Determining the mass eigenstate  $T$  and  $\bar{T}$  is the problem of vacuum alignment in the technifermion sector. This is essentially the same as diagonalizing the

<sup>6</sup> The  $V_8$  colorons enhance only  $t\bar{t}$  and  $b\bar{b}$  production.

<sup>7</sup> As this paper was being completed, I received two preprints discussing the possibility that a TeV mass  $Z^0$  boson affects high  $E_T$  jet production and the branching ratios for  $Z^0$  decay to  $b\bar{b}$  and  $c\bar{c}$  [24].

technifermion mass matrix (see, however, footnote 4 for a caveat on the use of chiral perturbation theory.) The top-pions  $t$  formed from  $(t;b)_L$  and  $t_R$  must be added to this large  $T$  (diagonalization calculation. Once mass eigenstates are determined, the  $T \rightarrow T T$  couplings can be determined by symmetry (see, e.g., [14]). Note that the  $T$  decay modes may include one or two weak bosons,  $W_L$  and  $Z_L^0$ . Vacuum alignment also determines the pattern of technifermion condensation, relevant for mixing between heavy and light quarks, and feeds into the Kobayashi-Maskawa matrix and other quark mixing angles and phases.

Vacuum alignment is carried out by minimizing the ground state energy of broken ETC and  $SU(3)_1 \times U(1)$  four-fermion operators and of second order QCD interactions [15]. In the absence of a concrete ETC model, the most that can be done is to make "reasonable" guesses for the coefficients of allowed operators | those already assumed plus others consistent with symmetries. Different assumptions for the relative strengths and signs of the operators will lead to different vacua, patterns of condensation, and  $T$  and  $T$  spectroscopies. Such studies should give us a plausible range of expectations for this aspect of TC2 phenomenology. Some issues of immediate concern are:

Typical masses of the charged top-pion and its mixing with technipions. The concern here is that the decay rates for  $t \rightarrow t b$  or  $T \rightarrow b$  may be too large [25].

Masses of the  $T$  and  $T$ . Technipion decays are mediated by ETC interactions connecting technifermions to quarks and leptons. Thus, the  $T$  are expected to decay to heavy quark and lepton pairs. The existence of "leptoquark" decay modes such as  $T \rightarrow b$  depends on whether ETC operators such as  $b_R D_R^{\frac{1}{2}} D_L^{\frac{1}{2}} \psi_L$  are allowed. Experiments at the LEP collider will soon be able to set limits in excess of 75 GeV for charged  $T$ . Mixing between gluons and color octet  $T$  leads to copious production of colored  $T$ ; Tevatron collider searches should be able to discover them with masses up to several hundred GeV. Production of color singlet  $T \rightarrow T T, W_L T, Z_L T, W_L W_L$ , and  $W_L Z_L$  should be accessible at the Tevatron for  $T$  masses of 100-200 GeV [14]. Another process to be searched for at the Tevatron is  $gg \rightarrow T^0 \rightarrow bb$  or  $tt$ , if  $M_{T^0} > 2m_t$ . For the longer term,  $T$  and  $T$  masses are needed for LHC and large  $e^+e^-$  collider studies.

Vacuum alignment may produce phases in quark (and technifermion) mixing matrices that induce detectable CP violation in the neutral K and B meson systems, in the neutron electric dipole moment, and so on. If this happens, it will be important to determine whether strong CP violation can be avoided.

These brief remarks only scratch the surface of the phenomenological aspects of the scenario I have presented. I do hope they give a flavor of the richness of topcolor{assisted technicolor. I do not expect the specific class of models described here to pass all the tests it faces. But, in facing them, I expect we will learn how to build more complete and more successful models.

#### Acknowledgements

I am grateful to Claudio Rebbi for providing a program to solve the nonlinear equations for the hypercharges. I thank Chris Hill, Estia Eichten, Sekhar Chivukula, Krishna Kumar and Elizabeth Simmons for helpful comments. I have benefited from the hospitality of the Aspen Center for Physics where this work was begun. This research was supported in part by the Department of Energy under Grant No. DE-FG02-91ER40676.

## Appendix: Anomaly Conditions and Hypercharge Solutions

There are 5 linear and 4 cubic equations for the hypercharges in Table 1 arising from the requirement that  $U(1)_i$  gauge anomalies cancel:

$$U(1)_{1,2} [SU(N)]^2 : \quad x_1 x_1^0 + y_1 y_1^0 + z_1 z_1^0 \\ x_2 x_2^0 + y_2 y_2^0 + z_2 z_2^0 = \frac{1}{2} (N-2) ( \quad )^0$$

$$U(1)_{1,2} [SU(3)_1]^2 : \quad d = N (u_1 - v_1)$$

$$U(1)_{1,2} [SU(3)_2]^2 : \quad b - b^0 = \frac{1}{2} N (u_1 - v_1)$$

$$U(1)_{1,2} [SU(2)]^2 : \quad 2(a + 3b) + (c + 3d) = N [\beta (u_1 + v_1) + x_1 + y_1 + z_1] \\ = N [\beta (u_2 + v_2) + x_2 + y_2 + z_2]$$

$$[U(1)_1]^3 : \quad 0 = \frac{1}{2} N (N-1) ( \quad )^3 \\ + 2 [2a^3 - a^0 b^3 + 6(b^3 - b^0 b^3)] + 2c^3 - c^0 c^3 + 6d^3 \\ + 2N (x_1^3 - x_1^0 x_1^3 + y_1^3 - y_1^0 y_1^3 + z_1^3 - z_1^0 z_1^3)$$

$$[U(1)_2]^3 : \quad 0 = \frac{1}{2} N (N-1) ( \quad )^3 \\ 2 [2a^3 - a^0 b^3 + 6(b^3 - b^0 b^3)] - (2c^3 - c^0 c^3 + 6d^3) \\ + 2N [x_2^3 - x_2^0 x_2^3 + y_2^3 - y_2^0 y_2^3 + z_2^3 - z_2^0 z_2^3 - \frac{9}{4} (u_2 + v_2) - \frac{3}{4} (x_2^0 + y_2^0 + z_2^0)] \\ + 2 [\beta (a^0 - a^2) + 3(a^0 - \frac{1}{2}a) + 3(b^2 - b^0) + 5b^0 - \frac{1}{2}b] \\ + 3(c^0 - c^2) + 3(c^0 - \frac{1}{2}c) + 3(d^2 - d^0) + 3d^0 - \frac{1}{2}d$$

$$[U(1)_1]^2 U(1)_2 : \quad 0 = \frac{1}{2} N (N-1) ( \quad )^3 \\ 2 [2a^3 - a^0 b^3 + a^2 - a^0 a^2 + 6(b^3 - b^0 b^3) + b^0 - b^2] \\ (2c^3 - c^0 c^3 + c^2 - c^0 c^2 + 6d^3 + d^0 - d^2) \\ + 2N (x_1^2 x_2 - x_1^0 x_2^0 + y_1^2 y_2 - y_1^0 y_2^0 + z_1^2 z_2 - z_1^0 z_2^0)$$

$$[U(1)_2]^2 U(1)_1 : \quad 0 = \frac{1}{2} N (N-1) ( \quad )^3 \\ + 2 [2a^3 - a^0 b^3 + 2(a^2 - a^0 a^2) + 6(b^3 - b^0 b^3) + 2(b^0 - b^2)] \\ + 2c^3 - c^0 c^3 + 2(c^2 - c^0 c^2) + 6d^3 + 2(d^0 - d^2)$$

$$\begin{aligned}
& + 2N [x_2^2 x_1 - x_2^0 x_1^0 + y_2^2 y_1 - y_2^0 y_1^0 + z_2^2 z_1 - z_2^0 z_1^0] \\
& \quad \frac{3}{4} (u_1 + v_1) - \frac{1}{4} (x_1^0 + y_1^0 + z_1^0)] \\
& + 2 \left( \frac{1}{2} a^0 - a^0 + \frac{1}{6} b^0 - \frac{5}{3} b^0 \right) + \frac{1}{2} c^0 - c^0 + \frac{1}{6} d^0 - d^0 : \tag{A 1}
\end{aligned}$$

These 4 cubic equations are not independent because the  $[U(1)_Y]^3 = [U(1)_1 + U(1)_2]^3$  anomaly cancellation is guaranteed by the  $U(1)_Y [SU(2)]^2$  condition. A convenient set of 3 independent cubic equations consists of  $[U(1)_1]^3$  plus  $[U(1)_1]^2 U(1)_Y$  and  $[U(1)_1]^3 + [U(1)_2]^3 - 3[U(1)_1]^2 U(1)_Y$ :

$$\begin{aligned}
[U(1)_1]^2 U(1)_Y : 0 = & 2(a^0 - a^2 + b^2 - b^0) + c^0 - c^2 + d^2 - d^0 \\
& + 2N [(x_1 + x_2)(x_1^2 - x_1^0) + (y_1 + y_2)(y_1^2 - y_1^0) + (z_1 + z_2)(z_1^2 - z_1^0)]
\end{aligned}$$

$$\begin{aligned}
[U(1)_1]^3 + [U(1)_2]^3 - 3[U(1)_1]^2 U(1)_Y : \\
0 = (u_1 - v_1) f 2N^2 [4(y_1 + y_2)^2 - (x_1 + x_2)^2 + \frac{3}{4}] - (5N + 2)g : \tag{A 2}
\end{aligned}$$

In the last equation, I used results from Eq. (5.13).

The 18 linear and 3 nonlinear equations satisfied by the 26 hypercharges do not determine them uniquely. I sought numerical solutions to them that have  $u = \frac{1}{2}(u_1 - v_1) \notin 0$  as follows: First, I set  $v = 0$  and  $c = a$ . Then I chose values for  $x, y_1$  and  $y_1 + y_2$ , and solved for  $u, a$  and  $x_1 + x_2$ . To obtain  $u_1 - v_1 = 0(1)$ , I input  $x_1, y_1 = 0(Nu)$ . For  $N = 4$  and  $y_1 + y_2 = 0$  (which implies  $x_1 + x_2 = \frac{1}{4}$ ) and  $x_1 = y_1 = 10$ , I obtained

$$\begin{aligned}
u = 1.075; \quad a = 1.040 \quad (\text{for } x_1 + x_2 = \frac{1}{4}) \\
u = 1.197; \quad a = 12.054 \quad (\text{for } x_1 + x_2 = \frac{1}{4}) : \tag{A 3}
\end{aligned}$$

As is apparent from Eqs. (A 2), these solutions scale linearly with the input values of  $x_1$  and  $y_1$ . Values of  $a$  as large as 12 are doubtless ruled out.

## References

- [1] C. T. Hill, *Phys. Lett. B* 345, 483 (1995).
- [2] C. T. Hill, *Phys. Lett. B* 266, 419 (1991) ;  
S. P. Martin, *Phys. Rev. D* 45, 4283 (1992); *ibid D* 46, 2197 (1992); *Nucl. Phys. B* 398, 359 (1993); M. Lindner and D. Ross, *Nucl. Phys. B* 370, 30 (1992);  
R. Bonisch, *Phys. Lett. B* 268, 394 (1991);  
C. T. Hill, D. Kennedy, T. Onogi, H. L. Yu, *Phys. Rev. D* 47, 2940 (1993).
- [3] Y. Nambu, in *New Theories in Physics, Proceedings of the XI International Symposium on Elementary Particle Physics, Kazimierz, Poland, 1988*, edited by Z. Ajduk, S. Pokorski and A. Trautmann (World Scientific, Singapore, 1989); Enrico Fermi Institute Report EFI 89-08 (unpublished);  
V. A. Miransky, M. Tanabashi and K. Yamawaki, *Phys. Lett. B* 221, 177 (1989);  
*Mod. Phys. Lett. A* 4, 1043 (1989);  
W. A. Bardeen, C. T. Hill and M. Lindner, *Phys. Rev. D* 41, 1647 (1990).
- [4] S. Weinberg, *Phys. Rev. D* 19, 1277 (1979);  
L. Susskind, *Phys. Rev. D* 20, 2619 (1979).
- [5] S. Dimopoulos and L. Susskind, *Nucl. Phys. B* 155, 237 (1979);  
E. Eichten and K. Lane, *Phys. Lett. B* 90, 125 (1980).
- [6] F. Abe, et al, The CDF Collaboration, *Phys. Rev. Lett.* 73, 225 (1994); *Phys. Rev. D* 50, 2966 (1994); *Phys. Rev. Lett.* 74, 2626 (1995) ;  
S. Abachi, et al, The D Collaboration, *Phys. Rev. Lett.* 74, 2632 (1995).
- [7] R. S. Chivukula, B. A. Dobrescu and J. Teming, *Phys. Lett. B* 353, 289 (1995).
- [8] D. Komiris, Boston University Preprint BUHEP-95-20, hep-ph/9506305 (1995).
- [9] K. Lane and E. Eichten, *Phys. Lett. B* 352, 382 (1995).
- [10] See E. Eichten and K. Lane in Ref. [5]
- [11] Particle Data Group, *Phys. Rev. D* 50, 1174 (1994).
- [12] B. Holdom, *Phys. Rev. D* 24, 1441 (1981); *Phys. Lett. B* 150, 301 (1985);  
T. Appelquist, D. Karabali and L. C. R. Wijewardhana, *Phys. Rev. Lett.* 57, 957 (1986); T. Appelquist and L. C. R. Wijewardhana, *Phys. Rev. D* 36, 568 (1987);  
K. Yamawaki, M. Bando and K. Matumoto, *Phys. Rev. Lett.* 56, 1335 (1986) ;  
T. Akiba and T. Yanagida, *Phys. Lett. B* 169, 432 (1986).
- [13] C. T. Hill and S. Parke, *Phys. Rev. D* 49, 4454 (1994)
- [14] K. Lane and E. Eichten, *Phys. Lett. B* 222, 274 (1989) ;  
K. Lane and M. V. Ramana, *Phys. Rev. D* 44, 2678 (1991).
- [15] R. Dashen, *Phys. Rev. D* 3, 1879 (1971); S. Weinberg, *Phys. Rev. D* 13, 974 (1976);  
E. Eichten, K. Lane and J. Preskill, *Phys. Rev. Lett.* 45, 225 (1980);  
K. Lane, *Physica Scripta* 23, 1005 (1981);  
M. Peskin, *Nucl. Phys. B* 175, 197 (1980); J. Preskill, *Nucl. Phys. B* 177, 21 (1981).

- [16] A. Longhitano, Phys. Rev. D 22 (1980) 1166; Nucl. Phys. B 188, (1981) 118;  
 R. Renken and M. Peskin, Nucl. Phys. B 211 (1983) 93;  
 B. W. Lynn, M. E. Peskin and R. G. Stuart, in Trieste Electroweak 1985, 213;  
 M. Golden and L. Randall, Nucl. Phys. B 361 (1990) 3;  
 B. Holdom and J. Terning, Phys. Lett. B 247 (1990) 88;  
 M. E. Peskin and T. Takeuchi, Phys. Rev. Lett. 65 (1990) 964;  
 A. Dobado, D. Espriu and M. J. Herrero, Phys. Lett. B 255 (1990) 405;  
 H. Georgi, Nucl. Phys. B 363 (1991) 301.
- [17] K. Lane, Proceedings of the 27th International Conference on High Energy Physics,  
 edited by P. J. Bussey and I. G. Knowles, Vol. II, p. 543, Glasgow, June 20-27, 1994.
- [18] B. Holdom, Phys. Lett. B 259, 329 (1991);  
 E. Gates and J. Terning, Phys. Rev. Lett. 67, 1840 (1991);  
 M. Luty and R. Sundrum, Phys. Rev. Lett. 70, 127 (1993);  
 T. Appelquist and J. Terning, Phys. Lett. B 315, 139 (1993).
- [19] K. Lane, Phys. Rev. D 52, 1546 (1995).
- [20] Studies along these lines are in progress by R. S. Chivukula and J. Terning.
- [21] P. Langacker, Theoretical Study of the Electroweak Interaction | Present and Future,  
 Proceedings of the 22<sup>nd</sup> INS Symposium on Physics with High Energy Colliders,  
 Tokyo, March 1994.
- [22] K. Kumar, E. Hughes, R. Holmes and P. Souder, "Precision Low Energy Weak Neutral  
 Current Experiments", Princeton University (October 30, 1995), to appear in  
 Mod. Phys. Lett. A.
- [23] E. Eichten, K. Lane and M. Peskin, Phys. Rev. Lett. 50, 811 (1983).
- [24] P. Chiappetta, et al., CPT-96/P.3304, hep-ph 9601306;  
 G. Altarelli, et al., CERN-TH/96-20, hep-ph 9601324.
- [25] This concern was first raised by Chris Hill. The problem is under study by B. Bajic.

Particle	SU (3) <sub>1</sub>	SU (3) <sub>2</sub>	Y <sub>1</sub>	Y <sub>2</sub>
$\nu_L^1$	1	1	a	$\frac{1}{2}$ a
$e_R^1, \mu_R^1$	1	1	$a^0$	1 $\hat{a}$
$q_L^1$	1	3	b	$\frac{1}{6}$ b
$u_R^1, c_R^1$	1	3	$b^0$	$\frac{2}{3}$ $\hat{b}$
$d_R^1, s_R^1$	1	3	$b^0$	$\frac{1}{3}$ $\hat{b}$
$\nu_L^h$	1	1	c	$\frac{1}{2}$ c
$\nu_R^h$	1	1	$c^0$	1 $\hat{c}$
$q_L^h$	3	1	d	$\frac{1}{6}$ d
$t_R^h$	3	1	$d^0$	$\frac{2}{3}$ $\hat{d}$
$b_R^h$	3	1	$d^0$	$\frac{1}{3} + d^0$
$T_L^1$	3	1	$u_1$	$u_2$
$U_R^1$	3	1	$v_1$	$v_2 + \frac{1}{2}$
$D_R^1$	3	1	$v_1$	$v_2 - \frac{1}{2}$
$T_L^2$	1	3	$v_1$	$v_2$
$U_R^2$	1	3	$u_1$	$u_2 + \frac{1}{2}$
$D_R^2$	1	3	$u_1$	$u_2 - \frac{1}{2}$
$T_L^1$	1	1	$x_1$	$x_2$
$U_R^1$	1	1	$x_1^0$	$x_2^0 + \frac{1}{2}$
$D_R^1$	1	1	$x_1^0$	$x_2^0 - \frac{1}{2}$
$T_L^t$	1	1	$Y_1$	$Y_2$
$U_R^t$	1	1	$Y_1^0$	$Y_2^0 + \frac{1}{2}$
$D_R^t$	1	1	$Y_1^0$	$Y_2^0 - \frac{1}{2}$
$T_L^b$	1	1	$z_1$	$z_2$
$U_R^b$	1	1	$z_1^0$	$z_2^0 + \frac{1}{2}$
$D_R^b$	1	1	$z_1^0$	$z_2^0 - \frac{1}{2}$
$L$	1	1		
$R$	1	1	0	0

TABLE 1: Lepton, quark and technifermion colors and hypercharges.