

Michael C . B irse

Theoretical Physics Group, Department of Physics and Astronomy

University of Manchester, Manchester, M13 9PL, UK

The contribution of nucleons to the quark condensate in nuclear matter includes a piece of first order in m_π , arising from the contribution of low-momentum virtual pions to the N sigma commutator. Chiral symmetry requires that no term of this order appears in the NN interaction. The mass of a nucleon in matter thus cannot depend in any simple way on the quark condensate alone. More generally, pieces of the quark condensate that arise from low-momentum pions should not be associated with partial restoration of chiral symmetry.

There has been much recent interest in the question of whether chiral symmetry is partially restored in nuclear matter.¹ The chiral isospin symmetry $SU(2) \times SU(2)$ is an approximate symmetry of the QCD Lagrangian, broken only by the small current masses of the up and down quarks. This symmetry is realised in the hidden (spontaneously broken) mode since the QCD vacuum is not invariant under it. The scalar density of quarks, often referred to as the "quark condensate," provides an order parameter describing the hidden symmetry. Any reduction of this condensate in matter has generally been interpreted as a signal of partial restoration of the symmetry.

¹For reviews of this topic and further references, see [1-3].

The quark condensate in vacuum can be related to pion properties through the Gell-Mann-Oakes-Renner (GOR) relation,

$$m^2 f^2 = 2m \langle \bar{q}q \rangle_0; \quad (1)$$

where m is the average of the current masses of the up and down quarks and $\langle \bar{q}q \rangle_0$ is the quark condensate (per quark flavour) in vacuum. The change in this condensate in matter, to first order in the density, can be estimated in a model-independent way from [4,6]

$$\frac{\langle \bar{q}q \rangle}{\langle \bar{q}q \rangle_0} = 1 - \frac{N}{f^2 m^2}; \quad (2)$$

where N is the pion-nucleon sigma commutator,

$$N = 2m_N \int d^3r \bar{q} \gamma_5 q N; \quad (3)$$

The current best determination of N from N scattering is 45 ± 7 MeV [7]. Using this in (2) suggests a 30% reduction in the quark condensate at normal nuclear matter densities.

There has been some debate about corrections to this result [5,8,9]. Estimates using both simple models [5,10,9] and relativistic BHF calculations with realistic NN forces [11,12] suggest that corrections are small up to normal nuclear densities, although they rapidly become large (and highly model-dependent) beyond that. There have also been questions about whether the effect is sufficiently short-ranged that the hard-core correlations between nucleons mean that one nucleon does not feel the change in the condensate produced by its neighbours [13]. However, the strong coupling of a scalar meson, which can be interpreted as an excitation of the condensate, to two-pion states suggests that the effect should be long-ranged and so not cut out by the correlations [16].

In contrast, there has been little discussion of whether changes in the quark condensate necessarily mean partial symmetry restoration, or indeed if those changes can have directly observable effects on particle properties in matter. It is these questions that I address here.

An indication that $\langle \bar{q}q \rangle$ may not be the most relevant quantity is provided by the chiral expansion of the energy of a nucleon in nuclear matter. To first order in the density the

mass of a nucleon in matter (at mean-field level) can be written in the form

$$M_N = M_N + V_S(0) ; \quad (4)$$

where $V_S(q^2)$ is the part of the NN potential (more specifically, the two-nucleon irreducible NN scattering amplitude) with a scalar-exchange character.² Suppose, for the sake of argument, that the mass in matter contained a piece that depended in some way on the quark condensate so that, to first order in the density, there was a term in M_N proportional to $\langle \bar{\psi}\psi \rangle = (f^2 m^2)$. This would imply the existence of a scalar piece of the NN interaction that was proportional to $\langle \bar{\psi}\psi \rangle = m^2$ at zero momentum transfer [16].

The chiral expansion of M_N has the form [17]

$$M_N = A m^2 + \frac{9}{16} \frac{g_{NN}^2}{2M_N} m^3 + \dots ; \quad (5)$$

where the constant A involves a counterterm corresponding to short-distance physics, whose value must be determined empirically from NN scattering. In contrast, the $O(m^3)$ term is nonanalytic in the current quark mass and so arises purely from the long-distance physics of the pion cloud surrounding a nucleon. Its coefficient is given entirely in terms of the NN coupling and nucleon mass. If there were a piece of the NN interaction proportional to $\langle \bar{\psi}\psi \rangle = m^2$ (with a coefficient independent of m), it would contain a corresponding term of order m . However Weinberg's dimensional counting rules [15]³ show that no such term can be present in the chiral expansion of the NN potential. Chiral symmetry thus rules out any interaction proportional to $\langle \bar{\psi}\psi \rangle = m^2$ on its own. Either there are additional terms that cancel out the piece of order m , or the coefficient of $\langle \bar{\psi}\psi \rangle = m^2$ is at least of order m^2 .

²This definition corresponds to a mass that could appear in a Dirac equation for the nucleon. The irreducible NN amplitude in (4) avoids the problem of the strong energy dependence of the full scattering amplitude produced by the deuteron pole just below threshold, discussed in [14]. The irreducible amplitude is also the one to which Weinberg's chiral counting rules apply [15].

³For a review see [18].

(in which case the term could make only a small contribution to the full scalar-exchange interaction). Hence there can be no direct dependence of the nucleon mass in matter on the quark condensate (2).

As an illustration of this point, consider the calculations in Ref. [16] using the linear sigma model. There I defined a "symmetry restoring amplitude" as a sum of NN diagrams in which the meson coupled directly to one of the nucleons. This has precisely the form just discussed, proportional to $\langle \bar{\psi}\psi \rangle = m^2$. However as pointed out in [16], the full NN scattering amplitude arising from exchange of two virtual pions involves a different combination of diagrams. In these there is a strong cancellation between diagrams in which the pions couple to a nucleon via a σ and others in which the pions couple directly, just like the "pair suppression" in N scattering. These cancellations ensure that the $O(m)$ term arising from the sigma commutator does not appear in any chirally symmetric calculation of the full NN amplitude [19,16].

In general, the contributions to the quark condensate from low-momentum pions need to be distinguished from those of other hadrons because of the special role pions play as approximate Goldstone bosons. The integrated scalar density of quarks in a pion can be estimated by applying the Feynman-Hellmann theorem to the pion mass and using the fact that $m^2 \propto \langle \bar{\psi}\psi \rangle$, as in (1). This leads to

$$2m \langle \bar{q}q \rangle = m^2 \frac{dm}{dm^2} = \frac{1}{2} m^2 \approx 70 \text{ MeV} : \quad (6)$$

This is even larger than the corresponding matrix element for the nucleon (3). Hence the scalar density of quarks in a pion is large, reflecting the collective nature of a Goldstone boson. Note that (6) applies to a pion state that has been unconventionally normalised to unity. For a more familiar covariantly normalised pion, the corresponding matrix element is m^2 . If one naively assumed that the scattering of a nucleon from a pion contained a term proportional to the scalar quark density of the pion, then one would expect this to give a contribution to NN scattering of order m^0 . However it has long been known that this is not the case. Chiral symmetry requires that the isospin-averaged amplitude for pion-nucleon

scattering is of order m^2 at threshold [20]. The contribution of these pions to observables such as scattering amplitudes or the masses of nucleons and other hadrons (except pions and kaons) in matter is thus suppressed by a factor of m^2 compared with their quark density.

In the particular case of the NN interaction, the leading nonanalytic dependence on the current quark mass, the $O(m^3)$ term, in ϵ_N arises from the longest-ranged part of the pion cloud, corresponding to virtual pions of very low momentum. These pions are responsible for the $O(m)$ piece of the scalar density of the nucleon but, as in the real NN scattering just discussed, their contribution to the scattering of another nucleon is suppressed by a factor of m^2 . This can also be seen by applying the counting rules [15] to a two-pion-exchange diagram with one NN vertex of dimension $d = 2$. From this one finds a contribution to the NN interaction of order m^3 . Hence, as we have already seen, the part of (2) which is of order m cannot contribute a piece of the same order to the energy of a nucleon in matter.

Note that short-ranged correlations between nucleons, which can also destroy any simple dependence of the nucleon mass on the quark condensate in matter [13,21], are not relevant to the question here. Since the parts of the NN interaction under discussion arise from long-ranged pion exchanges, they are not affected by such short-range correlations.

Analogous effects can be found in the properties of nucleons in vacuum. The quark condensate in vacuum includes a piece of order $m^2 \ln m$, also generated by virtual pions [22]. However logarithmic terms in the chiral expansion of the nucleon mass start at order $m^4 \ln m$ [23]. In any model where the nucleon mass contains a term proportional to the quark condensate, there are again strong cancellations of that with other pion cloud contributions, as illustrated by the one-loop calculation of ϵ_N in the linear sigma model [16].

The QCD sum rule approach [24] to hadron properties involves a short-distance operator product expansion (OPE). This brings in matrix elements of local operators like $\bar{q}q$. These cannot discriminate between contributions from low-momentum virtual pions and those from other components of the QCD vacuum. The OPE side of a sum rule for the nucleon mass [25] thus contains a piece of order $m^2 \ln m$ [26]. A careful treatment of the continuum,

including low-momentum π N states, is needed on the phenomenological side of the sum rule to obtain an expression for the mass without the ending logarithm [27].

Matter at finite temperature also demonstrates similar effects. At low temperatures and zero baryon density this is a pion gas. The scalar quark density of this produces a temperature dependence of the quark condensate that starts at order T^2 [28]. In contrast the change in the mass of a nucleon in this gas starts at order T^4 [29]. In finite-temperature QCD sum rules for the nucleon the $O(T^2)$ dependence of the condensate is cancelled by including the π N scattering on the phenomenological side of the sum rule [30].

A proper treatment of the π N continuum in matter would presumably remove the corresponding $O(m)$ term that arises on the OPE side of sum rules for a nucleon in matter [31]. Until this is done, the corresponding piece of $\bar{h}\bar{q}q$ should be regarded as an additional source of uncertainty. Its size can be estimated from the $O(m^3)$ term in χ_N , which is 25 MeV and so is about half of the coefficient of the density in (2). This presence of this piece in the sum rules for a nucleon in matter should be regarded as a significant extra uncertainty in their results.

If these contributions from low-momentum pions to changes in the quark condensate do not affect the masses of other hadrons, one should also ask whether they correspond to restoration of chiral symmetry. In the framework of a sigma model, one can imagine large-amplitude, low-momentum fluctuations of the pion fields around the chiral circle (or indeed a pion condensate [32]) that could significantly reduce the mean value of the field without moving the fields off the chiral circle $(\phi(x))^2 + (\sigma(x))^2 = f^2$. Since the field corresponds to the scalar quark density of the underlying QCD description, one might be tempted to conclude from the its reduction that chiral symmetry has been partially restored. However the fact that fields remain on the chiral circle indicates that this is not the case. Similarly a reduction in $\bar{h}\bar{q}q$ from its vacuum value is not enough for one to conclude that the symmetry has been partially restored, if part of this change arises from low-momentum pions.

As a corollary to this, one can imagine situations where the quark condensate vanishes but chiral symmetry is not restored. For example, this could happen if the linear density

dependence of (2) were to continue to hold up to around three times the density of nuclear matter, although that seems unlikely from present estimates of higher-order density dependence [11,12]. The pion would then be anomalously light, in the sense described in [33]. The leading term in the GOR relation would vanish and the pion mass would be proportional to the current quark mass rather than its square root.⁴ This could indicate proximity to a pion-condensed phase of the type described in [32]. In any case other condensates can play an increasingly important role as $\bar{h}q_i$ tends to zero.

In summary: the quark condensate in matter contains pieces arising from low-momentum virtual pions. These do not contribute to nucleon properties in matter and cannot be associated with a partial restoration of chiral symmetry.

ACKNOWLEDGEMENTS

I am grateful to B. Krippa for the question that prompted this work, and to him and J. McGovern for their comments on it. This work was supported by the EPSRC and PPARC.

[1] C. Adam and G. E. Brown, Phys. Reports 224 (1993) 1.

[2] M. C. Birse, J. Phys. G: Nucl. Part. Phys. 20 (1994) 1537.

[3] G. E. Brown and M. Rho, hep-ph/9504250, Phys. Reports (to appear).

[4] E. G. Drukarev and E. M. Levin, Nucl. Phys. A 511 (1990) 679; A 516 (1990) 715 (E).

[5] T. D. Cohen, R. J. Furnstahl and D. K. Griegel, Phys. Rev. C 45 (1992) 1881.

[6] M. Lutz, S. Klint and W. Weise, Nucl. Phys. A 542 (1992) 521.

⁴There is a school of chiral perturbation theory that suggests that this is already the situation in the normal vacuum. See, for example [34].

- [7] J. Gasser, H. Leutwyler and M. E. Sainio, Phys. Lett. B 253 (1991) 252, 260.
- [8] M. Ericson, Phys. Lett. B 301 (1993) 11.
- [9] M. C. Birse and J. A. McGovern, Phys. Lett. B 309 (1993) 231.
- [10] G. Chanfray and M. Ericson, Nucl. Phys. A 56 (1993) 427.
- [11] G. Q. Li and C. M. Ko, Phys. Lett. 338 (1994) 118.
- [12] R. Brockmann and W. Weise, Phys. Lett. B 367 (1996) 40.
- [13] T. E. O. Ericson, Phys. Lett. B 321 (1994) 312.
- [14] R. J. Furnstahl and T. Hatsuda, Phys. Rev. Lett. 75 (1994) 3128.
- [15] S. Weinberg, Phys. Lett. 251 (1990) 288; Nucl. Phys. B 363 (1991) 1; Phys. Lett. B 295 (1992) 114.
- [16] M. C. Birse, Phys. Rev. C 49 (1994) 2212.
- [17] J. Gasser, Ann. Phys. 136 (1981) 62;
J. Gasser and H. Leutwyler, Phys. Report 87 (1982) 77.
- [18] V. Bernard, N. Kaiser and U.-G. Meissner, Int. J. Mod. Phys. E 4 (1995) 193.
- [19] L. S. Celenza, A. Pantziris and C. M. Shakin, Phys. Rev. C 46 (1992) 2213;
C. Ordóñez, L. Ray and U. van Kolck, Phys. Rev. Lett. 72 (1994) 1982; hep-ph/9511380;
J. L. Friar and S. A. Coon, Phys. Rev. C 49 (1994) 1272;
C. A. da Rocha and M. R. Robilotta, Phys. Rev. C 49 (1994) 1818;
M. Robilotta, Nucl. Phys. A 595 (1995) 171;
J.-L. Ballot, M. R. Robilotta and C. A. da Rocha, hep-ph/9502369.
- [20] S. Weinberg, Phys. Rev. Lett. 17 (1966) 616.
- [21] J. Delorme, G. Chanfray and M. Ericson, University of Lyon preprint LYCEN 9604, nucl-th/9603005.

- [22] P. Langacker and H. Pagels, Phys. Rev. D 8 (1973) 4595;
J. Gasser and H. Leutwyler, Ann. Phys. 158 (1984) 142.
- [23] J. Gasser, M. E. Sainio and A. Svarc, Nucl. Phys. 307 (1988) 779.
- [24] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B 147 (1979) 385, 448, 519.
- [25] B. L. Io e, Nucl. Phys. B 188 (1981) 317; B 191 591 (E).
- [26] D. K. Griegel and T. D. Cohen, Phys. Lett. B 333 (1994) 27.
- [27] S. H. Lee, S. Choe, T. D. Cohen and D. K. Griegel, Phys. Lett. B 348 (1995) 263.
- [28] P. Gerber and H. Leutwyler, Nucl. Phys. B 321 (1989) 387.
- [29] H. Leutwyler and A. V. Smilga, Nucl. Phys. B 342 (1990) 302.
- [30] Y. Koike, Phys. Rev. D 48 (1993) 2313.
- [31] T. D. Cohen, R. J. Furnstahl and D. K. Griegel, Phys. Rev. Lett. 67 (1991) 961; T. D. Cohen,
D. K. Griegel and R. J. Furnstahl, Phys. Rev. C 46 (1992) 1507.
- [32] G. E. Brown, V. Koch and M. Rho, Nucl. Phys. A 535 (1991) 701.
- [33] T. D. Cohen and W. Broniowski, Phys. Lett. B 342 (1995) 25; B 348 12.
- [34] M. D. Scadron, Rep. Prog. Phys. 44 (1981) 213.