Statistical Properties of the Linear Sigm a M odel

J rgen Randrup

Nuclear Science Division, Law rence Berkeley National Laboratory University of California, Berkeley, California 94720

January 12, 1996

Abstract:

The statistical equilibrium properties of the linear sigm a model are studied, with a view towards characterizing the eld con gurations employed as initial conditions for num erical simulations of the formation of disoriented chiral condensates in high-energy nuclear collisions. The eld is decomposed into its spatial average, the order parameter, and the

uctuations, the quasi-particles, and enclosed in a rectangular box with periodic boundary conditions. The quantized quasi-particle modes are described approximately by Klein-G ordon dispersion relations containing an e ective m ass that depends on both the tem perature and the magnitude of the order parameter. The therm al uctuations are instrumental in shaping the e ective potential governing the order parameter, and the evolution of its statistical distribution with tem perature is discussed, as is the behavior of the associated e ective masses. As the system is cooled the eld uctuations subside, causing a sm ooth change from the high-tem perature phase in which chiral symmetry is approximately restored towards the norm alphase. Of practical interest is the fact that the equilibrium eld con qurations can be sampled in a simple manner, thus providing a convenient means for specifying the initial conditions in dynam ical simulations of the non-equilibrium relaxation of the chiral eld. The corresponding correlation function is brie y considered and used to calculate the spectral strength of radiated pions. Finally, by propagating sam ples of initial con gurations by the exact equation of motion, it has been ascertained that the treatment is su ciently accurate to be of practical utility.

This work was supported by the D irector, O \propto of Energy Research, O \propto of H igh Energy and N uclear P hysics, N uclear P hysics D ivision of the U S.D epartm ent of Energy under C ontract N o. D E -A C 03-76SF 00098, and by the N ational Institute for N uclear T heory at the U niversity of W ashington in Seattle.

1 Introduction

The possibility of producing and observing disoriented chiral condensates in highenergy collisions of hadrons and nuclei has stimulated considerable interest over the past few years (for a recent review, see ref. [1, 2]). The basic premise is that the collision generates an extended domain of space within which chiral symmetry is approximately restored. If this happens, macroscopic pion elds may be generated as a consequence of the subsequent non-equilibrium relaxation towards the normal state. Such isospin-aligned domains may manifest them selves in anomalous pion emission [3, 4, 5] of the type seen in the Centauro cosm ic ray events [6]. The experimental exploration of this phenomenon is of fundamental interest because it has a direct bearing on the mechanism of spontaneous chiral symmetry breaking. E orts are well underway to search for the associated pion multiplicity uctuations in proton-proton collisions [7].

In order to assess the prospects for such a phenom enon to actually occur and be detectable above the background of other pion production processes, it is necessary to perform extensive dynam ical calculations. This is a daunting task because the chiral degrees of freedom should be properly embedded in the complicated environm ent generated in a high-energy collision which evolves from being primarily partonic at the early stage to entirely hadronic in the course of the chiral relaxation process. Fortunately, the study of suitable idealized scenarios can yield valuable insights regarding the prospects for observing the e ect.

M ost dynam ical studies have been carried out within the fram ework of the linear m odel, treated in them ean-ekd approxim ation [8,9,10,11,12,13,14,15,16,17]. The physical problem divides naturally into three parts: 1) speci cation of the initial state in which chiral symmetry is approximately restored; 2) the dynamical evolution during which the chiral ekd cools and evolves towards the neighborhood of the physical vacuum; and 3) the analysis of the asymptotic state in terms of physical observables. The present work is particularly relevant for the rst issue, as it provides a wellbased fram ework for characterizing the chiral ekd con gurations that are employed as initial states in the numerical simulations. In particular, a convenient method is developed for sampling the statistical equilibrium ensemble of chiral con gurations and is expected to be of practical utility.

A fter a brief rem inder of the relevant features of the linear m odel (sect. 2), we describe how them al equilibrium can be treated approximately by means of a standard linearization procedure (sect. 3). We then discuss and illustrate the statistical distribution of the average eld strength, the order parameter (sect. 4), and subsequently turn to the properties of the quasi-particle degrees of freedom associated with the spatial eld uctuations (sect. 5). An impression of the validity of the treatment is then gained by evolving samples of eld con gurations by the exact equation of motion (sect. 6). Finally, a concluding discussion is given (sect. 7).

2 The linear model

To set the fram ework for the subsequent developm ents, we start by brie y recalling the most relevant features of the form all fram ework. The present study is carried out within the mean-eld approximation where the quantum eld operators are replaced by their expectation values, thereby bringing the treatment to the level of classical eld theory. This simplied treatment is expected to suce for exploratory calculations. Naturally, the mean-eld approximation is only a rst step towards a more complete description.

The basic object of study is then the chiral eld

$$(r;t) = (r;t) + i$$
 $(r;t);$ (1)

where the three elements of the vector are the Paulim atrices. Here the scalar eld

(r;t) and the vector eld (r;t) are both real and can conveniently be combined into the O (4) vector = (;).

In the linear model [18], one introduces a simple local e ective interaction energy density, ______#

$$V() = \frac{1}{h^{3}c^{3}} - \frac{1}{4}(^{2}v^{2})^{2} + H^{*}; \qquad (2)$$

where denotes the magnitude of the O (4) vector ,

2

$$= \int_{j=0}^{X^3} (r;t)^2 = {}^2 + ; \qquad (3)$$

with j = 0;1;2;3 referring to the four chiral directions. The interaction V contains three parameters: , the strength of the symmetric term; v, the location of its m inim um; and H, the strength of the symmetry-breaking term. As is commonly done, these parameters are xed by specifying the pion decay constant, f = 92 MeV, and the meson masses, $m = 138 \text{ MeV} = c^2$ and $m = 600 \text{ MeV} = c^2$, leading to

$$= \frac{m^2 c^2 m^2 c^2}{2f^2} = 20:14 ; \qquad (4)$$

$$\mathbf{v} = \frac{\mathbf{m}^2 - 3\mathbf{m}^2}{\mathbf{m}^2 - \mathbf{m}^2} \mathbf{f}^2 = 86:71 \, \mathrm{MeV} \; ; \tag{5}$$

$$H = m^2 c^2 f = (120.55 M eV)^3$$
: (6)

The precise values of the model parameters are immaterial, both in view of the simple nature of the model and, in particular, within the context of our present idealized study. The value for the hypothetical mass is the most commonly adopted one; for m we have used the weighted average of the three observed pion masses, and for the pion decay constant we have simply taken two thirds of the pion mass which gives a simple value within the range of experimental data (the current values being f = 92.4 0.26 MeV and $f_0 = 84$ 3 MeV [19]).

The Lagrangian density is given by

L (r;t) =
$$\frac{1}{h^3 c^3} = \frac{1}{2} (h \theta_t)^2 = \frac{1}{2} (h cr)^2 = \frac{1}{4} (c^2 - v^2)^2 + H^{(-)}$$
 (7)

The corresponding energy density is then

$$H (r;t) = \frac{1}{h^{3}c^{3}} \frac{1}{2}^{2} + \frac{1}{2}(hcr)^{2} + \frac{1}{4}(r^{2} v^{2})^{2} H ; \qquad (8)$$

where the time derivative hQ_t is the canonical conjugate of the eld strength.

The action generated by a given time evolution is given by S = drdtL(r;t). By demanding that S be stationary with respect to arbitrary variations of both the eld strength (r;t) and its rst derivatives, one obtains the associated equation of motion,

$$2 + (^{2} v^{2}) = H^{2};$$
 (9)

where $2 = h^2 \theta_t^2 + h^2 c^2 4$ is the d'A lam bert di erential operator, and $^$ denotes the unit vector in the direction. Since the equation of motion is of second order in time, the evolution of the system is fully determined once the initial values of the eld strength (r) and the associated time derivative (r) have been specified.

3 Approxim ate treatm ent

We now discuss how the chiral system can be treated in a convenient approximate manner. For this purpose, we con ne the system within a rectangular box and impose periodic boundary conditions.

3.1 Linearization

Our starting point is the equation of m otion (9) for the eld strength. It is natural to decompose the eld,

$$(r;t) = (t) + (r;t);$$
 (10)

where the rst term is the spatial average, $\langle \rangle$, so that $\langle \rangle = 0$. The average part of the eld, _, is often referred to as the order parameter, whereas the spatial uctuations, (r), may be considered as quasi-particle degrees of freedom representing elementary excitations relative to the constant eld con guration [10]. The O (4) direction of the order parameter plays a special role and it it convenient to employ a correspondingly aligned reference system, i.e. a system in which _ = (_0;0) and _ = (_k;_?). The therm all average is then a diagonal 4 4 tensor, as can be demonstrated by elementary means.

Taking the spatial average of the full equation of m otion (9) leads to an equation for time evolution of the order parameter [20],

$$h^{2}Q_{t}^{2} + [_{0}^{2} + 2 + 2]_{k}^{2} v^{2}]_{-} = H^{2};$$
 (11)

In deriving this equation, we have replaced the spatial averages < > by them al averages , as may be justiled for a system of dimensions larger than the correlation length (the uctuations in dimensions of the system are then independent and can therefore be assumed to reject the them all distribution from which the particular eld was sampled). We shall use $\frac{2}{2}$ to denote the diagonal elements of the isotropic 3 3 tensor $< \frac{2}{2}$, so that $2 = \frac{2}{k} + 3 = \frac{2}{2}$ is the total uctuation.¹ Furtherm ore, < 2 > vanishes by symmetry.

By subtracting the equation of motion for the order parameter, eq. (11), from the full equation of motion (9), it is possible to obtain approximate equations for the eld uctuations,

$$2 + \frac{2}{2}c^4 \qquad = 0;$$
 (13)

where the elective m asses $_{k}$ and $_{?}$ are determ ined by the auxiliary gap equations,

$${}^{2}_{k}c^{4}$$
 ${}^{3}_{0}{}^{2}_{0}+$ ${}^{2}_{k}+2$ ${}^{2}_{k}$ v^{2} ; (14)

$$c^{2}c^{4}$$
 $c^{2}+c^{2}+2$ $c^{2}+2$ c^{2} v^{2} : (15)

In arriving at this result, we have replaced products of two individual uctuations by their respective therm all ensemble average values, $j = j^0$; $j = j^0$, and, furtherm ore, products containing three uctuation factors have been contracted in the usualm anner [15],

Equations (12) and (13) describe the uctuations as independent quasi-particles having the respective e ective masses $_{\rm k}$ and $_{?}$, which in turn are given in terms of the eld uctuations, eqs. (14) and (15). This self-consistent relationship can be stated explicitly by invoking the expression for the associated therm al equilibrium uctuations,

$${}^{2}_{k} = \frac{h^{3}c^{3}X}{k} \frac{01}{k} \frac{1}{e^{k}} \frac{1}{1} \frac{1}{2^{2}} \frac{1}{k} c^{2} \frac{1}{n} K_{1} (n \frac{k^{2}}{T}); \quad (17)$$

and similarly for $\frac{2}{2}$. The system is enclosed in a rectangular box with the volum $e = L_x L_y L_z$ and periodic boundary conditions are imposed. The wave vectors k are then quantized $(k_x L_x = 2 K_x \text{ with } K_x = 0; 1; :::, \text{etc.})$, so that the individual quasi-particle modes are enumerated by $(K_x; K_y; K_z)$. The corresponding quasi-particle energy is determined by the dispersion relation $\frac{2}{k} = h^2 c^2 k^2 + \frac{2}{c^4}$, where the elective masses k and $\frac{2}{2}$ lead to the energies $\frac{k}{k}$ and $\frac{2}{k}$, respectively. Finally, the prime on the summation sign indicates that the mode having k = 0 should be om itted since it represents the order parameter rather than a spatial uctuation.

 $^{^{1}}$ W e call attention to the fact that in the present notation $^{2}_{2}$ denotes the variance of the ed uctuations in one of the transverse directions, whereas in refs. [10, 14] it represents three times that am ount, namely the sum of the variances in all three perpendicular directions.

In eq. (17), the symbol indicates the therm odynamic limit L ! 1 in which the quasi-particle modes form a continuum. The uctuations can then be expressed analytically in terms of the M odi ed Bessel Function K₁, as indicated. Generally, the uctuations decrease as the elective mass increases, for a xed temperature T, and $_{j}^{2} = T^{2}=12$ when the elective mass j vanishes. The global approximation $_{j}^{2} = (T^{2}=12)(1 + c^{2}=2 \text{ T})\exp((-c^{2}=T))$ is good to better than 2%.

The treatment above has been carried out under the simplifying assumption of them alequilibrium which su ces for our present purposes. However, it is interesting to note that the results can be readily generalized to non-equilibrium scenarios by simply replacing the them all averages by the corresponding spatial averages < >. In this manner, the system of equations would be closed and a conceptually simple dynam ical description emerges. However, since the direct interaction between the quasi-particles is then ignored, any relaxation can occur only via the mean eld, and the accuracy of such an approach should therefore be carefully assessed.

3.1.1 Sym m etries

It is in portant to note that the equation of motion (11) for the spatial average of the eld strength, the order parameter _, has four components, one for each of the chiral directions.

In the special case when the symmetry-breaking term vanishes, H = 0, the interaction term in (11) depends only on the magnitude of the order parameter, 0. The evolution of is then equivalent to that of a particle in a central eld in four dimensions and, accordingly, the dynamics separates into one radial and three angular modes. The rotational invariance in plies that the angular motion is massless, a manifestation of the Goldstone Theorem (see, for example, ref. [21]). The conserved currents associated with this 0 (4) invariance are the N oether iso-vector current. and iso-axial-vector current A = (related to rotations 77 = ß Ø Ø around the axis and rotations of the axis, respectively). In the general case, when H does not vanish, the m odel rem ains invariant with respect to rotations around the axis and the associated 0 (3) symmetry ensures the conservation of isospin.

We also note in this connection that the quasi-particles modes all have a nite e ective mass (except right at the critical point). This should pose no puzzle since they have always nite momenta, whereas the above modes carry no momentum.

3.2 E ective m asses

U tilizing the result (17), the coupled equations (14{15) for the e ective m asses can be solved for speci ed values of the tem perature T and the magnitude $_0$, provided that these parameters are su ciently large, and we then have $_k$? 0. It should be noted that the e ective m asses are independent of the parameter H, since the sym m etry-breaking term H is linear.²

²This simplifying feature holds only when , v, and H are considered as the primary model parameters. When H vanishes it is often custom ary to readjust the other two parameters, = $m^2c^2=2f^2=2127$ and v = f = 92 MeV, and then the elective masses change correspondingly.

In g.1 we show the resulting e ective masses $_{k}$ and $_{?}$ as functions of $_{0}$, for temperatures T up to well above critical. At any temperature, there is always a physical solution to the coupled equations (14-15) for the e ective masses when $_{0}$ v. This is easy to see from eq. (15): At T = 0, when the uctuations vanish, we have $_{?}^{2} = (_{0}^{2} v^{2})$ and so $_{?}^{2}$ vanishes at $_{0} = v$ and is positive for larger $_{0}$; an increase of T will always increase the uctuations, and hence the mass. Moreover, we always have $_{k}$?

Since the eld uctuations and the magnitude of the order parameter contribute to the eld uctuations in qualitatively similar ways, an increase of the temperature (and thus the uctuations) will permit a further decrease of $_0$, so that the point at which $_2$ vanishes is moved to ever smaller values of $_0$. The behavior of $_k$, considered as a function of $_0$, is nearly independent of temperature (as long as T is subcritical), except that each curve term inates at the point where the corresponding $_2$ vanishes. This limiting curve is indicated by the dotted curve on the interval (0;v) and it is elementary to calculate,

$${}^{2}_{0} = v^{2} \frac{5}{12}T^{2} {}^{2}_{k} v^{2} \frac{T^{2}}{12}(5 + e^{kc^{2} - T});$$
(18)

$${}^{2}_{k}c^{4} = (2v^{2} T^{2}) :$$
 (19)

W hen the tem perature reaches the critical value, $T = T_c$, there is a solution to eqs. (14-15) for all values of $_0$, and both e ective m asses vanish for $_0 = 0$. Then

 $_{j}^{2} = T_{c}^{2}$ =12 and so $T_{c}^{2} = 2v^{2}$, i.e. $T_{c} = 122.63 \text{ M} \text{ eV}$ with the adopted parameter values. The degeneracy in the masses, $_{k} = _{2}$, is a general consequence of the O (4) rotational sym metry that emerges for $_{0} = 0$ and it therefore remains as T is further increased, with the common mass value $_{0}$ increasing steadily. Since the elective mass at $_{0} = 0$ is given by $_{0}^{2}c^{4} = (6 \qquad _{j}^{2} \qquad v^{2})$, it becomes proportional to T at high temperatures, $_{0}c^{2} = 1.59 \text{ T}$ for T v.

The results displayed in g.1 have been calculated in the therm odynam ic lim it, L ! 1, in which the quasi-particle modes form a continuum. However, the systems of interest in connection with high-energy nuclear collisions have nite volum es. The quasi-particle energies are then discrete and, as a result, the elective masses are modiled. This elect is largest when the elective mass is small, i.e. near the critical conditions, because the absence of the constant mode in (17) is then most signil cant, and it generally leads to smaller masses. Figure 2 shows the tem perature dependence of the elective masses for either $_0 = 0$ or for $_0 = f$ (the value associated with the physical vacuum), as obtained either in the continuum limit or for a cubic box with a side length L of either 8 fm or 5 fm.

For $_0$ 0 and temperatures near critical, the elective masses are signily reduced. As a consequence, the associated critical temperature is increased to 154 MeV for L = 8 fm and to 172 MeV for L = 5 fm. However, for temperatures several tens of MeV above these thresholds, the elect is relatively small. The elect also disappears quickly as the order parameter is moved away from zero and by the time it reaches its vacuum value, $_{vac} = f = 92 \text{ MeV}$, there is hardly any elect for even the smallest box (L = 5 fm).

The e ect of the nite size on the uctuations in the eld strength is illustrated in g.3. The upper curves show the typical magnitude of the therm al uctuations, $\binom{2}{2}^{1=2}$, as functions of $_0$ over a range of tem peratures, for the same three box sizes as in g.2, while the low er curves show the continuum result for the contribution arising from uctuations directed along the order parameter _, $\binom{2}{k}^{1=2}$. The uctuations are remarkably independent of the volum e, except in the critical region T T_c and $_0$ 0, as just discussed. For $_0 = 0$ the O (4) symmetry implies that the uctuations are the same in all directions so that each of the low er curves starts out at half the value of the corresponding upper curve. For larger values of $_0$ the relative contribution from uctuations along _ decreases progressively, since _k grow smuch larger than $_2$ (see g.1).

3.3 Energy

It is instructive to exam ine the total energy of the system . It is obtained by integrating the energy density (8) over the volume of the box,

E (t) =
$$dr H (r;t) = < \frac{1}{2}h^2 + \frac{1}{2}h^2c^2(r)^2 + V >$$
 (20)

$$= (E_{0} + E_{qp} + V) :$$
 (21)

The total energy is conserved in time when the eld is evolved with the equation of motion (9). The energy is composed of the three basic contributions exhibited in the second expression: one arising from the time dependence of the eld strength, another associated with the spatial variation of the eld, and a third resulting from the interaction. In the last expression, these contributions have been reorganized into three di erent term s which we will now discuss in turn.

The rst term is the bare energy density, i.e. the energy density that would arise if the eld had no spatial variation, = and = ,

$$E_{0} = \frac{1}{h^{3}c^{3}} \begin{bmatrix} 1 \\ 2 \end{bmatrix}_{0}^{2} + \frac{1}{4} \left(\begin{bmatrix} 2 \\ 0 \end{bmatrix} + \frac{1}{4} \left(\begin{bmatrix} 2 \\ 0 \end{bmatrix} + \begin{bmatrix} 2 \\ 4 \end{bmatrix} \right)^{2} + \begin{bmatrix} 2 \\ 0 \end{bmatrix} = \begin{bmatrix}$$

We have here introduced $_0$ to denote the angle of disorientation, i.e. the angle between the four-dimensional order parameter __ and the __ direction. The spatial average of __ is then __ = __ ^ = __ cos __ , and the corresponding mean magnitude of the pion eld is __ = __ sin __ . In the last expression K __ denotes the kinetic part of E __ and V__ is the bare potential energy density.

The second term in (21) denotes the energy density associated with the quasiparticle degrees of freedom, in the approximation that these form a gas of independent particles,

$$E_{qp} = \frac{1}{h^{3}c^{3}} \sum_{j=0}^{X^{4}} \frac{1}{2} < (h (\theta_{t}))^{2} + (h cr)^{2} + \sum_{j=0}^{2} c^{4} - \sum_{j=0}^{2} >$$
(23)

$$= \frac{1}{h^{3}c^{3}} \sum_{j=0}^{X^{4}} \frac{1}{2} \sum_{k=0}^{X} (j_{k}) j_{k}^{(j)} j_{j}^{2} + (h^{2}k^{2}c^{2} + j_{j}^{2}c^{4}) j_{k}^{(j)} j_{j}^{2}) : \qquad (24)$$

We have here invoked the general Fourier expansions of (r) and (r),

$$(r;t) = \sum_{k}^{X} (t) e^{ik} r;$$
 (25)

$$(r;t) = \sum_{k}^{X^{*}} (t) e^{ik} r:$$
 (26)

The expansion coe cients are generally complex and are given simply in terms of spatial averages, $_{k} = \langle (r) \exp((ik r) \rangle$ and $_{k} = \langle (r) \exp((ik r) \rangle$. We note that the components with k = 0 represent the order parameter, $_{0} = _$ and $_{0} = _$. Furtherm ore, the fact that the chiral eld is real imposes the relations $_{k} = _{k}$ and $_{k} = _{k}$.

The third term in (21) corrects for the fact that the quasi-particles in fact interact via the fourth-order term in V ,

$$V = \frac{1}{4} \begin{bmatrix} 4 \\ 2 \end{bmatrix} = \frac{2}{2} \begin{bmatrix} 2 \\ 2 \end{bmatrix} = \frac{2}{2} \begin{bmatrix} 2 \\ 2 \end{bmatrix} = \frac{2}{3} \begin{bmatrix} 2 \\ 2 \end{bmatrix} =$$

The last result can be obtained by using the Gaussian approximation to express < ⁴ > in terms of squares of quadratic terms.

The energy of the system can be calculated as a function of the order parameter, assuming that the quasi-particle degrees of freedom are in them al equilibrium. The resulting energy density is

$$\underline{E}(\underline{;}) = E_{0}(_{0};_{0};_{0}) + E_{qp}(_{0}) + V(_{0}) = K_{0} + V_{T}(_{0};_{0}) : (28)$$

Here the therm al equilibrium value of the quasi-particle energy density is given by

$$E_{qp}^{(j)} = \frac{1}{k} \frac{X}{e^{k-T}} \frac{0}{1} \frac{k}{n^{3}c^{3}} \frac{(c^{2})^{4}}{2^{2}} \frac{1}{n>0} \frac{X}{n^{2}} \frac{1}{n^{2}} \frac{X}{n} \frac{1}{n^{2}} \frac{1}{n^{2}$$

for each of the four chiral directions j, with $T = {}_{j}c^{2}$. It can be expressed analytically in the continuum limit, as indicated. Furtherm ore, the therm al equilibrium value of the correction term is quite well represented by its G aussian approximation,

In the last expression in (28) the energy has been split into the kinetic energy associated with the time dependence of the order parameter, $K_0 = {}_0^2 = 2h^3c^3$, and the remainder which can be regarded as an elective potential,

$$V_{T}(_{0};_{0}) = V_{0} + E_{qp} + V :$$
 (31)

This elective potential has been plotted in g.4 for the particular situation when the order parameter _ is directed along the axis, for a range of tem peratures T. The directional dependence of V_T is simply given by the term H $_0 \cos _0 = h^3 c^3$. At zero tem perature the elective potential can only be displayed for order parameters

whose magnitude $_0$ exceed v, since only then can the elective masses be calculated, as explained in sect. 3.2: For smaller values of $_0$ some of the low-energy quasiparticle modes are unstable and so have no well-de ned therm all equilibrium. As the tem perature is increased, the domain of stability extends ever lower towards $_0 = 0$. The limiting curve connecting the end points of all the subcritical curves is determined by the condition that $_2$ vanish, implying $_k^2 = (2v^2 - T^2)$. The result is shown on the gure as the dotted curve between $_0 = 0$ and $_0 = v$.

For H = 0 (g. 4a) the elective potential has 0 (4) symmetry and so the minima form the surface of a 4-sphere; its intersections with the axis are indicated by the two arrows. The addition of the linear symmetry-breaking term (g. 4b) tilts the potential towards the direction and, consequently, there is only one minimum for any given temperature and it is located on the axis.

For all values of H, the m inim a of the elective potential move gradually towards larger values of the order parameter. In this respect, the results are similar to the behavior of the energy of nuclear matter as a function of the density (see, for example, ref. [22]). However, while the energetics alone thus favors ever larger values of either $_0$ or as the temperature is raised, the proper incorporation of the phase space (by m eans of the appropriate entropy) turns the picture around, as we shall now see.

4 The order param eter

We can now consider the distribution of the chiral eld in statistical equilibrium at a given temperature T. Since the equation of motion (9) is of second order in time, we need to specify both the eld strength (r) and its time derivative (r) $h \mathcal{Q}_t$ (r) in order to fully characterize a state. A swe have already exploited above, the quasiparticle degrees of freedom can be regarded as a Bose-E instein gas having therm al distributions determ ined by their respective elective masses, in the appropriate domain of $_0$. Their behavior can then be calculated once the order parameter has been specified.

4.1 The partition function

It is natural to start the discussion of statistical properties by considering the partition function for the chiral eld. It is obtained by sum m ing the canonical weights for all the possible states,

$$Z_{T} \qquad \sum_{z} D[;] e^{\frac{T}{T}E[;]} = \sum_{k}^{Y} d^{4}_{k} d^{4}_{k} e^{\frac{T}{T}(E_{0}+V+E_{qp})}$$
(32)

$$d^{4} d^{4} e^{T} (E_{0} + V) Z_{qp} = d^{4} d^{4} W_{T} (::) ; \qquad (33)$$

where E is the total energy of the system given in eqs. (20-21). The functional integral over all states ((r); (r)) has rst been expressed as a multiple integral over the four-dimensional Fourier amplitudes $_{k}$ and $_{k}$ (see eqs. (25-26)). If the integration

over the eld uctuations (i.e. those amplitudes having $k \in 0$) is performed, the remaining integrand is the statistical weight W_T expressing the relative probability for nding the system with a speci ed value of the order parameter. In order to derive an expression for W_T , we have replaced the correction term V (eq. (27)) by its approximate thermalaverage V, for each value (;) of the order parameter (it depends only on $_0$). The integral over the quasi-particle degrees of freedom can then be performed and yields the quasi-particle partition function $Z_{\rm qp}$.

This quantity factorizes into contributions from each of the four chiral directions, $Z_{qp} = {}^Q_{j} Z_{qp}^{(j)}$, where

Since the complex Fourier amplitudes ${}_{k}^{(j)}$ and ${}_{k}^{(j)}$ are subject to symmetry relations under k! k, as already noted above, is it more convenient to use the representation in terms of trigonom etric functions, eqs. (46) and (48). The dimension wave numbers are then fully decoupled, and for each k the integration variables are the (positive) amplitude C_k and the associated phase k, as indicated in the last expression above. The phases are unimportant here since they each merely contribute a factor of 2. Moreover, the combination $n_k = {}_kC_k^2 = h^3c^3$ can be interpreted as the average number of quanta in the mode k (see eq. (47)). Since we want to take account of the quantal nature of the quasi-particle degrees of freedom (see 5.1.1), we replace the continuous integral over n_k by a discrete sum over the possible integer values of the occupation number, thereby obtaining the standard expression for an ideal B ose-E instein gas,

$$Z_{qp}^{(j)} = \begin{pmatrix} Y_{04} & X^{d} \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & \\ & & & & \\ &$$

We have used the fact that the quasi-particle energy is additive, $E_{qp}^{(j)} = {}^{P}_{k} n_{k \ k}$. Furthermore, $f_{k} = 1 + f_{k}$ where $f_{k} = (e^{k=T} - 1)^{-1} = n_{k}$ is the mean number of quanta (in the chiral direction j) having a wave number of magnitude k, and the corresponding energy $_{k}$ is determined by ${}^{2}_{k} = h^{2}c^{2}k^{2} + {}^{2}_{j}c^{4}$.

It follows that the free energy density of the quasi-particles is $F_{qp} = \int_{j}^{P} F_{qp}^{(j)}$ with

$$F_{qp}^{(j)} = \frac{T}{m} \sum_{qp}^{(j)} = \sum_{k=1}^{N-0} h_{k} f_{k} T (f_{k} \ln f_{k} f_{k} \ln f_{k})^{i} = E_{qp}^{(j)} T S_{T}^{(j)} : (36)$$

In the resulting expression, $E_{qp}^{(j)}$ is the average therm all energy density of the quasi-particle m odes associated with the chiral direction j, eq. (29), and $S_T^{(j)}$ is their entropy density. The total entropy density at the tem perature T is then $S_T = S_T^k + 3S_T^2$, where S_T^k ($_0$) represents the entropy density of the eld uctuations along the direction of [23],

$$S_{T}^{k} = -\frac{1}{k} \int_{k}^{\infty} f_{k}^{k} \ln f_{k}^{k} f_{k}^{k} \ln f_{k}^{k} \int_{k}^{k} \ln f_{k}^{k} dk f_{k}^{$$

$$\frac{1}{h^{3}c^{3}} \int_{k^{2}}^{z} \frac{1}{2^{2}} \left(\int_{k^{2}}^{2} \frac{1}{2} \int_{k^{2}}^{z} \frac{1}{2} \int_{k^{2}}^{h} f(x) \ln f(x) f(x) \ln f(x) + \int_{k^{2}}^{1} \frac{1}{2} \int_{k^{2}}^{1} \frac{1}{2} \int_{k^{2}}^{h} f(x) \ln f(x) + \int_{k^{2}}^{1} \frac{1}{2} \int_{k^{2}}^{1} \frac{$$

and the contribution to the entropy density from uctuations in a perpendicular direction, S_T^2 ($_0$), can be calculated analogously by replacing $_k$ with $_2$.

It is evident from g.4 that the energetics favors rather large m agnitudes of the order parameter, as would be expected from the form of the interaction potential V. However, the density of quasi-particle states is higher at small values of $_0$, since the elective m assess are then smaller, for a given temperature.³ This interplay between energy and phase-space is quantiled in the statistical weight (see eq. (33)),

$$W_{T}(\underline{;}) \qquad D \begin{bmatrix} 0; \\ 0 \end{bmatrix} e^{E\begin{bmatrix} 0; \\ 0 \end{bmatrix} = T - 4} (\underline{0} -)^{-4} (\underline{0} -)$$
(39)

$$e^{T}(K_{0}+V_{0}+V_{0}+V_{0}+F_{qp}) = e^{T}(K_{0}+F_{T})$$
(40)

We have used the approximate decomposition (21) of the total energy which causes the partition function to factorize and, in the last expression, we have introduced the free energy density for the order parameter, F_T () = V_T (0; 0) TS_T (0).

The free energy density is illustrated in g.5 for a range of tem peratures above which the trend is obvious. Since the entropy density of a free gas at xed T goes up when the particle m ass is reduced, the entropy provides a restoring force towards symmetry (recall that a reduction of the order parameter $_0$ leads to smaller e ective m asses). As a result, for H = 0 and tem peratures well above T_c the free energy density then has its minimum at $_0 = 0$. As the tem perature drops, the symmetric minimum growsevermore shallow.NearT 175MeV a secondary minimum appears 50 M eV, and it becom es the lowest one at T 171 M eV. As T decreases at 🔐 further the m in imum gently approaches $_0 = v$. The abrupt change in the location of the free energy m in im um is characteristic of a rst-order phase transition. In contrast, when H takes on a realistic nite value (see eq. (6)) the minimum of F_T is always located on the positive part of the axis and it moves smoothly outwards as the system is cooled, with no drastic change at any tem perature. The resulting behavior is shown by the solid curve in q.6.

4.2 Distributions

The preceding analysis in plies that the statistical weight $W_T(\underline{;})$, which is defined on the eight-dimensional phase space of the order parameter, in fact depends on only three quantities: the speed $_0 = \underline{j}$ \underline{j} the magnitude $_0 = \underline{j}$ \underline{j} and the disalignment angle $_0$. Moreover, since the entropy depends only on the elective masses, which in turn are determined solely by $_0$, the dependence of the free energy on the direction of ______ arises only through the symmetry-breaking H term in the interaction V (eq. (2)). Consequently, the corresponding projected probability distribution factorizes,

$$P_{T}(_{0};_{0};_{0}) = P(_{0})P(_{0})P(_{0};_{0}) = \frac{2}{3} \frac{2}{3} \frac{2}{3} \sin^{2} (_{0})W_{T}(_{0};_{0}) = \frac{2}{3} \frac{2}{3} \sin^{2} (_{0})W_{T}(_{0};_{0};_{0}) = \frac{2}{3} \frac{2}{3} \sin^{2} (_{0})W_{T}(_{0};_{0};_{0}) = \frac{2}{3} \frac{2}{3} \sin^{2} (_{0})W_{T}(_{0};_{0};_{0}) = \frac{2}{3} \frac{2}{3} \frac{2}{3} \sin^{2} (_{0})W_{T}(_{0};_{0};_{0}) = \frac{2}{3} \frac{2}{3} \frac{2}{3} \frac{2}{3} \sin^{2} (_{0})W_{T}(_{0};_{0};_{0}) = \frac{2}{3} \frac{2}$$

 $^{^{3}}$ The importance of the quasi-particle entropy in determining the statistical properties of the model was already emphasized by Bardeen and Moshe [24] a decade ago.

Here $_{0}^{3}$ and $_{0}^{3}$ are the Jacobians associated with the transform ation from the fourdimensional vectors _ and _ to their magnitudes $_{0}$ and $_{0}$, respectively, and \sin^{2}_{0} is the Jacobian arising from the three-dimensional nature of the space perpendicular to the order parameter _. Furthermore, the three normalized probability densities are given by

P (₀) = N
$$_{0}^{3} \exp\left[\frac{1}{h^{3}c^{3}}\frac{2}{2T}\right];$$
 (42)

$$P(_{0}) = N _{0}^{3} \exp[((\frac{1}{T}V_{T}(_{0}; _{0} = 0) S_{T}(_{0}))]; \qquad (43)$$

$$P(_{0}; _{0}) = N(_{0}) \sin^{2} _{0} \exp\left[\frac{1}{h^{3}c^{3}} \frac{H}{T} _{0}(1 \cos _{0})\right]:$$
(44)

For the normalization constants we have N $^{1} = 2 ^{2}$, with the convenient abbreviation

 $Th^{3}c^{3}=$, and $N^{1}=(=)$ [() exp(), with $H_{0}=$.

The kinetic part, P ($_0$), is the projection of an isotropic four-dimensional normal distribution having a total variance of $_0^2 = 4$. The average kinetic energy associated with the time evolution of the order parameter is then $_q \frac{K}{9} = 4T$, as would be expected in four dimensions. The mean speed is $_0 = 9 = 8$.

Considered as a function of the four-dimensional order parameter _, the probability density has a maximum where the free energy has a minimum, as is evident from eq. (39). However, the jacobian factor 3_0 associated with the projection from _ to $_0$ increases the most probable value of the magnitude $_0$. This elect depends on volume, as is illustrated in g.6. The results for a nite box with L = 8 fm (dashed curve) are nearly identical to those obtained for L ! 1, while L = 5 fm (dots) leads to a more significant increases in $_0$ at the higher temperatures. The most rapid increase of $_0$ occurs near T = 200 M eV which is well above the critical temperature T_c (arrow) at which the elective masses drop to zero.

The directional distribution P ($_0$; $_0$), which depends param etrically on the m agnitude $_0$, grow s broader at high tem perature where $_0$ decreases. In the tem perature range considered here, up to several hundred M eV, P ($_0$) is well approximated by the projection of a three-dimensional norm all distribution with a total variance of $3^{=}$, and the average magnitude of the disalignment angle $_0$ is then $_0$ $8^{=}$.

A lthough the locations of the m inim a in the free energy appear qualitatively different when comparing H = 0 with H > 0 (gs. 5a and 5b, respectively), the corresponding curves for the most probable $_0$ are rather similar when nite volumes of nuclear size are considered, due to statistical uctuations. This is also illustrated in g. 6. The thin curve is analogous to the solid curve and shows the dependence of $_0$ in the therm odynam ic limit. (Recall that the vacuum value is $_0 = v = 86:71 \text{ M eV}$ for H = 0.) At T 171 M eV the value changes abruptly. The open dots are analoguos to the solid dots and show the most probable $_0$ for the box having L = 5 fm. The bias by the jacobian factor $_0^3$ (see eq. (43)) causes the behavior to become quite similar to that obtained for H > 0. In particular, there is no discontinuity near T 171 M eV where the minimum in the free energy jumps between $_0 = 0$ and

56 M eV.W e nally note that for relatively low temperatures the Jacobian causes the most probable value of $_0$ to slightly exceed the vacuum value.

4.2.1 Distribution of the order parameter

For any nite volume , the order parameter the exhibits uctuations around its most probable value. The full width at half maximum of the $_0$ distribution is shown by the horizontal bars in g.6 for the smallest box (L = 5 fm) where the uctuations are the largest. The relative smallness of the uctuations indicates that the order parameter is distributed within a rather limited range of values. This feature is further illustrated in g.7 which depicts the entire distribution, P ($_0$), over a range of tem peratures, for a cubic box with side length L = 8 fm, which is our standard scenario. The solid curves in g.7 have been obtained by scaling the continuum results to the nite volume. The dashed curves show the e ect of properly quantizing the problem (i.e. summing over the discrete modes rather than integrating). The e ect is very small, because the distributions are peaked well inside the respective dom ains of stability.

For small temperatures, the distribution is narrow by peaked near the vacuum value $v_{ac} = f$ (indicated by the arrow). As the temperature is increased, the distribution broadens and gradually begins to move inwards towards smaller values of $_0$. The width of the distribution increases from zero at T = 0, exhibits a maximum near T = 220 MeV, and then slow by shrinks as T + 1. The somewhat counterintuitive decrease of the uctuations at high temperatures is due to by the fact that the grow ing thermal uctuations cause the interaction to become progressively more repulsive, causing the elective potential V_T to become ever more con ning.

We recall that for subcritical temperatures, $T < T_c$, our treatment can only be carried out above a temperature-dependent minimum value of $_0$. As it turns it, this principal limitation is of little practical import. For example, for T = 80 M eV (the lowest temperature shown in g. 7) the distance from the centroid (at 91 M eV) to the boundary (at 69 M eV) is over ten times the dispersion of the distribution (2 M eV). Thus, for any temperature, the distribution P ($_0$) is su ciently narrow to make incursions into the respective unstable regime extrem ely unlikely.

A more global in pression of the statistical distribution of the order parameter can be obtained by considering a contour diagram of the projected probability density $P(_{0};_{0}) = P(_{0}) P(_{0};_{0})$. Such a plot is displayed in q.8 for a box with L = 8 fm. The abscissa is the projection of the order parameter onto the axis, $_{0}$ $^{=}$ $_{0} \cos _{0}$, and the ordinate is the magnitude of its perpendicular com ponent, 0 $_0 \sin _0$. For each tem perature, the dots show the location of the maximum and the half-maximum contours are traced out. The solid contours and dots refer to the continuum treatment, whereas the dashed contours and the open dots (shown for two tem peratures only) are obtained with a quantized treatment, which is seen to have little e ect. The proles in q.7 are the projections of P ($_0; _0$) onto the magnitude $_0$. It is evident from the above results that the O (4) symmetry is far from restored at the temperatures expected in the planned high-energy nuclear collisions. Instead the order parameter is in fact distributed within a fairly limited dom ain, with both its magnitude $_0$ and its angle of disalignment $_0$ exploring only rather narrow ranges.

4.2.2 Distribution of the e ective masses

The elective m asses are functions of the magnitude of the order parameter, $_k(_0)$ and $_2(_0)$. Consequently, the thermal uctuations in the order parameter will cause the elective m asses to uctuate as well. Since their distributions may be of some interest (see sect. 5.3), we consider brie y the corresponding probability densities for the elective m asses,

$$P_{?}(c^{2}) = d^{4} P() (_{?}(_{0})c^{2} - c^{2});$$
 (45)

and analogously for P_k (c^2), where $P(\underline{)} = P(\underline{0})P(\underline{0}; \underline{0})$ is the probability density for the order parameter __. In order to gain an impression of these distributions, we show in g.9 the most probable elective masses as functions of the tem perature.

The transverse elective mass $_{?}$ increases steadily from its free value m as the temperature is raised, whereas the parallel elective mass $_{k}$ drops by nearly fly per cent before nally turning upwards. For temperatures above T 300 MeV the two elective masses are practically degenerate and gently approach their asymptotic form

 c^2 1:59 T. We note that the elective m asses always exceed the temperature, so it is never reasonable to ignore the elective m asses. We also note that the transverse m ass has practically no strength near the free pion m ass, until the temperature is below T_c (see g.13).

The parallele ective mass $_{k}$ exhibits a pronounced minimum near T 240 M eV. It is in this region that the order parameter exhibits its most rapid evolution with T (see g.6) and the temperature at which the minimum in $_{k}$ occurs may be interpreted as the elective critical temperature of the model [25].

4.3 Sam pling of the order param eter

The above analysis provides a convenient basis for sampling the order parameter in accordance with the statistical weight W $_{\rm T}$ (_;_) given in eq. (39), and we describe brie y how this can be accomplished in a manner that is both quick (i.e. the computationale ort is small) and e cient (i.e. no e ort is expended on rejection).

As noted already, the time derivative _____ is governed by a four-dimensional normaldistribution which is isotropic and entirely decoupled from the other degrees of freedom (see eq. (42)). Thus it is elementary to sample this quantity.

The most complicated sampling concerns the magnitude $_0$, due to the intricate structure of its probability distribution, as discussed above. However, the numerical e ort required is quite modest. The most e cient method requires a precalculation of the e ective masses as functions of $_0$, for the particular T of interest. This is quickly done by proceeding as described in sect. 3.2. The $_0$ -independent part of the e ective potential, V_T ($_0$;0), can then be obtained together with the corresponding entropy S_T ($_0$). In elect, the probability distribution for $_0$ can be pretabulated, P ($_0$) (see eq. (43)), and it is then a numerically trivial task to sample $_0$.

Once the magnitude $_0$ has been selected, it is straightforward to sample the disalignment angle $_0$, using either the exact form (44) or its G aussian approximation.

In order to orient ____ in the subspace, there remains the task of selecting the remaining 0 (3) spherical angles $\#_0$ and \prime_0 , upon which the order parameter is given by = $(_0 \cos _0; _0 \sin _0 \sin \#_0 \cos'_0; _0 \sin _0 \sin \#_0 \sin'_0; _0 \sin _0 \cos \#_0)$.

5 The quasi-particles

We turn now to the discussion of the quasi-particle degrees of freedom associated with the spatial variations of the chiral eld, (r). Once the magnitude of the order parameter, $_0$, has been chosen (see sect. 4.3), the quasi-particle degrees of freedom are fully characterized, via the elective masses $_k(_0)$ and $_2(_0)$, and it is then possible to sample them them appropriately.

5.1 Sam pling

On the basis of the above developments, it is possible to devise a simple and e cient method for performing the statistical sampling of the chiral eld, thus putting the initialization of dynamical simulations on a formally sound basis.

Since the di erent quasi-particle m odes can be regarded as e ectively decoupled, the sam pling is best done by m aking an expansion into the elementary m odes,

$$_{k}(\mathbf{r};t) = \frac{2}{k} \int_{k}^{\frac{1}{2}X} O_{k} O_{k} O_{k} C_{k}^{k} O_{k} C_{k}^{k} O_{k} C_{k}^{k} O_{k} O_{k}$$

and similarly for the three transverse chiral components $_{?}(r;t)$. Here the energy $_{k} = h!_{k}$ is determined by the K lein-G ordon dispersion relation, $_{k}^{2} = h^{2}c^{2}k^{2} + {}^{2}c^{4}$ (with being the appropriate elective mass, $_{k}$ or $_{?}$). The phase $_{k}$ is random in the interval (0;2) and is thus trivial to sample. Furthermore, the real (and positive) amplitude C $_{k}$ can be related to the number of quanta n_{k} by considering the energy carried by the mode,

$$E_{k} = n_{k-k} = \frac{\frac{2}{k}}{h^{3}c^{3}}C_{k}^{2} \quad) \quad C_{k} = h^{3}c^{3}\frac{n_{k}}{k}^{\frac{1}{2}} :$$
(47)

We have here on itted the zero-point energy [26], thereby elim inating the associated ultraviolet divergence. A lthough not quite correct, this approach is justiled a posteriori by the apparent good quality of the resulting approximate treatment. (The dynamical tests discussed in sect. 6 are here very important, since the statistical samples would not remain stationary under time evolution if the treatment were substantially wrong.)

Thus the problem has been reduced to sampling the number of quanta n_k . Since the probability for nding a particular number of quanta in the mode k is given by P $(n_k) = (1 \exp(k=T))\exp(n_k =T)$, it is elementary to sample the integer n_k appropriately.⁴ W e note that the therm allowerage of n_k is equal to the occupancy f_k employed in the calculation of the entropy, $n_k = f_k$.

 $^{^{4}}$ T he corresponding algorithm for this task is very simple because n_{k} can be regarded as counting the number of successive times the sampling of a standard random number (i.e. uniform on (0;1)) yields a value below exp($_{k}$ =T).

Once the amplitudes and phases have been selected, the expansion (46) readily yields the initial value of the eld uctuations, (r;0). The corresponding conjugate momentum, hQ, readily follows since (46) in plies

$$_{k}(r;t) = \frac{2}{k} \int_{k}^{\frac{1}{2}X} \int_{k}^{k} C_{k}^{k} \sin(k r + kt + k) :$$
(48)

The entire state of the chiral eld, ((r); (r)), has then been selected at the time t = 0. W hen the equations of motion are propagated by a leap-frog method, the eld strength (r) is calculated at the times $t_n = n$ twhile the momentum (r) is obtained at the intermediate times. The appropriate initial (r) can then easily be obtained by evaluating the expansion (48) at t = t=2, after C _k and _k have been selected.

Finally, since the sampling has been done in a system aligned with the order parameter _ (in which the mass tensor is diagonal), an O (4) rotation of the sampled eld con guration is required in order to express the state with respect to the chiral directions. This is readily accomplished on the basis of the angles ($_0; \#_0; '_0$) specifying the direction of .

5.1.1 Occupation numbers

Since the quasi-particles represent bosonic modes, it is useful to know how large the occupancies can become. The thermal occupation numbers are given by $n_k = f_k = 1 = (\exp(k=T))$, and since $\frac{1}{2} < \frac{1}{k}$ in any given scenario the largest occupancies occur for the transverse modes, $f_k^2 > f_k^k$. Moreover, for a given temperature T, f_k is largest when the momentum vanishes, $f_0 = 1 = (\exp(c^2 = T))$, which then provides an upper bound on f_k (recall that k > 0 for the quasi-particles modes).

Considered as functions of T, the bounds f_0^k and f_0^2 start out from zero, display maxima well above T_c, and then drop o towards a common constant value at high temperatures. The (common) limiting occupancy is $f_0 = 0.26 \text{ since } c^2 = 1.59 \text{T}$ when T ! 1. (This feature is a direct consequence of the repulsive self-interaction of the chiral eld and is in marked contrast to the ever increasing occupancy characteristic of free bosons in a thermal bath.) The maximum values attained by f_0^k and f_0^2 are about 0.38 and 0.48 and occur at approximately a temperature of 265 and 235 M eV, respectively. Since these values are well below unity, the system is never very degenerate.

Nevertheless, there is an important advantage to using Bose-E instein rather than Boltzm ann statistics for the eld uctuations, as we will now discuss. When classical statistics is used, the occupation probability is $n_k = T = k$ when k = T and therefore the total quasi-particle density grows as the square of the upper limit on k, a manifestation of the Rayleigh-Jeans divergence. Such a description would be entirely wrong in the present context and, moreover, it would be numerically ill-behaved. By contrast, the quantal occupation probability falls of exponentially and the density of quanta is nite. By adopting the Bose-E instein treatment, we then get a much more realistic form aldescription of the statistical properties of the system and, in addition,

the num erical treatm ent becom es straightforward. O fourse, if a conguration sam – pled on the basis of the quantal statistics is propagated for a su cient length of time, it will eventually exhibit classical features, since the equation of motion is entirely classical. It is therefore fortunate that our treatm ent is expected to be applied only to processes that are far faster than the time scales associated with the reversion to classical statistics. In particular, for the form ation of disoriented chiral condensates in high-energy collisions the relevant time scales are of the order of a few fm =c, while our num erical studies exhibit no ultraviolet run-away for at least several tens of fm =c.

5.2 Correlation function

It is particularly interesting to calculate the correlation function of the chiral eld since this quantity determ ines the spectral distribution of the em itted eld quanta.

The density matrix associated with the quasi-particle degrees of freedom is a 4 - 4 tensor,

C
$$(r_{12};t_{12})$$
 $(r_1;t_1)$ $(r_2;t_2)$: (49)

The average is over the ensemble of system s considered, in the present case a therm all ensemble held at the temperature T. Since a system in equilibrium is invariant in time, the correlation function depends only on the time dimension $t_{12} = t_1 - t_2$. A nalogously, the translational symmetry of the scenario implies that the spatial dependence is via the separation $r_{12} = r_1 - r_2$. Moreover, to the extent that there is invariance under spatial rotations, only the magnitude $r_{12} = jr_{12}j$ enters. In principle, these spatial symmetries are broken when a nite box is considered but the elect is insignicant and can be disregarded in the present study.

Utilizing the expansion (46), it is elementary to show that the correlation tensor C is diagonal with the elements C_k and C_2 , where

$$C_{k}(r;t) = \frac{h^{3}c^{3}X}{k} \frac{01}{k} \frac{\cos(k r kt)}{e^{k} 1} \frac{1}{2^{2}} \frac{hc}{r} \frac{2}{k} \frac{1}{e^{2}} d \frac{\sin kr}{e 1} \cos kt; \quad (50)$$

with ${}^2_k = h^2!{}^2_k = h^2k^2c^2 + {}^2_kc^4$. An analogous expression holds for C₂ (r;t). Since $h^3c^3f_k = {}_k$ is equal to the therm allowerage A^2_k , we recognize that the result (50) corresponds to the expression (146.10) given in ref. [27].

The usual correlation function is the trace of C ,

C
$$(r_1;t_1)$$
 $(r_2;t_2) = trC = C_k(r_{12};t_{12}) + 3C_2(r_{12};t_{12}) : (51)$

Its overall magnitude is set by its value at zero, which is simply the corresponding uctuation in the eld strength, C (0;0) = 2 . It is therefore convenient to divide C by this quantity and so de ne the reduced function C₁₂ C (r_{12} ; t_{12}) = 2 . Since C₁₂ is unity when $r_1 = r_2$ and $t_1 = t_2$, it expresses the space-time attenuation of the correlation between the eld-strength uctuations at di erent space-time points. In general, C₁₂ ($l=r_{12}$) exp($cr_{12}=h$) in the limit of large separations, r_{12} ! 1, so that h = c provides a simple measure of the correlation length. In the special case when the eld-strength uctuation length. In the special case

on analytical form, C_{12} (3=) (coth 1=), where $T_1 \underline{x}$ =hc. In this extreme case, the correlation function falls o only as $1=r_{12}$.

Figure 10 shows the reduced correlation function C_{12} obtained at the critical tem – perature, $T = T_c$. Its appearance depends on the m agnitude of the order parameter, 0, through the elective m asses. As $_0$ is increased from zero to its vacuum value f = 92 MeV, the corresponding elective m assess increase from zero to their free values and the correlation function falls oliver more rapidly. The correlation function thus exhibits a signilicant sensitivity to the order parameter. We note in particular that for the most probable value, $_0$ 87 MeV, the attenuation of C_{12} is considerably faster than for $_0 = 0$ and its width has dropped by over a factor of two. It is therefore in portant to take proper account of the order parameter when calculating the correlation function.

Figure 11 shows how the reduced correlation function evolves with temperature when the most probable magnitude of the order parameter is employed. At high tem – peratures the elective masses grow nearly in proportion to T so then the correlation length tends to zero, as is borne out by the steady shrinking of C_{12} . For temperatures below critical, the eld uctuations are predom inantly associated with the transverse modes, since those have the smallest elective mass, $\frac{2}{2}$, and the correlation length grows ever larger.

It is common to characterize the system by the \correlation length", de ned as the rst moment of the equal-time correlation function which diverges when ! 0. For the present discussion, it is more convenient to characterize C_{12} by its full width at half maximum for equal times, $_{12}$, since this quantity is always easy to extract, even when the mass vanishes. Figure 12 shows this measure of the correlation length as a function of temperature, using the most probable value of $_0$. For temperatures near and below T 200 M eV the dominant uctuations are perpendicular to the order parameter since the corresponding elective mass is relatively small. For higher temperatures the asymptotic regime is approached where the chiral symmetry is approximately restored and the uctuations are similar in all four chiral directions.

It is sometimes of interest to also consider correlation functions involving $\,$, the time derivative of the local eld strength [11]. Those can be obtained in a similar manner.

5.3 Radiation spectra

The evolving chiral eld m ay give rise to mesonic radiation, in analogy with the emission of photons by a time dependent electrom agnetic eld. The rate for production of a eld quantum having energy E and momentum p is proportional to the square of the corresponding Fourier am plitude [28, 29, 30],

$$E \frac{dN_{j}}{dp} \qquad dr \quad dt_{j}(r;t) = \frac{i}{h} (p r E t)^{2} : \qquad (52)$$

Here j denotes the particular O (4) direction considered so that j = 0 gives rise to isoscalar -like m esons and j > 0 represent three components of the isovector pion-like m esons (with j = 3 corresponding to $_0$, say).

A uniform system in equilibrium has both temporal and spatial invariance and the speci c radiation rate, $_{j}(p; E)$ (i.e. the production per unit volum e and per unit time), is then given by the Fourier transform of the correlation function [2],

$$_{j}(p;E) = \frac{1}{t_{0}} E \frac{dN_{j}}{dp} = dr^{2} dt C_{j}(r;t) e^{\frac{i}{h}(p r Et)} C_{j}(p;E); (53)$$

where is the volume of the system and t_0 is the time interval considered. The Fourier transform of the correlation function is easy to obtain in the continuum limit,

$$C_{j}(p; E) = h^{3}c^{3}\frac{1}{e^{E}-1} (m^{2}c^{4} - \frac{2}{j}c^{4});$$
 (54)

where we have assumed that the energy and momentum of the radiated mesons are related by E 2 = p^2c^2 + m $^2c^4$.

The above result holds for a speci ed value of the order parameter $_0$ which determ ines the elective mass $_j$. As we have discussed, the order parameter has in general a statistical distribution, P ($_0$), giving rise to corresponding distributions of the elective masses, P $_k$ () and P $_2$ () (see sect. 4.2.2). The resulting production rate can then be obtained by integrating over the appropriate therm alm ass distribution. Thus, for example, the speci c production rate for $_0$ m esons is

$$\circ (p; E) \qquad \overset{Z}{d^{2}c^{4}} P_{2} (^{2}c^{4}) C_{2} (p; E) = h^{3}c^{3} \frac{P_{2} (m^{2}c^{4})}{e^{E} 1}; \qquad (55)$$

and the rates for the charged pions are sim ilar.⁵ This result is easy to interpret: The spectral distribution of the radiated m esons is of them allows E instein form, with the tem perature given by the value characterizing the source itself, and the overall norm alization of the radiation rate is proportional to the probability that the space and time evolution of the eld matches the particular dispersion relation for the type of m eson considered.

As an illustration, g. 13 displays the strength function P_2 (m² c⁴) obtained by evaluating eq. (45) for = m as a function of the source temperature. As the temperature is decreased towards zero, the centroid of the $_2$ distribution moves down towards the free value m and at the same time its width keeps shrinking towards zero (see g. 9). Through the supercritical regime the pion strength then grows approximately exponentially. Then a plateau is reached where the increase caused by the approach of the centroid to m is counterbalanced by the decrease due to the shrinkage. As a result, the strength is nearly constant from 90 to 30 M eV. Finally, after the centroid has practically reached m, the free strength exhibits a rapid rise as the $_2$ distribution approaches a function. On the basis of this result, one would expect radiation of free pions to be unim portant until the tem perature has dropped below T_c .

⁵The rates considered here pertain to the idealized scenario of a macroscopically uniform system (possibly enclosed in a periodic box), and hence they di er from those describing the emission from a nite source into the surrounding vacuum. In particular, the familiar kinematic enhancement of the faster-moving ejectiles is absent.

6 Dynam ical tests

Once the initial state has been prepared, for example by means of the statistical sampling described above, the chiral eld may be propagated in time by means of the equation of motion (9), which is straightforward to implement in either x or k representation.

It is possible to exploit the dynam ical evolution to test the validity of our approxim ate statistical treatment. If the system is ergodic, as would be expected because of its non-linear interaction, a dynam ical trajectory will explore the space of possible

eld con gurations in accordance with the appropriate m icrocanonical weight. Conversely, an ensemble of eld con gurations that has been sampled statistically should not exhibit any change under time evolution. These features provide a convenient m eans for checking our treatment and we give two illustrations below.⁶

6.1 A verage eld strength

Perhaps most vividly, we show in g.14 the early trajectories of the order parameter for a sample of eight congurations, considering a box with L = 8 fm and either T = 200 MeV or T = 240 MeV. The dashed contours are those already given in g. 8 indicating where the projected probability density has fallen to half its maximum value and the centroids are indicated as well (open dots). For each individual trajectory, the initial location is indicated by the solid dot and the attached solid curve shows the trajectory up to t = 1 fm = c. The fact that the initial points reject the calculated statistical distribution provides an elementary test of the num erical sampling algorithm s. Less trivial is the fact that the dynam ical trajectories indeed appear to explore the region predicted by the approxim ate statistical distribution.

This correspondence can be made more quantitative by studying how the distribution of the order parameter evolves in the course of time. This analysis is illustrated in g.15. The approximate distribution P ($_0$) given in (43) is indicated by the solid curves for T = 180 and T = 240 M eV. Forty individual systems have then been prepared by sam pling their eld con gurations as described above and they have subsequently been propagated by the equation of motion (9) up to the time t = 10 fm =c. In the course of the evolution, the value of $_0$ is extracted at regular intervals and binned into slots that are 5 M eV wide. In this manner the time-averaged distribution of $_0$ can be determined and the dashed curves display the result (which is not sensitive to an increase of either the maximum time or the sample size). The overall agreem ent with the initial distribution is very good. There is generally is slight shift outwards, am ounting typically to 1-2 M eV, which suggests that our approxim ate therm aldistributions are centered at correspondingly too low values of $_0$. Consequently, for applications where such a relatively m inor displacem ent is unimportant, the test dem onstrates the validity of our approxim ate m ethod for obtaining the statistical properties of the model.

 $^{^{6}}$ It is important to recognize that whereas the statistical properties have been obtained by assum – ing that the quasi-particle degrees of freedom are electively decoupled, no such assumption is being made in the dynamics, since the trajectories are obtained by solving the full equation of motion (9).

6.2 Field uctuations

Figure 16 displays the correlation function for the pion components, C_{12} (s₁₂), for a box with L = 5 fm prepared at T = 240 M eV. The short-dashed curves show the correlation function obtained on the basis of ten sampled con gurations, and the long-dashed curves then indicate the corresponding result after those systems have been evolved up to the time t = 10 fm =c. The two curves that go up again have been obtained by aligning the relative separation r_{12} along one of the cartesian axes, while it is directed diagonally for the other two curves (the periodicity is then $\frac{1}{3L}$ and so their eventual rise is only barely visible). For reference is shown the exact them al correlation function for either the nite box considered (solid curves) or the continuum limit (dots). W hile this latter curve tends to zero for large separations (and in fact falls o m onotonically), the correlation function for a nite box drops to a negative value, because its spatial average m ust vanish.

The correlation function remains remarkably invariant in the course of time. This indicates that our treatment, including the sampling procedure, in fact yields a good approximation to the correlated eld uctuations. As a quantitative measure, one may consider the full width at half maximum $_{12}$. The continuum value is 1.33 fm, slightly larger than the thermal result for the nite box, 1.30. For the sample of ten initial con gurations we nd 1.256 and 1.271 for the cartesian and diagonal directions, respectively, which have evolved into 1.263 and 1.290 at t = 10 fm = c. So there is no signi cant change in the width of the correlation function.

7 Concluding rem arks

The present work was motivated by the current interest in disoriented chiral condensates, particularly the various dynam ical simulations carried out with the linear model [8, 9, 11, 13, 14, 16]. Those calculations follow the non-equilibrium evolution of the cooling chiral eld in order to ascertain the degree to which coherent dom ains develop. Since the dynam ics is inherently unstable, with the low-momentum modes experiencing rapid amplication, one may expect a signi cant sensitivity of the results to the initial conditions, with a commensurate degree of di culty regarding their interpretation. Consequently, caution is required when characterizing the ensemble of initial eld con gurations employed.

In order to provide a useful fram ework for this aspect of the problem, we have explored the statistical properties of the linear model, by conning the system to a rectangular box held at a xed temperature. A lthough this problem can be treated exactly [31], we have found it preferable to linearize the equations of motion since our view is towards practical calculations. The resulting treatment then becomes very simple and appears to be su ciently accurate in the present context.

The problem separates into one concerning the spatial average of the eld, the order parameter, and another dealing with the eld uctuations, referred to as the quasi-particle degrees of freedom. The latter are described approximately in terms of elective masses that depend on both the order parameter and the temperature, but

are independent of the symmetry breaking H term; these were presented in g.1.

The partition function then takes on a corresponding separable form and, as a consequence, it is possible to develop a simple method for performing a statistical sampling of the thermal equilibrium eld con gurations, including their time derivatives, at any temperature. This result, having a clear physical basis, is expected to be directly useful as practical means for initializing the dynam ical simulations of the chiral eld of the type carried out recently by several groups [8, 9, 11, 13, 14, 16], thus making it easier to interpret the num erical results.

M oreover, our speci c illustrations provide useful insight into the equilibrium properties. In particular, it appears to be unrealistic to start the order parameter o with a value equal to zero. Indeed, at the critical tem perature the most probable order parameter is much closer to its vacuum value f than to zero. The relation-ship between tem perature and order parameter was summarized in g.6 and a more global impression of the distribution of the the order parameter (including its degree of m isalignment) can be gained from the contour plots in g.8.

Since the order parameter is thus very unlikely to vanish, the e ective quasiparticle masses remain nite. Consequently, the statistical equilibrium distribution remains well behaved at all temperatures and the change from the \restored" phase to the norm allone is fairly gradual. However, the nite size generally reduces the e ective masses, thereby bringing the system closer to criticality.

O focurse, the statistical properties are ofm ost practical interest at the relatively high tem peratures characteristic of the initial stage of the high-energy collision. Once the chiral eld has been initialized accordingly, any instabilities and associated am plications will be autom atically included in the dynam ical propagation and the system can generally be expected to quickly move out of equilibrium. The equilibrium results can then provide a meaningful reference against which to analyze the deviation from equilibrium at any stage in the dynam ical relaxation process.

A dditionally, we illustrated brie y the equilibrium form of the correlation function which is an object of prim ary interest. Indeed, it is the correlation length that properly expresses the \dom ain size" governing the conjectured anom abus pion radiation. E sæntially, what one would expect to see at the end is a stretched version of the initial correlation function since the long wave lengths are the most unstable and so will contribute in an ever larger propertion. This underscores the importance of starting out with chiral elds having physically reasonable correlation properties. To illustrate the use of the correlation function, we derived the rate at which real pion m esons are created by the eld, and subsequently we calculated the dependence of the free pion strength on the tem perature of the system.

Finally, we sought to assess the degree of validity of our approach by subjecting sampled eld con gurations to the exact time evolution. This convenient means of testing suggested that the approximate treatment is of su cient accuracy to be of practical utility. We therefore anticipate that it may not use in simulation studies, such as those exploring disoriented chiral condensates in high-energy collisions.

The author is pleased to acknow ledge helpful discussions with m any colleagues, including P.Bedaque, LP.C semai, S.G avin, J.I.K apusta, Y.K luger, V.K och, L.

McLerran, R.D. Pisarski, R. Vogt, and X.N. Wang.

References

- [1] K.Rajagopal, in Quark-Gluon Plasma 2, ed.R.Hwa, World Scientic (1995)
- [2] S.Gavin, Nucl. Phys. A 590 (1995) 163c
- [3] A A. Anselm, Phys. Lett. B 217 (1989) 169;
 A A. Anselm and M. G. Ryskin, Phys. Lett. B 266 (1991) 482
- [4] J.-P. B laizot and A. Krzywicki, Phys. Rev. D 46 (1992) 246
- [5] K.Rajagopaland F.W ilczek, Nucl. Phys. B 399 (1993) 395
- [6] C M G.Lattes, Y.Fujim oto, and S.Hasegawa, Phys.Rep. 65 (1980) 151;
 JR.Ren et al., Phys.Rev.D 38 (1988) 1417;
 L.T.Baradzei et al., Nucl. Phys.B 370 (1992) 365
- [7] K L.Kowalski and C C. Taylor, Report CW RUTH-92-6 (1992);
 JD.Bjorken, K L.Kowalski, and C C. Taylor, Report SLAC-PUB-6109 (1993)
- [8] K.Rajagopaland F.W ilczek, Nucl. Phys. B404 (1993) 577
- [9] S.Gavin, A.Gocksch, and R.D.Pisarski, Phys. Rev. Lett. 72 (1994) 2143
- [10] S.Gavin and B.M uller, Phys. Lett. B 329 (1994) 486
- F.Cooper, Y.K luger, E.M ottola, and JP.Paz, Phys. Rev. D 51 (1995) 2377;
 Y.K luger, F.Cooper, E.M ottola, JP.Paz, and A.Kovner, Nucl. Phys. A 590 (1995) 581c
- [12] J.P.B laizot and A.K rzywicki, Phys. Rev. D 50 (1994) 442
- [13] Z.Huang and X.-N.W ang, Phys.Rev.D 49 (1994) 4335
- [14] M.Asakawa, Z.Huang, and X.-N.W ang, Phys. Rev. Lett. 74 (1995) 3126
- [15] D.Boyanovsky, H.J. de Vega, and R.Holm an, Phys. Rev. D 51 (1995) 734
- [16] LP.Csemai and IN.M ishustin, Phys. Rev. Lett. 74 (1995) 5005
- [17] S.M rowczynski and B.M uller, Phys. Lett. 363B (1995) 1
- [18] M.Gell-Mann and M.Levy, Nuovo Cimento 16 (1960) 705
- [19] M.Aguilar-Benitez et al., Phys. Rev. D 50 (1994) 1173
- [20] G.Baym and G.Grinstein, Phys. Rev. D 15 (1977) 2897
- [21] C. Itzykson and J.B. Zuber, Quantum Field Theory, McGraw-Hill, New York (1980)
- [22] E.M edeiros de Lim a and J.R andrup, Nucl. Phys. A 529 (1991) 115

- [23] M. Toda, R. Kubo, and N. Saitô, Statistical Physics I, Springer, New York (1992)
- [24] W A.Bardeen and M.Moshe, Phys. Rev. D 34 (1986) 1229
- [25] R D . Pisarski, Phys. Rev. D 52 (1995) R 3773
- [26] J.I.Kapusta, Finite-tem perature eld theory, Cambridge University Press, New York (1989)
- [27] LD.Landau and EM.Lifshitz, Statistical Physics, Pergamon, New York (1980)
- [28] D.Hom and R.Silver, Ann. Phys. 66 (1971) 509
- [29] M.Gyulassy, S.K.Kau mann, and L.W. Wilson, Phys. Rev. C 20 (1979) 2267
- [30] R D. Amado, F. Cannata, J.P. Dedonder, M.P. Locher, and Bin Shao, Phys. Rev. Lett. 72 (1994) 970
- [31] J.Bernstein and S.Dodelson, Phys. Rev. Lett. 66 (1991) 683





The elective masses $_{k}$ (solid) and $_{?}$ (dashed), as functions of the magnitude of the order parameter, $_{0}$, for a range of temperatures: $T = 0, 40, 80, 100, 122.63 (= T_{c})$, 160, 200, 240, 300, 400, 500 M eV, calculated in the therm odynamic limit where the box size is large, L ! 1. For a temperature above T_{c} , the two elective mass curves start out at $_{0} = 0$ with degenerate values, whereas below T_{c} they only exist if $_{0}$ is su ciently large. The corresponding starting points for $_{k}$ are connected by the dotted curve and, since $_{k}$ is then nearly independent of T, only the curve for T = 0 has been shown. The vertical arrow points to the vacuum value of the order parameter, $_{vac} = f = 92 \text{ MeV}$, and the free mass values $_{k} = m = 600 \text{ MeV} = c^{2}$ and $_{?} = m = 138 \text{ MeV} = c^{2}$ are indicated by the horizontal arrows. The locations of the corresponding points in the diagram are shown by the two solid symbols.



Figure 2: E ect of nite size on the e ective masses.

The elective masses are shown as functions of the temperature for either $_0 = 0$, when the O (4) symmetry is restored, or for $_0 = f$, the physical vacuum value. Three di erent scenarios have been considered: either the therm odynamic limit of large volume in which the quasi-particle spectrum is continuos (solid curve), corresponding to the scenario of g.1, or a nite cubic box with side length L = 8 fm (dashed) or L = 5 fm (dotted), where the quasi-particle modes are quantized.





The typical magnitude of the spatial uctuations in the eld strength, 2 $^{1=2}$, as a function of the magnitude of the order parameter, $_{0}$, for speci ed tem peratures T: 122.63 (= T_c), 160, 200, 240 M eV. The system is enclosed in a cubic box of side length L ! 1 (solid curves), L = 8 fm (dashed), or L = 5 fm (dotted). The lower four curves display the uctuations along the chiral direction of the order parameter , while the upper curves represent the total uctuation.





The elective potential energy density V_T along the axis for either the O (4) symmetric model having H = 0 (part a) or with the adopted positive value of H (part b). The solid curves show the results for a number of tem peratures: 100, 122.63 (= T_c), 160, 200, 240 M eV. For T < T_c the elective potential curve starts at a certain minimum value of $_0$ between 0 and v, These starting points are connected by the dotted curve, while the dashed curve shows the bare potential V_0 obtained when uctuations are neglected. The arrow spoint to the minim a of V_T ; for H > 0 there is only a single minimum (located at $_0 = f$), while for H = 0 the degenerate ground-state minim a form the surface of the 4-sphere determined by = v. The value of the elective potential corresponding to other orientations of the order parameter can be easily obtained by noting that the directional dependence of V_T is given by H $_0 \cos _0 = h^3 c^3$.





The elective potential energy density V_T along the axis for either the O (4) symmetric model having H = 0 (part a) or with the adopted positive value of H (part b). The solid curves show the results for a number of tem peratures: 100, 122.63 (= T_c), 160, 200, 240 M eV. For T < T_c the elective potential curve starts at a certain minimum value of $_0$ between 0 and v, These starting points are connected by the dotted curve, while the dashed curve shows the bare potential V_0 obtained when uctuations are neglected. The arrow spoint to the minim a of V_T ; for H > 0 there is only a single minimum (located at $_0 = f$), while for H = 0 the degenerate ground-state minim a form the surface of the 4-sphere determined by = v. The value of the elective potential corresponding to other orientations of the order parameter can be easily obtained by noting that the directional dependence of V_T is given by H $_0 \cos _0 = h^3 c^3$.



Figure 5: Free energy (part a: H = 0).

The free energy density F_T (_) along the axis for either the O (4) symmetric model having H = 0 (part a) or with the adopted positive value of H (part b). The solid curves show the results for a number of temperatures. For T < T_c the curve starts at a certain m inimum value of the magnitude $_0$ and these starting points are connected by the dotted curve, while the dashed curve shows the result obtained when the temperature is neglected. The arrows point to the m inim a of F_T ; for H > 0 there is only a single m inimum, while for H = 0 the degenerate m inim a form the surface of a 4-sphere. For each temperature the location of the m inim a are indicated by the solid dots. (For H = 0 the shallow secondary m inim a at $_0 = 0$ are shown by open dots.)



Figure 5: Free energy (part b: H > 0).

The free energy density F_T (_) along the axis for either the O (4) symmetric model having H = 0 (part a) or with the adopted positive value of H (part b). The solid curves show the results for a number of temperatures. For T < T_c the curve starts at a certain m inimum value of the magnitude $_0$ and these starting points are connected by the dotted curve, while the dashed curve shows the result obtained when the temperature is neglected. The arrows point to the m inim a of F_T ; for H > 0 there is only a single m inimum, while for H = 0 the degenerate m inim a form the surface of a 4-sphere. For each temperature the location of the m inim a are indicated by the solid dots. (For H = 0 the shallow secondary m inim a at $_0 = 0$ are shown by open dots.)



Figure 6: Tem perature dependence of the order param eter.

The most probable value of $_0$, the magnitude of the order parameter, in the standard model where H > 0. In the limit of a very large box, L ! 1 (solid curve) $_0$ is constrained to the value for which the free energy density has its minimum (see g. 5b). The bars show the full width at half maximum of the thermal distribution of $_0$ in the system with L = 5 fm; those for L = 8 fm are about half that size (see g. 7). The open dots show the centroids for H = 0 for the box with L = 5 fm, while the light curve shows the corresponding result for a large box.



Figure 7: D istribution of the magnitude of the order parameter. For a cubic box of side length L = 8 fm is shown the probability density for the magnitude of the order parameter, P ($_0$), for a range of speci ed tem peratures T (indicated), obtained either by scaling the continuum result (solid curves) or by quantizing the quasi-particle modes (dashed curves).



Figure 8: The joint distribution P ($_0$; $_0$).

The projection of the probability density P (_) onto the variables $_0$ (the magnitude of the order parameter) and $_0$ (the disalignment angle) is displayed as a function of $_0 = _0 \cos_0$ and $_0 = _0 \sin_0$, for a cubic box of side length L = 8 fm. For each temperature T, the solid dot indicates the location of the maximum of P ($_0$; $_0$) and the solid curve traces out the half-maximum contour, obtained by scaling the continuum results. For the temperatures 200 and 240 m eV is indicated the corresponding result of quantizing the quasi-particle modes (dashed contours and open centroid dots).



Figure 9: Most probable e ective masses.

The most probable e ective masses $_k$ (solid) and $_?$ (dashed) are shown as functions of the temperature T, with the error bars indicating the full width at half maximum of the distribution resulting for a box with a side length of L = 8 fm. The results obtained by scaling the continuum results are indistinguishable from those of the corresponding quantized treatment.



Figure 10: Correlation function at $T = T_c$. The reduced correlation function $C_{12} = (r_1) (p) = {}^2$ calculated at the critical temperature for various magnitudes of the order parameter, ranging from $_0 = 0$ to the vacuum value, $_{vac} = f = 92 \text{ MeV}$. The most probable value at $T = T_c$ is $_0 = 87 \text{ MeV}$.



F igure 11: Tem perature dependence of correlation function. The reduced correlation function C_{12} for a range of tem peratures T, employing for each one the most probable value of $_0$, the magnitude of the order parameter.



Figure 12: Correlation length.

The correlation length $_{12}$ (the full width of C $_{12}$ at half maximum) as a function of tem perature, employing the most probable value of $_0$ for each T.





The strength function P_2 (m c^2) determ ining the overall rate at which ⁰ m esons are being produced (see eq. (55)) in a source in them al equilibrium, as a function of its tem perature T. This result is obtained by perform ing a Fourier transform of the quasi-particle correlation function associated with a given order parameter ₀ and subsequently averaging over its distribution, $P_0(_0)$. The overall norm alization is arbitraty.



Figure 14: Early dynam ics of the order param eter.

The early trajectories of the order parameter is shown for a sample of eight congurations, considering a box with L = 8 fm with a temperature of either 200 or 240 MeV. The display is similar to that in g.8 and the half-density contours (dashed) as well as the centroids (open dots) are those already given there. For each individual trajectory, the initial location is indicated by the solid dot and the attached solid curve traces out the dynam ical path up to the time t = 1 fm = c.



Figure 15: T in e-averaged distribution of the order parameter.

This gure illustrates the in uence of the time evolution on the distribution of the magnitude of the order parameter, $_0$, for a bow with L = 8 fm and for the tem – peratures of 180 and 240 M eV. The solid curves show the initial distribution of $_0$, as given by approximate statistical distribution P ($_0$) (see eq. (43)). A sample of 40 systems are then followed up to the time t = 10 fm = c and the order parameter is binned at regular time intervals throughout the evolution, leading to the dashed curves.



Figure 16: T in e evolution of the correlation function.

The pion correlation function C_{12} (s_{12}) at the tem perature T = 240 M eV. The dotted curve is the continuum limit (L ! 1) and the solid curve is the corresponding therm all result for a quantized nit box with a side length of L = 5 fm. The correlation function for a sample of ten initial con gurations are shown by the short-dashed curves, and the long-dashed curves show the corresponding result after they have been propagated self-consistently up to the time t = 10 fm = c. The dashed curves have been obtained in two di erent ways: The curves that go up again result from aligning the separation r_{12} along one of the cartesian directions, while the other two are obtained for separations directed along a diagonal. The aligned qurves have a periodicity equal to L, whereas the periodicity of the diagonal curves is 3 tim es larger.