

TM UP-HEL-9604

February 1996

THREE FLAVOR NEUTRINO OSCILLATION
CONSTRAINTS FROM ACCELERATOR,
REACTOR, ATMOSPHERIC AND SOLAR
NEUTRINO EXPERIMENTS

Osamu Yasuda¹ and Hisakazu Minakata²

Department of Physics, Tokyo Metropolitan University

1-1 Minamiosaka Hachioji, Tokyo 192-03, Japan

Abstract

We discuss constraints on three flavor neutrino mixings from the accelerator and reactor experiments, the Kamiohka multi-GeV data, and the solar neutrino observations. The LSND result is excluded at 90% CL by the constraints imposed by all the data of reactor and accelerator experiments and the Kamiohka multi-GeV data if the mass scale required for the solution to the solar neutrino problem is hierarchically small. The region of a set of the effective two-flavor mixing parameters (Δm^2 , $\sin^2 2\theta$) is given for the

¹Email: yasuda@phys.metro-u.ac.jp

²Email: minakata@phys.metro-u.ac.jp

channel $\pi^+ \pi^-$ which is allowed at 90% CL by the multi-GeV KAMIOKANDe data alone.

Recently the LSND collaboration has claimed that they have found candidate events for $\bar{\nu}_\mu \rightarrow \nu_e$ oscillation [1] (See also [2]). If their result turns out to be correct, it gives us an important information form masses and mixing angles of neutrinos. In this paper a possibility is explored that all the experimental data, solar, atmospheric, reactor and accelerator data including LSND can be explained within a framework of three flavor neutrino mixing. It turns out that the LSND result is excluded at 90% CL by the constraint imposed by all the data of reactor and accelerator experiments and the Kamikande multi-GeV data. The statement is true even without invoking actual solar neutrino data as far as the mass scale required for the solution is hierarchically small.

Among the atmospheric neutrino data [3] [4] [5] [6] [7] [8] we discuss only the Kamikande multi-GeV data [4] throughout this paper. The reason for this restriction is three-fold; (1) It is the only experiment that gives us a nontrivial zenith-angle dependence which leads to the upper and lower bounds for the mass-squared difference of neutrinos. (2) The result of Monte-Carlo simulation for neutrino energy spectrum is published only for the the Kamikande multi-GeV data. (3) It seems difficult to reconcile it with NUSEX [6] and Frejus [7] data.

The Dirac equation for three flavors of neutrinos in vacuum is given by

$$i \frac{d}{dx} (\mathbf{x}) = U \text{diag}(E_1; E_2; E_3) U^{-1} (\mathbf{x}); \quad (1)$$

which is easily solved:

$$(\mathbf{x}) = \sum_{j=1}^3 U_j U_j^{-1} e^{iE_j x} (\mathbf{x}); \quad (2)$$

where $E_j = \sqrt{p^2 + m_j^2}$ ($j = 1; 2; 3$) is the energy of neutrinos in the mass basis, $(\mathbf{x}) = (\nu_e(x); \nu_\mu(x); \nu_\tau(x))^T$ ($= e; \mu; \tau$) is the wave function of

neutrinos in the flavor basis, and

$$U = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13} \\ s_{12}c_{23} & c_{12}c_{23} & s_{23}c_{13} \\ s_{12}s_{23} & c_{12}s_{23} & c_{23}c_{13} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad (3)$$

with $c_{ij} = \cos \theta_{ij}$; $s_{ij} = \sin \theta_{ij}$ is the orthogonal mixing matrix of neutrinos. We will not discuss the CP violating phase of the mixing matrix here for simplicity. The probability of $\nu_\alpha \rightarrow \nu_\beta$ transition is given by

$$P(\nu_\alpha \rightarrow \nu_\beta; E, L) = \begin{cases} \frac{1}{4} \sin^2 \frac{E_{ij}L}{2} U_{\alpha i}^2 U_{\beta j}^2 & \text{for } \alpha = \beta \\ \frac{1}{4} \sin^2 \frac{E_{ij}L}{2} U_{\alpha i} U_{\beta j} U_{\alpha j} U_{\beta i} & \text{for } \alpha \neq \beta \end{cases} \quad (4)$$

where $E_{ij} = E_i - E_j$, $(m_i^2 - m_j^2) = 2E_{ij}$ is the difference of the energy of two mass eigenstates.

The number of neutrinos (ν_e, ν_μ, ν_τ) is measured in terms of charged leptons e, μ, τ which come out from a scattering $X \rightarrow \nu X^0$. In appearance experiments of ν_α the expected number of the charged leptons ℓ is given by

$$N(\nu_\alpha \rightarrow \ell; L) = n_T \int_0^L dE \int_0^{q_{max}} dq (q) F(E) \frac{d}{dq} (E; q) P(\nu_\alpha \rightarrow \ell; E, L); \quad (5)$$

whereas in disappearance experiments we measure attenuation of beam

$$N(\ell; L) = N(\nu_\alpha \rightarrow \ell; L) = n_T \int_0^L dE \int_0^{q_{max}} dq (q) F(E) \frac{d}{dq} (E; q) (1 - P(\nu_\alpha \rightarrow \nu_\alpha; E, L)); \quad (6)$$

where

$$N(L) = n_T \int_0^{Z_1} dE \int_0^{q_{max}} dq (q) F(E) \frac{d\sigma(E; q)}{dq} : \quad (7)$$

$F(E)$ is the flux of neutrino with energy E , n_T is the number of target nucleons, (q) is the detection efficiency function for charged leptons of energy q , $d\sigma(E; q)/dq$ is the differential cross section of the interaction $\nu + X \rightarrow \ell + X'$, $P(\nu_\mu \rightarrow \nu_\tau; E; L)$ is the probability of $\nu_\mu \rightarrow \nu_\tau$ transitions with energy E after traveling a distance L . The results of experiments are usually expressed in terms of a set of the oscillation parameters $(\Delta m^2; \sin^2 2\theta)$ in the two-flavor analysis. The probability in the two-flavor mixing is given by

$$P(\nu_\mu \rightarrow \nu_\tau; E; L) = \begin{cases} \sin^2 2\theta \sin^2 \frac{\Delta m^2 L}{4E} & \text{for } \Delta m^2 \ll 4E \\ \sin^2 2\theta & \text{for } \Delta m^2 \gg 4E \end{cases} \quad (8)$$

Introducing the notation

$$\frac{1}{N(L)} n_T \int_0^{Z_1} dE \int_0^{q_{max}} dq (q) F(E) \frac{d\sigma(E; q)}{dq} \sin^2 \frac{\Delta m^2 L}{4E} ; \quad (9)$$

we have the expected number of charged leptons in the two-flavor scenario

$$\begin{aligned} \frac{N^{(2)}(\nu_\mu \rightarrow \nu_\tau; L)}{N(L)} &= \sin^2 2\theta \left(\frac{\Delta m^2}{4E} \right) \sin^2 \frac{\Delta m^2 L}{4E} \quad \text{(disappearance experiments)} \\ \frac{N^{(2)}(\nu_\mu \rightarrow \nu_\tau; L)}{N(L)} &= \sin^2 2\theta \left(\frac{\Delta m^2}{4E} \right) \sin^2 \frac{\Delta m^2 L}{4E} \quad \text{(appearance experiments)}; \end{aligned} \quad (10)$$

where we have denoted the m^2 dependence of the mixing angle $\theta(m^2)$ explicitly to indicate that $\theta(m^2)$ is a function of m^2 in the two-flavor analysis.

The boundary of the excluded region (for negative results) in the $(m^2; \sin^2 2\theta)$ plot is determined by

$$= \frac{N^{(2)}(\theta; L)}{N(L)} = \sin^2 2\theta(m^2) \sin^2 \frac{m^2 L}{4E}; \quad (11)$$

for appearance experiments where the charged leptons are detected at one point at a distance L , or

$$= \frac{N^{(2)}(\theta; L_1)}{N(L_1)} = \sin^2 2\theta(m^2) \sin^2 \frac{m^2 L_2}{4E} \sin^2 \frac{m^2 L_1}{4E} \quad (12)$$

for disappearance experiments where the charged leptons are detected at two points at distances L_1 and L_2 ($L_1 < L_2$). In (11) and (12) θ denotes the largest fraction of the appearance events allowed by a given confidence level, i.e., $N^{(2)}(\theta; L) = N(L) \times \theta$ for appearance experiments, and the largest fraction of beam attenuation, $N^{(2)}(\theta; L_1) = N(L_1) \times \theta$, $N^{(2)}(\theta; L_2) = N(L_2) \times \theta$ for disappearance experiments.

From (11) and (12), we can read off the value of $\sin^2 \frac{m^2 L}{4E}$ for arbitrary m^2 from the figure of the two-flavor mixing parameters $(m^2; \sin^2 2\theta)$ given in each experimental paper as long as $\sin^2 2\theta(m^2) \neq 0$:

$$\sin^2 \frac{m^2 L_2}{4E} = \frac{\sin^2 \frac{m^2 L_1}{4E}}{\sin^2 2\theta(m^2)} \quad (\text{disappearance experiments})$$

³In case of the CP violating phase, which will not be discussed in the present paper, this is not the case. To discuss the CP violating effect, one needs the information of $\sin \frac{m^2 L}{2E}$, which cannot be obtained from the information in published papers only.

$$\sin^2 \frac{m^2 L}{4E} = \frac{\sin^2 2\theta (m^2)}{\sin^2 2\theta (m^2)} \quad (\text{appearance experiments}): \quad (13)$$

In (13) and in the following, $\sin^2 2\theta (m^2)$ or $\sin^2 2\theta (m^2)$ stands for the value of $\sin^2 2\theta (m^2)$ on the boundary of the allowed region in the $(m^2; \sin^2 2\theta)$ plot in the two-flavor analysis.

Similarly we can express the number of the expected charged leptons in the three-flavor mixing:

$$\begin{aligned} 1 \frac{N^{(3)}(\nu_\mu; L)}{N(L)} &= 4 \sum_{i < j} U_{\mu i}^2 U_{\mu j}^2 \sin^2 \frac{m_{ij}^2 L}{4E} \quad \text{for } \nu_\mu \\ \frac{N^{(3)}(\nu_\mu; L)}{N(L)} &= 4 \sum_{i < j} U_{\mu i} U_{\mu j} U_{\mu i} U_{\mu j} \sin^2 \frac{m_{ij}^2 L}{4E} \quad \text{for } \nu_\mu : (14) \end{aligned}$$

From (13) we observe that the quantity $\sin^2 (m^2 L = 4E) = \frac{D}{E}$ is equal to $\sin^2 2\theta (m^2)$ which can be read off from the published literatures, and we can express the conditions for the three-flavor mixing parameters in case of negative results:

$$\begin{aligned} &> \frac{N^{(3)}(\nu_\mu; L_1)}{N(L_1)} - \frac{N^{(3)}(\nu_\mu; L_2)}{N(L_2)} = \sum_{i < j} \frac{U_{\mu i}^2 U_{\mu j}^2}{\sin^2 2\theta (m_{ij}^2)} \\ &\quad (\text{disappearance experiments}) \\ &> \frac{N^{(3)}(\nu_\mu; L)}{N(L)} = 4 \sum_{i < j} \frac{U_{\mu i} U_{\mu j} U_{\mu i} U_{\mu j}}{\sin^2 2\theta (m_{ij}^2)} \\ &\quad (\text{appearance experiments}): (15) \end{aligned}$$

Notice that the left-hand side of (13) is defined independent of the number of flavors of neutrinos.

Throughout this paper we assume that a single mass scale is involved in the solution of the solar neutrino problem, which is hierarchically small compared to others. Namely, we assume that

$$m_{21}^2, m_{32}^2 < m_{31}^2; \quad (16)$$

where we have assumed $m_1^2 < m_2^2 < m_3^2$ without loss of generality. In the case where $m_{32} = m_{21}^2 < m_{31}^2$, we can show that we obtain the same conclusions, although we will not give the calculation here.

The hierarchy (16) is satisfied in the two-flavor mixing solution to the solar neutrino problem [9] [10] which requires,

$$\begin{aligned} & (m^2; \sin^2 2\theta) \\ & \begin{cases} (m_{21}^2; \sin^2 2\theta_{12}) \\ \geq (O(10^{-11} \text{eV}^2); O(1)); & \text{(vacuum solution)} \\ \geq (O(10^{-5} \text{eV}^2); O(10^{-2})); & \text{(small angle MSW solution)} \\ \geq (O(10^{-5} \text{eV}^2); O(1)) & \text{(large angle MSW solution):} \end{cases} \end{aligned} \quad (17)$$

These mass scales are much smaller than those which appear in the atmospheric neutrino observations [4], or in the LSND experiment [1]. In fact the only condition on $(m_{21}^2; \sin^2 2\theta_{12})$ which is essential to the discussions below is (16), and our conclusions remain unchanged irrespective of the value of θ_{12} as long as (16) holds.

One can show [12] [13] that under the mass hierarchy (16) the relation

$$U_{e3}^2 = 1; \quad (18)$$

must hold in order to have solar neutrino deficit under the constraints from the accelerator and the reactor experiments. In this setting it can also be demonstrated that the solar neutrino problem is indeed solved by a two-flavor framework in the MSW and the vacuum solutions [12].

In the present case the formulas in (14) become much simpler [11]:

$$\begin{aligned} 1 - P(\nu_e \rightarrow \nu_e) &= 4 \sin^2 \frac{m_{21}^2 L}{4E} U_1^2 U_2^2 + 4 \sin^2 \frac{m_{31}^2 L}{4E} U_3^2 (1 - U_3^2) \\ P(\nu_e \rightarrow \nu_e) &= 4 \sin^2 \frac{m_{21}^2 L}{4E} U_1 U_2 U_1 U_2 + 4 \sin^2 \frac{m_{31}^2 L}{4E} U_3^2 U_3^2 : \end{aligned} \quad (19)$$

For the cases discussed below, we have $j m_{21}^2 L = 4E j \ll 1$ and the first term in each formula in (19) can be ignored. Hence we get

$$\begin{aligned} 4U_3^2 (1 - U_3^2) &= \sin^2 2 \theta (m_{31}^2) \quad (\text{disappearance experiments}) \\ 4U_3^2 U_3^2 &= \sin^2 2 \theta (m_{31}^2) \quad (\text{appearance experiments}) \end{aligned} \quad (20)$$

for negative results, respectively.

The data by the LSND group suggests that the allowed region for the mass-squared difference m_{31}^2 is approximately $m_{31}^2 > 5 \times 10^{-2} \text{eV}^2$. If we combine the data on the same channel $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ (and $\nu_\mu \rightarrow \nu_e$) by the E776 group [14], however, the region $m_{31}^2 > 2.5 \text{eV}^2$ seems to be almost excluded, and we have $5 \times 10^{-2} \text{eV}^2 < m_{31}^2 < 2.5 \text{eV}^2$ at 90% confidence level.

As has been pointed out in Ref. [11], strong constraints on the mixing angle come from the reactor experiment [15]. Using (20) we have the constraint from the reactor experiment [15]

$$\sin^2 2 \theta_{13} = 4U_{e3}^2 (1 - U_{e3}^2) = \sin^2 2 \theta_{\text{Bugey}} (m_{31}^2); \quad (21)$$

where $\sin^2 2 \theta_{\text{Bugey}} (m_{31}^2)$ stands for the value of $\sin^2 2 \theta$ on the boundary of the allowed region in the $(m_{31}^2; \sin^2 2 \theta)$ plot in [15]. For the entire region $5 \times 10^{-2} \text{eV}^2 < m_{31}^2 < 2.5 \text{eV}^2$, $\sin^2 2 \theta_{\text{Bugey}} (m_{31}^2)$ is small [15]:

$$\frac{1}{50} < \sin^2 2 \theta_{\text{Bugey}} (m_{31}^2) < \frac{1}{10}; \quad (22)$$

From (21) we have either small U_{e3}^2 or large U_{e3}^2 , but the latter possibility is excluded by (18). Therefore we have

$$s_{13}^2 = U_{e3}^2 < \frac{1}{4} \sin^2 2 \theta_{\text{Bugey}} (m_{31}^2); \quad (23)$$

On the other hand, we have another constraint from the disappearance experiment of ν_μ [16]

$$4s_{23}^2 c_{13}^2 (1 - s_{23}^2 c_{13}^2) = 4U_3^2 (1 - U_3^2) = \sin^2 2 \theta_{\text{CDHSW}} (m_{31}^2); \quad (24)$$

where $\sin^2 2_{\text{CDHSW}}(m_{31}^2)$ stands for the value of $\sin^2 2$ on the boundary of the allowed region in the $(m^2; \sin^2 2)$ plot in [16]. The mixing angle in this case is constrained for $0.7\text{eV}^2 < m_{31}^2 < 13\text{eV}^2$ as

$$\sin^2 2_{\text{CDHSW}}(m_{31}^2) < 0.2; \quad (25)$$

If we consider the probability $P(\nu_\mu \rightarrow \nu_\mu)$ in the atmospheric neutrino experiments for the mass region above, we have small deviation of $P(\nu_\mu \rightarrow \nu_\mu)$ from unity

$$1 - P(\nu_\mu \rightarrow \nu_\mu) \approx 2U_3^2(1 - U_3^2) < 0.1; \quad (26)$$

where we have averaged over rapid oscillations. (26) is obviously inconsistent with the atmospheric neutrino observations [3] [4] [5] [8], since we cannot have gross deficit of ν_μ in this case. In fact we have verified explicitly, by the same calculation as in Ref. [17], that the region $m_{31}^2 > 0.47\text{eV}^2$ with the constraints (21) and (24) is excluded at 95% confidence level by the Kamikande multi-GeV data ($\chi^2 = 17.6$ for 3 degrees of freedom, where $\chi^2_{\text{min}} = 3.2$, $(\chi^2 - \chi^2_{\text{min}}) = 7 \times 2.1$ implies 2 σ).

So we are left with the region $5 \times 10^{-2}\text{eV}^2 < m_{31}^2 < 0.47\text{eV}^2$. Applying the formula (19) to the case of LSND, we have

$$s_{23}^2 \sin^2 2_{13} = 4U_{e3}^2 U_3^2 = \sin^2 2_{\text{LSND}}(m_{31}^2); \quad (27)$$

where $\sin^2 2_{\text{LSND}}(m_{31}^2)$ stands for the value of $\sin^2 2$ within the allowed region in the $(m^2; \sin^2 2)$ plot in [1], and the LSND data indicates

$$1.5 \times 10^{-3} < \sin^2 2_{\text{LSND}}(m_{31}^2) < 1; \quad (28)$$

From (21) and the constraint $s_{23}^2 < 1$, it follows that

$$\sin^2 2_{\text{LSND}}(m_{31}^2) = \sin^2 2_{\text{Bugey}}(m_{31}^2); \quad (29)$$

so that we have

$$m_{31}^2 > 0.25 \text{eV}^2; \quad (30)$$

with

$$1.5 \cdot 10^{-3} < \sin^2 2_{\text{LSND}} \left(\frac{m_{31}^2}{\text{eV}^2} \right) < 3.8 \cdot 10^{-2}; \quad (31)$$

where we have used the $(m_{31}^2; \sin^2 2)$ plots in [15] and [1]. However, we have verified explicitly again that the region $m_{31}^2 > 0.25 \text{eV}^2$ is excluded at 90% confidence level by the Kamikande multi-GeV data ($\chi^2 = 15$ for three degrees of freedom, $(\chi^2_{\text{min}} = 7, 1:7 \text{ implies } 1.6 \sigma)$). Therefore, we conclude that the LSND data cannot be explained by neutrino oscillations among three flavors, if all the accelerator and reactor data as well as the Kamikande multi-GeV data are taken for granted.

The present LSND data allows conflicting interpretations either as a possible evidence for neutrino oscillation [1], or a stringent bound for the mixing parameters [2]. It might be possible that the allowed region of the set of the parameters $(m_{31}^2, \sin^2 2)$ implied by the LSND data changes in the future. We have looked for the region of $(m_{31}^2, \sin^2 2)$ for $\nu_e \rightarrow \nu_e$ (or $\bar{\nu}_e \rightarrow \bar{\nu}_e$), in which $\nu_e \rightarrow \nu_e$ oscillation is consistent with all the experiments (except LSND), including the Kamikande multi-GeV data. To obtain the $(m_{31}^2, \sin^2 2_{\nu_e})$ plot of the two-flavor analysis for general $\nu_e \rightarrow \nu_e$ oscillations with a mixing angle $\theta_e(m_{31}^2)$, we use the following correspondence between the rates for the $\nu_e \rightarrow \nu_e$ process in the three and the two-flavor frameworks:

$$\begin{aligned} \frac{N(\nu_e \rightarrow \nu_e; L)}{N_{\nu_e}(L)} &= 4U_{e3}^2 U_3^2 \sin^2 \frac{m_{31}^2 L}{4E} \\ &= s_{23}^2 \sin^2 2_{13} \sin^2 \frac{m_{31}^2 L}{4E} \\ &= \sin^2 2_{\nu_e}(m_{31}^2) \sin^2 \frac{m_{31}^2 L}{4E}; \end{aligned} \quad (32)$$

where we have used the hierarchical condition (16). From (21) and (32) we have

$$\sin^2 2_{\theta_e} (m_{31}^2) \leq s_{23}^2 \sin^2 2_{13} \leq s_{23}^2 \sin^2 2_{\text{Bugey}} (m_{31}^2): \quad (33)$$

The Kamikande multi-GeV data has better fit for larger values of $\sin^2 2_{23}$, so (33) shows that $\sin^2 2_{\theta_e} (m_{31}^2)$ cannot be larger than $\sin^2 2_{\text{Bugey}} (m_{31}^2) = 2$, if $\sin^2 2_{13} \leq 1^4$. The result is shown in Fig.1.

(Insert Fig.1 here.)

The region suggested by the LSND experiment [1] is close to the 90% CL region obtained in our analysis. If the 20% systematic uncertainty mentioned in Ref. [1] shifts the allowed region of LSND in the direction of smaller mixing, there may be a chance that all the neutrino anomalies are explained by three flavor neutrino oscillations. Hopefully further data from the LSND group will clarify the situation.

Acknowledgement

One of the authors (O.Y.) would like to thank K.S. Babu, P.I. Krastev, C.N. Leung, and A. Smirnov for discussions. We would like to thank K. Tida for collaboration in the early stages of this work and members of the Physics Department of Yale University for their hospitality during our stay while a part of this work was done. This research was supported in part by a Grant-in-Aid for Scientific Research, Ministry of Education, Science and Culture, # 05302016, # 05640355, and # 07044092.

⁴This is not the case for $m_{31}^2 < 3 \times 10^{-2} \text{eV}^2$, where $1 - P(\nu_e \rightarrow \nu_e) = 4s_{23}^2 c_{13}^2 (1 - s_{23}^2 c_{13}^2) \approx 0$ (1) becomes possible even for $2_{23} < \pi/4$.

References

- [1] LSND Collaboration, C. Athanassopoulos, et al., Phys. Rev. Lett. 75 (1995) 2650.
- [2] J.E. Hill, Phys. Rev. Lett. 75 (1995) 2654.
- [3] Kamioke Collaboration, K. S. Hirata et al., Phys. Lett. B 205 (1988) 416; *ibid.* B 280 (1992) 146.
- [4] Kamioke Collaboration, Y. Fukuda et al., Phys. Lett. B 335 (1994) 237.
- [5] IMB Collaboration, D. Casper et al., Phys. Rev. Lett. 66 (1989) 2561; R. Becker-Szendy et al., Phys. Rev. D 46 (1989) 3720.
- [6] NUSEX Collaboration, M. Aglietta et al., Europhys. Lett. 8 (1989) 611.
- [7] Frejus Collaboration, Ch. Berger et al., Phys. Lett. B 227 (1989) 489; *ibid.* B 245 (1990) 305; K. Daum et al., Z. Phys. C 66 (1995) 417.
- [8] Soudan 2 Collaboration, M. Goodman et al., Nucl. Phys. B (Proc. Suppl.) 38 (1995) 337.
- [9] See, e.g., J.N. Bahcall and R.K. Ulrich, Rev. Mod. Phys. 60 (1988) 297; J.N. Bahcall and M.H. Pinsonneault, Rev. Mod. Phys. 64 (1992) 885; J.N. Bahcall, R. Davis, Jr., P. Parker, A. Smirnov, R. Ulrich eds., SOLAR NEUTRINOS: the first thirty years Reading, Mass., Addison-Wesley, 1994 and references therein.
- [10] S.P. Mikheyev and A. Smirnov, Nuovo Cim. 9C (1986) 17; L. Wolfenstein, Phys. Rev. D 17 (1978) 2369.

- [11] G.L. Fogli, E. Lisi, D. Montanino, Phys. Rev. D 49 (1994) 3626; H. Minakata, Phys. Lett. B 356 (1995) 61; Phys. Rev. D 52 (1995) 6630; S.M. Bilenky, A. Bottino, C. Giunti, C.W. Kim, Phys. Lett. B 356 (1995) 273; preprint DFTT 2/96 (hep-ph/9602216); S.M. Bilenky, C. Giunti, C.W. Kim, preprint DFTT-30-95 (hep-ph/9505301); K.S. Babu, J.C. Pati and F.W. Ilczek, Phys. Lett. B 359 (1995) 351, B 364 (1995) 251 (E); M. Narayan, M.V.N. Murthy, G. Rajasekaran, S. Uma Sankar, preprint IMSC-95-96-001 (hep-ph/9505281); G.L. Fogli, E. Lisi and G. Scioscia, Phys. Rev. D 52 (1995) 5334; S. Goswami, K. Kar and A. Raychaudhuri, preprint CUPP-95-3 (hep-ph/9505395).
- [12] The first paper of H. Minakata in [11].
- [13] The first paper of S.M. Bilenky et al. in [11].
- [14] L. Borodovsky et al., Phys. Rev. Lett. 68 (1992) 274.
- [15] B. Acker et al., Nucl. Phys. B 434, (1995) 503.
- [16] CDHSW Collaboration, F. Dyak et al., Phys. Lett. 134B (1984) 281.
- [17] O. Yasuda, preprint TM UP-HEL-9603 (hep-ph/9602342).

Figures

Fig.1 Below the solid and dashed lines are the regions for $\nu_\mu \rightarrow \nu_e$ (or $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$) oscillation which are compatible with all the experiments (except LSND), including the atmospheric multi-GeV data of Kamikande at 68% CL (solid) and 90% CL (dashed), respectively. To get this plot of the two-flavor mixing parameters, we have used the correspondence $\sin^2 2\theta_{\text{atm}} = \sin^2 2\theta_e (\tan^2 \theta_{31}) = s_{23}^2 \sin^2 2\theta_{13}$ (cf. (32)). The shadowed area stands for the region allowed by all the accelerator and reactor experiments including LSND (The dotted, dashed and dot-dashed lines stand for the LSND [1], E776 [14], and Bugey [15] experiments, respectively).




Fig.1