

High E_T jets at pp collisions and triple gauge boson vertex

B. A. Arbuzov

Institute for High Energy Physics, Protvino,
Moscow region, 142284, Russia

Abstract

Triple gauge boson vertex, which is inherent to the model of a dynamical breaking of the electroweak symmetry, leads to an effective four-fermion interaction of quarks. Calculation of the effective coupling constant in the framework of the model results in agreement with recent information on enhancement of inclusive high E_T jet cross section at $\sqrt{s} = 1.8 \text{ TeV}$. Thus this effect may be connected with triple gauge boson vertex with coupling constant 0.7, which will clearly manifest itself at the soon forthcoming LEP II.

PASC : 12.15; 14.80 Er

Keywords: gauge boson vertex; four-fermion interaction; jets

In the recent Fermilab experiment [1] the effect of an enhanced cross section of inclusive jets production at $\sqrt{s} = 1.8 \text{ TeV}$ with $E_T > 200 \text{ GeV}$ is obtained. The authors state, that this effect could be connected with consequences of quark compositeness, leading e.g. to effective four-fermion interaction [2]

$$L_{\text{eff}} = \frac{4}{2c^2} q_L^a q_L^a q_L^b q_L^b; \quad (1)$$

where q corresponds to light quark pair $u; d$ and currents in interaction (1) are isoscalar. The best agreement of data [1] with (1) is achieved for

$$c = 1.6 \text{ TeV}; \quad (2)$$

In this letter we would point out, that effective four-fermion interaction appears not only in the case of quarks substructure but also in the case of an existence of triple gauge vertex ($\epsilon^{abc} W^a W^b W^c$), which appears in a selfconsistent way in the model of a dynamical breaking of the electroweak symmetry [3-6]

$$\begin{aligned} \epsilon^{abc}(p; q; k) = & \frac{g}{M_W^2} \epsilon^{abc} F(p; q; k) - g(p(qk) - q(pk)) + \\ & + g(q(pk) - k(pq)) + g(k(pq) - p(qk)) + k p q - q k p; \quad (3) \end{aligned}$$

where form factor F restricts the small momentum region of action of the vertex by an effective cut-off E_W , which is of few TeV order of magnitude. According to estimates [5], [6], which are consistent with t -quark production cross section, $Z \rightarrow b\bar{b}$ width and experimental limitations [7], [8]

$$E_W = 4.5 \text{ TeV}; \quad 0.7 : \quad (4)$$

Note, that additional gauge boson vertices are repeatedly introduced basing on various arguments (see e.g. [9]).

Let us calculate effective four-fermion terms in one loop approximation using vertex (3) and usual EW interaction

$$\frac{g}{2} \bar{q}_L^a q_L^a + \sum_j Q_L^j \bar{q}_L^a Q_L^j W^a; \quad (5)$$

where $j = 1, 2$ marks generations of heavier quarks. We use here the electroweak interaction without symmetry breaking, because we deal with TeV energy region, corresponding to restoration of the symmetry.

We have two types of diagrams: s -type and t -type. They both contain triangle diagram with one vertex (3) and two vertices (5). Calculation of this diagram gives the following result (see also [10])

$$\frac{g^3}{64\pi^2 M_W^2} \ln \frac{E_W^2}{M_W^2} \bar{q}_L^a (k^2 - k \hat{k}) q_L W^a; \quad (6)$$

where k is the momentum of W . The second term in brackets is proportional to M_q , so it is negligible. The first one is proportional to k^2 , so being multiplied by W propagator $(k^2 - M_W^2)^{-1}$ it leads to contact four-fermion interaction for sufficiently large $k^2 - M_W^2$. We have two s -type diagrams (with triangle to the left and to the right) and two t -type ones. Now we obtain the effective four-fermion interaction, of which we write down terms, containing at least one qq factor and thus giving contribution to pp processes

$$\begin{aligned} L_{eff} = & \frac{G}{2} \bar{q}_L^a q_L^a \bar{q}_L^0 q_L^0 + 2 \sum_j Q_L^j \bar{q}_L^a Q_L^j + \\ & + \bar{q}_L^b q_L^0 \bar{q}_L^0 q_L^b; \end{aligned} \quad (7)$$

$$G = \frac{\frac{2}{M_W^2}}{\frac{2}{M_W^2}} \ln \frac{E_W^2}{M_W^2}; \quad g_w = \frac{g^2}{4} = \frac{1}{\sin^2 \theta_w};$$

Here we introduce primes to distinguish initial (not primed) and final (primed) light quarks in our diagrams. Really the last term in Eq.(7) corresponds to t -type diagrams and it is convenient to perform Fierz transformation

$$\bar{q}_L^b q_L^0 \bar{q}_L^0 q_L^b = \frac{3}{2} \bar{q}_L^a q_L^0 \bar{q}_L^0 q_L^a - \frac{1}{2} \bar{q}_L^a q_L^0 \bar{q}_L^a q_L^0; \quad (8)$$

From Eqs. (7) and (8) we have final form of the effective interaction

$$L_{\text{eff}} = \frac{G}{2} \left[\frac{3}{2} q_L^a q_L^a q_L^a + \frac{1}{2} q_L^a q_L^a q_L^a + 2 q_L^a q_L^a Q_L^j \right] : \quad (9)$$

The first term in lagrangian (9) coincides with Eq.(1) provided

$$c = \frac{8}{3G} = \frac{2M_W}{3 \ln(E_W/M_W)} : \quad (10)$$

Substituting values (4) into Eq.(10) we have $c = 2.87 \text{ TeV}$, that is not so far from experimental value (2).

However other terms in Eq.(9) also give contribution to the cross section. For example, we may start reasoning in the following way. Interference terms with leading QCD diagrams, which are describing the main effect of experiment [1] in the region $200 \text{ GeV} < E_T < 300 \text{ GeV}$, are caused by the first two terms of Eq.(9) and the contribution of the second term is equal to that of the first one. This result is due to simple relation

$$\text{Trace } a^a a^a = 3 \text{Trace } I : \quad (11)$$

Therefore the result for the effective c has to be divided by $\frac{P}{2}$ (amplitudes of production of heavy quarks do not interfere with QCD terms)

$$c = \frac{2.87 \text{ TeV}}{\frac{P}{2}} = 2.03 \text{ TeV} : \quad (11)$$

Result (11) is really close to value (2) and agrees with data [1] within their uncertainties. Let us mention, that the boundary value of restrictions [8] $= 0.85$ leads instead of (11) to $c = 1.84 \text{ TeV}$, that almost coincides with value (2).

To conclude we would state that exciting results [1] could be interpreted not only in terms of quark compositeness, but also in terms of anomalous gauge vertices, which also lead to an effective four-fermion interaction. Parameter in this case has to be sufficiently large, just on the boundary of experimental limitations [7], [8]. Such "strong" triple gauge vertex without doubt can be detected at the soon forthcoming LEP II facilities in the reaction of W pair production, so the way to verify the variant being discussed here is quite straightforward.

The present work is partially supported by the Russian Foundation of Fundamental Researches under project 95-02-03704.

References

- [1] F Abe et al. (CDF Coll.), <http://www-cdf.fnal.gov/physics/pub96/pub96.html> Phys Rev Lett., to be published.

- [2] E J Eichten, K D Lane and M E Peskin, *PhysRevLett.* 50 (1983) 811.
- [3] B A Arbuzov, *PhysLett. B* 288 (1992) 179.
- [4] B A Arbuzov, in: *Advanced Study Conference on: HEAVY FLAVOURS*, Pavia (Italy), ed. G Bellini et al, *Frontiers, G if-sur-Y vette*, 1994, p. 227.
- [5] B A Arbuzov and S A Shichanin, *P is'm a v ZhETF* 60 (1994) 75; *JETP Lett.* 60 (1994) 79.
- [6] B A Arbuzov, *PhysLett. B* 353 (1995) 532.
- [7] F Abe et al. (CDF Coll), *PhysRevLett.* 74 (1995) 1936.
- [8] F Abe et al. (CDF Coll), *PhysRevLett.* 75 (1995) 1017.
- [9] K Hagiwara, R D Peccei, D Zeppenfeld and K Hikasa, *NuclPhys. B* 282 (1987) 253.
- [10] K Hagiwara, S Ishihara, R Szalapski and D Zeppenfeld, *PhysRev. D* 48 (1993) 2182.