

Natural Unification with a Supersymmetric $SO(10)_{\text{GUT}} \times SO(6)_H$ Gauge Theory

T. Hotta, Izawa K.-I. and T. Yanagida

Department of Physics, University of Tokyo, Bunkyo-ku, Tokyo 113, Japan

(April 16, 2024)

Abstract

We propose a unified model of elementary particles based on a supersymmetric $SO(10)_{\text{GUT}} \times SO(6)_H$ gauge theory. This model completely achieves natural unification of the strong and electroweak interactions without any fine-tunings.

11.30.Pb, 12.10.Dm, 12.60.Jv

It has been long noticed that the quarks and leptons fit into complete multiplets of a simple gauge group such as $SU(5)_{GUT}$ [1] and $SO(10)_{GUT}$ [2] which unifies the standard gauge group. This grand unification (GUT) approach seems to yield an attractive candidate of a unified model beyond the standard one. In fact, recent high-precision measurements of the three gauge coupling constants at low energies strongly support reality of the supersymmetric (SUSY) unification of this kind [3]. However, as for the Higgs doublet, which also takes a crucial part in the standard model, the unification does not work naturally. The doublet tends to be accompanied by its triplet partner due to simplicity of the gauge group, which poses a fine-tuning problem of the doublet-triplet splitting [4].

In this letter, we propose a supersymmetric unified model based on a semisimple gauge group $SO(10)_{GUT} \times SO(6)_H$, and show that it completely achieves natural unification of elementary particles without any fine-tunings [5], keeping all the excellent successes of minimal $SO(10)$ SUSY-GUT's. In particular, the doublet-triplet splitting in the Higgs sector is realized naturally in this model. At first glance, non-simplicity of the gauge group appears to spoil unification of the standard three gauge coupling constants, which would be achieved by means of a simple gauge group. However, that is not the case. It suffices that the $SO(6)_H$ gauge interaction is strong enough at the GUT scale [8,9], which needs no fine-tuning (in a gauge theory with asymptotic freedom).

Let us first consider a SUSY $SO(10)_{GUT} \times SO(N_C)_H$ gauge theory with eleven flavors of hyperquarks Q^A ($A = 1; \dots; 11; \dots; N_C$) which transform as the vector N_C representation under the hypercolor $SO(N_C)_H$. The first ten hyperquarks Q^I ($I = 1; \dots; 10$) form the 10 representation and the last one Q^{11} is a singlet of the $SO(10)_{GUT}$. The reason why we take the $SO(N_C)_H$ (rather than the $SU(N_C)_H$) is that the anomaly-free condition is automatically satisfied for the orthogonal groups (except $SO(6) \times SU(4)$) [10]. We also introduce two kinds of $SO(N_C)_H$ -singlet chiral superfields: a 10 Higgs H_I and a 54 Higgs S_{IJ} ($I, J = 1; \dots; 10$) of the $SQ_c(10)$.

All the quark and lepton superfields constitute spinor 16 representations of the $SO(10)_{GUT}$ and singlets of the $SO(N_C)_H$. We choose $N_C = 6$ to guarantee the asymp-

otic freedom of the hypercolor gauge interaction. In the following, we restrict ourselves to the minimum case $N_C = 6$.

Imposing a global $U(1)_A$ symmetry

$$Q^I \rightarrow Q^I; Q^{11} \rightarrow e^{2i} Q^{11}; H_I \rightarrow e^{2i} H_I; S_{IJ} \rightarrow S_{IJ}; \quad (1)$$

we forbid such (mass) terms as $H_I H_I$ and $Q^{11} Q^{11}$ in the superpotential. Notice that this global $U(1)_A$ has an anomaly — it is broken by quantum effects. However, the broken $U(1)_A$ still plays a crucial role to guarantee the masslessness of Higgs doublets [9].

The tree-level superpotential is, then, given by

$$W = \sum_{I,J} Q^I Q^J S_{IJ} + m_Q Q^I Q^I + h Q^I Q^{11} H_I + \frac{1}{2} m_S \text{Tr}(S^2) + \frac{1}{3} m_S \text{Tr}(S^3); \quad (I, J = 1, \dots, 10); \quad (2)$$

Classically there is an $SO(10)_{GUT}$ -unbroken vacuum, whereas it does not exist quantum mechanically. If $\langle S_{IJ} \rangle = 0$, the low-energy physics below the scale $m_Q \neq 0$ would be effectively described by an $SO(6)_H$ gauge theory with one massless hyperquark Q^{11} . For this case ($N_f = N_C = 5$), however, there is no stable SUSY vacuum [11]. Here, N_f denotes the number of flavors of massless hyperquarks.

We now show that there is indeed a desired SUSY vacuum in the present model. We consider vacua specified by

$$\langle S_{IJ} \rangle = v \begin{array}{c} 0 \\ \vdots \\ \frac{3}{2}1 \\ \vdots \\ \frac{3}{2}1 \\ \vdots \\ 1 \\ \vdots \\ 1 \\ \vdots \\ 1 \end{array}; \quad 1 = \begin{array}{cc} 0 & 1 \\ \vdots & \vdots \\ 1 & 0 \\ \vdots & \vdots \\ 0 & 1 \end{array}; \quad (3)$$

in which the $SO(10)_{GUT}$ is broken down to the Pati-Salam gauge group $SO(6) \times SU(2)_L \times SU(2)_R$ [12]. The vacuum expectation value v is dynamically chosen as $Q v = m_Q$ or $\frac{2}{3}m_Q$ so that there are seven or five massless hyperquarks, respectively [13]. We adopt the former vacuum, since the latter has a phenomenological problem [14].

In the vacua $v = m_Q = 0$, the first four hyperquarks Q^i ($i = 1; \dots; 4$) are massive and the rest Q^a ($a = 5; \dots; 11$) are massless. Therefore, we may treat this theory as an $SO(6)$ gauge theory with seven hyperquarks at low energies. Fortunately, the quantum moduli space is the same as the classical one for $N_f = N_C + 1$ in the $SO(N_C)_H$ gauge theory [11]. This allows us to analyze the vacua with the aid of the classical theory in Eq.(2), since the numbers of flavors and colors for the massless hyperquarks Q^a amount to $N_f = 7$ and $N_C = 6$, respectively.

It is now a straightforward task to seek all of the vacua with Eq.(3). The F-term flatness conditions for the vacua are given by

$$\begin{aligned} m_S S^{IJ} + S(S^2)^{IJ} - \frac{1}{10} \text{Tr}(S^2) S^{IJ} + Q^I Q^J - \frac{1}{10} \text{Tr}(Q^I Q^J) S^{IJ} &= 0; \\ 2(Q^I S_{IJ} + m_Q \delta_{IJ}) Q^J + h Q^{11} H_I &= 0; \\ h Q^I H_I &= 0; \\ h Q^I Q^{11} &= 0; \end{aligned} \quad (4)$$

From these equations together with the D-term flatness conditions, we find the following vacuum :

$$h Q^I i = v_Q \begin{array}{c} 0 \\ \vdots \\ 1 \\ \vdots \\ 1 \\ \vdots \\ 1 \end{array} \begin{array}{c} 1 \\ 0 \\ \vdots \\ 1 \\ \vdots \\ 1 \\ \vdots \\ 1 \end{array} \begin{array}{c} C \\ \vdots \\ A \end{array}; \quad h Q^{11} i = 0; \quad h H_I i = 0; \quad v_Q = \sqrt[4]{\frac{5(2 m_Q m_S + m_S^2)}{4 Q^3}}; \quad (5)$$

with Eq.(3). We take the Yukawa couplings $Q = 0$ and $v = \bar{v}$, assuming $m_S = m_Q$. The GUT scale $v = \bar{v}$ is suggested to be 10^6 GeV from the renormalization group analysis on the gauge coupling constants of the low-energy gauge groups [3].

In this vacuum, the Q^I condensation in Eq.(5) breaks the hypercolor $SO(6)_H$ gauge symmetry (i.e. a Higgs phase is realized) and the total gauge group is broken down to the $SO(6)_C \times SU(2)_L \times SU(2)_R$. Here, the unbroken $SO(6)_C$ is the diagonal subgroup of the hypercolor $SO(6)_H$ and the $SO(6)$ subgroup of $SO(10)_{GUT}$. The quark-lepton superfields

transform as $(4, 2, 1)$ and $(4, 1, 2)$ under this unbroken $SO(6)_C \times SU(2)_L \times SU(2)_R$, as they do in the original Pati-Salam model. The gauge coupling constant of the $SO(6)_C$ is given by

$$g_6 = \frac{g_{GUT}}{1 + \frac{g_{GUT}}{g_H}} : \quad (6)$$

In general $g_6 \neq g_{GUT}$ and hence we do not have the GUT unification of the gauge coupling constants of the low-energy gauge groups. However, as stressed before, the GUT unification is achieved naturally in a strong coupling limit of the hypercolor $SO(6)_H$, i.e. $g_H \rightarrow 1$ [15].

Since the masses m_Q and m_S in the superpotential in Eq.(2) are assumed to be of order the GUT scale, the vacuum expectation value v_Q of Q^a ($a = 5; \dots; 10$) also takes a value of the GUT scale. In view of Eq.(2), the colored Higgs H_a ($a = 5; \dots; 10$) obtain the GUT scale masses together with Q^{11} but the Higgs H_i ($i = 1; \dots; 4$) remain massless as long as $\langle Q^{11} \rangle = 0$. These massless Higgs multiplets transform as $(2, 2)$ under the $SU(2)_L \times SU(2)_R$. They are identified with two Higgs doublets in the SUSY standard model when the $SO(6)_C \times SU(2)_L \times SU(2)_R$ is broken down to the $SU(3)_C \times SU(2)_L \times U(1)_X$. The masslessness of H_i is guaranteed by the $U(1)_A$ symmetry in Eq.(1) [9]. On the other hand, the mass term for the colored Higgs H_a and Q^{11} is allowed since they have the opposite $U(1)_A$ charges each other. Notice that the presence of the vacuum with unbroken $U(1)_A$ in Eqs.(3) and (5) is a dynamical consequence of the present model.

We are now at a point to discuss further breaking of the remaining $SO(6)_C \times SU(2)_L \times SU(2)_R$ down to the $SU(3)_C \times SU(2)_L \times U(1)_X$. To this end, we introduce additional Higgs superfields A_{IJ} (45), (16), (16) and a singlet of the $SO(10)_{GUT}$ [16]. Their superpotential is given by

$$W^0 = m_A \text{Tr}(A^2) + \lambda_A \text{Tr}(A^2 S) + g_1 (\lambda_{IJ}) A_{IJ} + g_2 (\lambda^2) : \quad (7)$$

Then, the total superpotential is given by the sum of Eq.(2) and Eq.(7).

In order to show the presence of a desired vacuum, we consider the following forms of $\langle A_{IJ} \rangle$ and $\langle S \rangle$:

$$hA_{IJ}i = \begin{pmatrix} 0 & 1 \\ B & C \\ a_{i_2} & \\ & a_{i_2} \\ & b_{i_2} \\ & b_{i_2} \\ & b_{i_2} \end{pmatrix}; h i = v \begin{pmatrix} " & " & " & " & " \end{pmatrix}; \quad (8)$$

where [17]

$$i_2 = \begin{pmatrix} 0 & 1 \\ B & C \\ a & \\ b & \\ c & \end{pmatrix} : \quad (9)$$

The vacuum expectation values a, b, v, v_Q and satisfy equations

$$\begin{aligned} 5m_Q m_S Q^1 + 2_A (a^2 - b^2) - \frac{5}{2} m_Q^2 Q^2 + 2_Q v_Q^2 &= 0; \\ C_1 a - g v^2 &= 0; \\ C_2 b - g v^2 &= 0; \\ g (4a + 6b)v + g v &= 0; \\ g (v^2 - Q^2) &= 0; \end{aligned} \quad (10)$$

where we define

$$C_1 = 2m_A + 3_A m_Q Q^1; \quad C_2 = 2m_A - 2_A m_Q Q^1; \quad (11)$$

Here, we have used $v = m_Q = Q$ in Eq.(3). From these equations we find a vacuum

$$\begin{aligned} v_Q &= \frac{5}{2} m_Q m_S Q^2 + \frac{5}{4} m_Q^2 Q^3 - g^2 A Q^1 (C_1^2 - C_2^2)^{-1}; \\ a &= g^{-2} C_1^{-1}; \\ b &= g^{-2} C_2^{-1}; \\ &= 10g^2 A m_Q Q^1 + 2m_A (g C_1 C_2)^{-1}; \\ v^2 &= Q^2; \end{aligned} \quad (12)$$

where the total gauge group $SO(10)_{GUT} \rightarrow SO(6)_H$ is now broken down to just the standard gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$ at the GUT scale.

It is important that the H_i^{11} still vanishes as it does in Eq.(5). Thus, the masslessness of the Higgs doublets H_i is maintained in this natural vacuum.

Now, we proceed to the ordinary matter sector. We assign one unit of the $U(1)_A$ charge to the quark-lepton superfields so that the Higgs doublets H_i couple to them and produce their masses after spontaneous breaking of the $SU(2)_L \times U(1)_Y$ symmetry. Then, there are two issues about quark and lepton masses as in the minimal $SO(10)_{GUT}$ models: vanishing mixing angles in the quark sector and mass degeneracy of the neutrinos and up-type quarks.

These issues are resolved by taking into account nonrenormalizable interactions. Let us define the $U(1)_A$ transformation of ψ and ψ^c as

$$\psi \rightarrow e^{i\alpha} \psi; \quad \psi^c \rightarrow e^{-i\alpha} \psi^c; \quad (13)$$

Then, the superpotential of the model may contain nonrenormalizable interactions

$$W \supset \frac{f}{M_P} \psi \psi^c; \quad (14)$$

where M_P denotes the Planck scale as a cutoff and $\psi = (1, 2, 3)$ are the three families of quark-lepton multiplets. The vacuum expectation value of ψ of the GUT scale gives the right-handed neutrinos Majorana masses of the order of $f \sqrt{v^2} = M_P \sqrt{f} \sim 10^3$ GeV, resulting in small masses for the left-handed neutrinos [18]. Small masses of the quarks and leptons which generate sizable mixings may be also induced by nonrenormalizable interactions [19].

It might appear that the non-vanishing $U(1)_A$ charges of ψ 's would spoil the masslessness of the Higgs doublets H_i since their vacuum expectation values break the $U(1)_A$ symmetry. Fortunately enough, there remains an unbroken $U(1)$ symmetry, $U(1)_{A^0}$, which is a linear combination of the original $U(1)_A$ and a $U(1)$ subgroup of the $SO(10)_{GUT}$. The global $U(1)_A$ and $U(1)_{A^0}$ symmetries are physically equivalent since the $SO(10)_{GUT}$ is a gauge symmetry. Therefore, the Higgs H_i are kept massless by the unbroken $U(1)_{A^0}$ symmetry.

We conclude that the low-energy spectrum in the present vacuum is nothing other than that in the SUSY standard model with the right-handed neutrinos.

Now, several comments are in order.

First, the dangerous dimension-ve operators [20] which induce the proton decay are forbidden in our model by the $U(1)_{A^0}$ symmetry.

Second, the $SO(10)_{GUT}$ is not asymptotically free above the threshold of the GUT superfields such as S_{IJ} , A_{IJ} and Φ 's. However, the gauge coupling constant g_{GUT} may not blow up until the Planck scale even if all the GUT superfields have the GUT scale masses.

Third, a SUSY-invariant mass term of the Higgs doublets may be generated when the $U(1)_{A^0}$ symmetry is broken spontaneously at an intermediate scale [21].

Finally, we mention a possibility that the $SO(10)_{GUT}$ is broken down to the $SU(5)_{GUT} \times SO(6)_H$ theory at the Planck scale. Then our model can be regarded as the $SU(5)_{GUT} \times SO(6)_H$ theory up to the Planck scale. That is, we may construct another model based on the $SU(5)_{GUT} \times SO(6)_H$ gauge group which also achieves natural unification of the strong and electroweak interactions, as does the $SO(10)_{GUT} \rightarrow SO(6)_H$ model.

REFERENCES

- [1] H. Georgi and S.L. Glashow, *Phys. Rev. Lett.* **32**, 438 (1974).
- [2] H. Georgi, In *Particles and fields* {1974, (APS/DPF Williamsburg), ed. C.E. Carlson (AIP, New York, 1975) p.573; H. Fritzsch and P. Minkowski, *Ann. Phys. (N.Y.)* **93**, 193 (1975).
- [3] P. Langacker and M.-X. Luo, *Phys. Rev. D* **44**, 817 (1991); U. Amaldi, W. de Boer and H. Furstenau, *Phys. Lett. B* **260**, 447 (1991); J. Ellis, S. Kelley and D.V. Nanopoulos, *Phys. Lett. B* **260**, 131 (1991); W. J. Marciano, *Annu. Rev. Nucl. Part.* **41**, 469 (1991).
- [4] There have been several attempts to solve this problem. See E. Witten, *Phys. Lett. B* **105**, 267 (1981); A. Masiero, D.V. Nanopoulos, K. Tamvakis and T. Yanagida, *Phys. Lett. B* **115**, 380 (1982); B. Grinstein, *Nucl. Phys. B* **206**, 387 (1982); S. Dimopoulos and F. Wilczek, University of California at Santa Barbara, Institute for Theoretical Physics Report No. NSF-ITP-82-07, 1981 (unpublished); For recent models, see also K.S. Babu and S.M. Barr, *Phys. Rev. D* **50**, 3529 (1994).
- [5] There has been already an attempt for semisimple unification along the present line based on an $SU(5)_{GUT} \times SU(3)_H$ gauge group [6,7]. Although it retains phenomenological successes, it predicts deviation from the minimal $SU(5)$ SUSY-GUT. See also Ref. [8,9], which deal with a $U(1)_H$ gauge group sacrificing the automatic hypercharge quantization.
- [6] T. Hotta, Izawa K.-I. and T. Yanagida, hep-ph/9511431.
- [7] T. Hotta, Izawa K.-I. and T. Yanagida, hep-ph/9601320.
- [8] T. Yanagida, *Phys. Lett. B* **344**, 211 (1995).
- [9] T. Hotta, Izawa K.-I. and T. Yanagida, hep-ph/9509201 (to be published in *Phys. Rev. D*).

[10] Of course, one may construct an anomaly-free model choosing real representations of $SU(N_C)_H$ for the hyperquarks Q .

[11] K. Intriligator and N. Seiberg, Nucl. Phys. B 444, 125 (1995).

[12] J.C. Pati and A. Salam, Phys. Rev. D 8, 86 (1971).

[13] For $Q v \neq m_Q$ or $\frac{2}{3}m_Q$, there is no SUSY vacuum since the low-energy theory again has $N_f = 1$.

[14] In the vacuum $Q v = \frac{2}{3}m_Q$, the massless hyperquarks Q^{11} ($= 1; \dots; 4$) may be identified with a pair of Higgs doublets in the SUSY standard model. However, they do not have renormalizable Yukawa couplings to quarks and leptons [9].

[15] When the $SO(6)_C \times SU(2)_L \times SU(2)_R$ is broken down to the standard $SU(3)_C \times SU(2)_L \times U(1)_Y$ at the GUT scale, the gauge coupling constant of $SU(3)_C$ equals to g_6 there.

[16] We introduce only to break the $SO(10)_{GUT}$ down to the $SU(5)_{GUT}$. Instead of Δ , nonrenormalizable interactions such as $(\Delta^c)^2$ serve for that purpose.

[17] For the $SO(10)$ spinor notations, see J. Hisano, H. Murayama and T. Yanagida, Phys. Rev. D 49, 4966 (1994).

[18] T. Yanagida, in Proceedings of the Workshop on Unified Theory and Baryon Number in the Universe, ed. O. Sawada and A. Sugamoto, (KEK, Tsukuba, Japan, 1979), p.95; M. Gell-Mann, P. Ramond and R. Slansky, in Supergravity, Ed. F. van Nieuwenhuizen and D. Freedman (North Holland, 1979), p.315.

[19] For example, G. Anderson, S. Dimopoulos, L.J. Hall, S. Raby and G.D. Starkman, Phys. Rev. D 49, 3660 (1994).

[20] N. Sakai and T. Yanagida, Nucl. Phys. B 197, 533 (1982); S. Weinberg, Phys. Rev. D 26, 287 (1982).

[21] H. Murayama, H. Suzuki and T. Yanagida, Phys. Lett. B 291, 418 (1992). See also Ref.

[9].