

FTUV/96-19

IFIC/96-21

Nucleosynthesis constraints on heavy χ in the presence of annihilations to majorons

S. Pastor¹

Instituto de Física Corpuscular - C.S.I.C.
Departament de Física Teòrica, Universitat de València
46100 Burjassot, València, SPAIN

Contribution to the IV International Workshop
on Theoretical and Phenomenological Aspects of
Underground Physics, Toledo, Spain, 17-21 September 1995.

Abstract

We show that in the presence of sufficiently strong χ annihilations to majorons, primordial nucleosynthesis constraints can not rule out any values of the χ mass up to the present laboratory limit.

¹Supported by Conselleria d'Educació i Ciència of Generalitat Valenciana. Work done in collaboration with J.C. Romão (IST, Lisbon), A.D. Dolgov and J.W.F. Valle (IFIC, Valencia).

The tau-neutrino is the only one which can have mass in the MeV range. Among all studied limits on m_τ , the kinematical ones are the only model-independent bounds. The last result from ALEPH [1] is

$$m_\tau < 23 \text{ MeV} \quad (1)$$

In addition to this, there are stronger bounds from cosmological considerations. From the contribution of stable τ to the present relic density one gets $m_\tau < 92 h^2 \text{ eV}$ [2], where $h = H_0/(100 \text{ km s}^{-1} \text{ Mpc}^{-1})$. This means that a massive τ with mass in the MeV range must be unstable with lifetimes smaller than the age of the Universe.

Moreover, if massive τ 's are stable during nucleosynthesis (lifetime longer than 100 sec), one can constrain their contribution to the total energy density, using the observed amount of primordial helium. This bound can be expressed through an effective number of massless neutrino species N_{eff} . Using $N_{\text{eff}} < 3.4$ [3, 6], the mass range

$$0.5 \text{ MeV} < m_\tau < 35 \text{ MeV} \quad (2)$$

has been excluded [3, 4]. This forbids all masses on the few MeV range. However, such masses are theoretically viable [5], and interesting for a possible solution of the structure formation problem as described in [6].

It is possible to weaken the constraints on masses of (2), by adding new interactions of τ beyond the standard ones. One may consider either that the τ 's are unstable during nucleosynthesis [7] or that they possess new channels of annihilation beyond the standard ones. The last case is what we have considered.

We studied the effect of the presence of Majorana annihilations to majorons (J). The interaction term of the Lagrangian that describes the coupling between neutrinos and majorons can be expressed [5] as $L_{\nu J} = g J^\text{T} \nu$, where g is a 3×3 matrix containing all possible couplings. In particular for the main processes are those described in figure (1). We assume that the coupling constants g_{23} (non-diagonal) and g_{33} (diagonal) are such that during nucleosynthesis time only annihilations are important. In previous works only the decay processes were taken into account, though in general g_{23} and g_{33} may be simultaneously important.

We consider the interactions of massive τ 's stable during the epoch just before nucleosynthesis. The τ 's interact with leptons via the standard weak interactions, $\tau \rightarrow \nu_\tau, e^+ e^-$, as in [3, 4]. We assume that, in addition,




Figure 1: Non-diagonal and diagonal couplings of Majorana tau neutrinos and Majorana jons.

the ν_τ 's annihilate to Majorana jons via the diagonal coupling

$$\mathcal{L} = g J^T \nu_\tau \quad (3)$$

where $g = g_{33}$ and ν_τ represents a two-component Majorana spinor. The corresponding elastic processes do not change the particle densities, but as long as they are effective they maintain all species with the same temperature.

Even after the ν_τ 's have decoupled from standard weak interactions, they remain in contact with Majorana jons. Therefore, as happens with photons and neutrinos when the e^+e^- pairs annihilate, after the decoupling there are two plasmas with different temperatures, one constituted of ν_τ 's and J 's and the other of the rest of the particles.

The evolution of the number densities of tau neutrinos (n_ν) and Majorana jons (n_J) is given by the solution of a set of Boltzmann differential equations

$$\begin{aligned} \dot{n}_\nu + 3H n_\nu &= \sum_{i=e,\mu} h_{i\nu} (n_i^2 - (n_i^{\text{eq}})^2) \\ &\quad - h_{J\nu} (n_\nu^2 - (n_\nu^{\text{eq}})^2 \frac{n_J^2}{(n_J^{\text{eq}})^2}) \end{aligned} \quad (4)$$

$$\dot{n}_J + 3H n_J = h_{J\nu} (n_\nu^2 - (n_\nu^{\text{eq}})^2 \frac{n_J^2}{(n_J^{\text{eq}})^2}) \quad (5)$$

where $\langle \sigma v \rangle$ is the thermally averaged cross section of the annihilation process of ν 's to particle-type β], where $\beta = e; \mu; J$. In the above equations we assumed Boltzmann statistics and $n_i = n_i^{\text{eq}}$ for $i = e; \mu$.

Now let us briefly describe the calculations. First we normalized the number densities to the number density of massless neutrinos, $n_0 = 0.18 T^3$. We introduced $r = n/n_0$, where $n = \nu; J$, and the corresponding equilibrium functions r^{eq} . On the other hand we performed the integrations using the dimensionless variable $x = m/T$. The set of equations (4) and (5) is completed with the evolution of the temperature, which is obtained from one of Einstein's equations

$$\dot{\rho} = -3H(\rho + P) : \quad (6)$$

Here ρ and P are the energy density and pressure of tau neutrinos and majorns, respectively.

We have integrated numerically the coupled differential equations in (4-6), obtaining the solutions of r for each pair of values $(m; g)$. The initial conditions are, for sufficiently high temperatures, $r = r^{\text{eq}}$, $r_J = r_J^{\text{eq}}$ and $T = T_J = T_0$.

The value of $r(m; g)$ is used to estimate the variation of total energy density $\rho_{\text{tot}} = \rho_R + \rho_\nu$. In ρ_R all relativistic species are taken into account, including majorns and two massless neutrinos, whereas ρ_ν is the energy density of massive ν 's.

We can calculate now the effective number of massless neutrino species (N_{eff}) corresponding to each $r(m; g)$. First we run a program which calculates the evolution of the neutron fraction (x_n), as presented e.g. in [9], varying the value of N_{eff} . Then we incorporate ρ_{tot} to the same program and perform the integration for each pair of values $(m; g)$. Comparing x_n obtained in each case at $T = 0.065 \text{ MeV}$ (the moment when practically all neutrons are wound up in ^4He), we can relate $(m; g)$ and N_{eff} .

The results are shown in figure (2). One can see that for a fixed $N_{\text{eff}}^{\text{max}}$, a wide range of tau neutrino masses is allowed for large enough coupling constants g . One sees that all masses below 23 MeV are allowed by nucleosynthesis arguments, provided that the coupling between ν 's and J 's exceeds a value of a few times 10^{-4} . Such values are reasonable for many majron models [5]. A more detailed study of this problem will be presented elsewhere [10].

Our conclusion is that the constraints on the mass of a Majrona from primordial nucleosynthesis can be relaxed if annihilations $\nu \bar{\nu} \rightarrow JJ$ are present.




Figure 2: Effective number of massless neutrinos equivalent to the contribution of massive neutrinos with different values of g expressed in units of 10^{-5} . For comparison, the dashed line corresponds to the case when $g = 0$ and no Majorons are present.

References

- [1] D .Buskulic et al., Phys. Lett. B 349 (1995) 585.
- [2] S.S.Gerstein and Ya.B .Zeldovich, JETP Lett. 4 (1966) 17; R .Cowsik and J.M cC lelland, Phys. Rev. Lett. 29 (1972) 669.
- [3] E.W .Kolb, M .S.Turner, A .Chakravorty and D.N .Schramm , Phys. Rev. Lett. 67 (1991) 533.
- [4] A.D .Dolgov and I.Z .Rothstein, Phys. Rev. Lett. 71 (1993) 476.
- [5] J.W .F.Valle, proceedings of TAUP 95, Toledo (Spain), September 1995, Nucl. Phys. B (Proc. Suppl.) , in press.
- [6] A.D .Dolgov, proceedings of TAUP 95, Toledo (Spain), September 1995, Nucl. Phys. B (Proc. Suppl.) , in press.
- [7] M .Kawasaki, G .Steigman and H.-S.Kang, Nucl. Phys. B 402 (1993) 323, B 419 (1994) 105; S.Dodelson, G .Gyuk and M .S.Turner, Phys. Rev. D 49 (1994) 5068.
- [8] P.Gondolo and G .Gelmini, Nucl. Phys. B 360 (1991) 145.
- [9] D.A .Dicus et al., Phys. Rev. D 26 (1982) 2694.
- [10] A.D .Dolgov, S.Pastor, J.C .Romao and J.W .F.Valle, in preparation.