

KEK-TH-473

SNU-TP-96-025

YUMS-96-005

(Modified version)

Average Kinetic Energy of Heavy Quark in Semileptonic B Decay

Dae Sung Hwang¹, C. S. Kim^{2,3} and Wuk Namgung⁴

1: Dept. of Physics, Sejong Univ. Seoul 143, Korea (dshwang@phy.sejong.ac.kr)

2: Dept. of Physics, Yonsei Univ. Seoul 120, Korea (kim@cskim.yonsei.ac.kr)

3: Theory Division, KEK, Tsukuba, Ibaraki 305, Japan (cskim@kek.vax.kek.jp)

4: Dept. of Physics, Dongguk Univ. Seoul 100, Korea (ngw@cakra.dongguk.ac.kr)

Abstract

Within the ACCMM model the average kinetic energy of heavy quark in a heavy-light meson is calculated as $\langle p_F^2 \rangle = \frac{3}{2} p_F^2$, solely from the fact that the Gaussian momentum probability distribution has been taken in the ACCMM model. Therefore, the Fermi momentum parameter p_F of the ACCMM model is not a truly free parameter, but is closely related to the average kinetic energy of heavy quark, which is theoretically calculable in principle. In this context, we determine p_F by comparing the theoretical prediction of the ACCMM model with the model independent lepton energy spectrum of $B \rightarrow e X$ from the recent CLEO analysis, and find that $p_F = 0.54^{+0.16}_{-0.15}$ GeV. We also calculate p_F in the relativistic quark model by applying the quantum mechanical variational method, and obtained $p_F = 0.5 - 0.6$ GeV. We show the correspondences between the relativistic quark model and the heavy quark effective theory. We then clarify the importance of the value of p_F in the determination of $\langle \bar{\psi} \psi \rangle_{ub} = V_{cb}^2$.

1. Introduction

In the standard $SU(2) \times U(1)$ gauge theory of Glashow, Salam and Weinberg the fermion masses and hadronic flavor changing weak transitions have a somewhat less secure role, since they require a prior knowledge of the mass generation mechanism. The simplest possibility to give mass to the fermions in the theory makes use of Yukawa interactions involving the doublet Higgs field. These interactions give rise to the Cabibbo-Kobayashi-Maskawa (CKM) matrix: Quarks of different flavor are mixed in the charged weak currents by means of a unitary matrix V . However, both the electromagnetic current and the weak neutral current remain flavor diagonal. Second order weak processes such as mixing and CP violation are even less secure theoretically, since they can be affected by both beyond the Standard Model virtual contributions and new physics direct contributions. Our present understanding of CP violation is based on the three-family Kobayashi-Maskawa model [1] of quarks, some of whose charged current couplings have phases. Over the past decade, new data have allowed one to refine our knowledge about parameters of this matrix V .

In the minimal Standard Model CP violation is possible through the CKM mixing matrix of three families, and it is important to know whether the element V_{ub} is non-zero or not accurately. Its knowledge is also necessary to check whether the unitarity triangle is closed or not [2]. However, its experimental value is very poorly known presently and its better experimental information is urgently required. At present, the only experimental method to measure V_{ub} is through the end-point lepton energy spectrum of the inclusive B-meson semileptonic decays, e.g. CLEO [3] and ARGUS [4], and their data indicate that V_{ub} is non-zero. Recently it has also been suggested that the measurements of hadronic invariant mass spectrum [5,6] as well as hadronic energy spectrum [7] in the inclusive $B \rightarrow X_{c(u)} l$ decays can be useful in extracting $|V_{ub}|$ with better theoretical understandings. In

future asymmetric B factories with vertex detector, the hadronic invariant mass spectrum will offer alternative ways to select $b \rightarrow u$ transitions that are much more efficient than selecting the upper end of the lepton energy spectrum, with much less theoretical uncertainties.

The simplest model for the semileptonic B -decay is the spectator model which considers the decaying b -quark in the B -meson as a free particle. The spectator model is usually used with the inclusion of perturbative QCD radiative corrections [8]. Then the decay width of the process $B \rightarrow X_{q1} l$ is given by

$$\Gamma_B(B \rightarrow X_{q1} l) = |V_{qb}|^2 \Gamma_B^e(B \rightarrow X_{q1} l) \\ \Gamma_B^e(b \rightarrow q1 l) = |V_{qb}|^2 \frac{G_F^2 m_b^5}{192 \pi^3} f(z) \left[1 - \frac{2}{3} g(z) \right] \frac{m_q}{m_b} ; \quad (1)$$

where m_q is the mass of the final q -quark decayed from b -quark. Here $f(z)$ is the phase-space factor, and $g(z) = (1 - 3z^2)(1 - z)^2 + 1.5$ is the corresponding single gluon QCD correction [9]. As can be seen, the decay width of the spectator model depends on m_b^5 , therefore small difference of m_b would change the decay width significantly. The model of Altarelli et al. [10] (ACCM model) is an improvement on the naive free-quark decay spectator model, but at the cost of introducing several free parameters: the final (charm) quark mass m_c , the spectator mass m_{sp} , and the most important Fermi momentum function $\langle p; p_F \rangle$ that includes both binding and final state interaction effects.

In Section 2, we determine the Fermi momentum parameter p_F by comparing the theoretical prediction of the ACCM model with the model independent lepton energy spectrum of $B \rightarrow X_{cl}$ for the whole region of electron energy, which has been recently extracted by CLEO [11]. Previously, the comparison had been hampered by the cascade decay of $b \rightarrow c \rightarrow sl$, and only the part of lepton energy spectrum ($E_l > 1.8 \text{ GeV}$) could be compared to give $p_F = 0.3 \text{ GeV}$. However, we argue that the value $p_F = 0.3 \text{ GeV}$, which has been commonly used in experimental analyses, has no theoretical or experimental clear justification. Therefore, it is strongly recommended to determine the value of p_F more reliably and inde-

pendently, when we think of the importance of its role in experimental analyses. A better determination of p_F is also interesting theoretically since it has its own physical correspondence related to the average kinetic energy ($\langle p^2 \rangle$) of heavy quark inside heavy meson. In this context we calculate theoretically the value of $\langle p^2 \rangle$ in the relativistic quark model using quantum mechanical variational method in Section 3. We also compare our model with the heavy quark effective theory (HQET) in expansion of $1/M_Q$. The value of p_F is particularly important in the determination of the value of $\langle V_{ub} \rangle = \langle V_{cb} \rangle$ as we explain in Section 4. Section 5 contains our conclusions.

2. Determination of p_F from the Experimental Spectrum

Altarelli et al: [10] proposed for the inclusive B-meson semileptonic decays their ACCMM model, which incorporates the bound state effect by treating the b-quark as a virtual state particle, thus giving momentum dependence to the b-quark mass. The virtual state b-quark mass W is given by

$$W^2(p) = m_B^2 + m_{sp}^2 - 2m_B^q \frac{p^2 + m_{sp}^2}{p^2 + m_{sp}^2} \quad (2)$$

in the B-meson rest frame, where m_{sp} is the spectator quark mass, m_B is the B-meson mass, and p is the momentum of the b-quark inside B-meson.

For the momentum distribution of the virtual b-quark, Altarelli et al: considered the Fermi motion inside the B-meson with the Gaussian momentum probability distribution

$$(p; p_F) = \frac{4}{p_F^3} e^{-p^2/p_F^2}; \quad (3)$$

where the Gaussian width, p_F , is treated as a free parameter. Then the lepton energy spectrum of the B-meson decay is given by

$$\frac{d_B}{dE_1}(p_F; m_{sp}; m_q; m_B) = \int_0^{p_{max}} p^2 dp (p; p_F) \frac{d_b}{dE_1}(m_b = W; m_q); \quad (4)$$

where p_{max} is the maximum kinematically allowed value of $p = |\mathbf{p}|$. The ACCMM model, therefore, introduces a new parameter p_F for the Gaussian momentum distribution of the b -quark inside B -meson, instead of the b -quark mass of the spectator model. In this way the ACCMM model incorporates the bound state effects and reduces the strong dependence on b -quark mass in the decay width of the spectator model.

The Fermi momentum parameter p_F is the most essential parameter of the ACCMM model, as we explained in the above. However, the experimental determination of its value from the lepton energy spectrum has been very ambiguous, because various parameters of the ACCMM model, such as p_F , m_q and m_{sp} , are fitted all together from the limited region of end-point lepton energy spectrum ($E_l > 1.8 \text{ GeV}$) to avoid the cascade decay of $b \rightarrow c \rightarrow s l$, and because the perturbative QCD corrections are very sensitive in the end-point region of the spectrum. Recently, CLEO [11] extracted the model independent lepton energy spectrum of $B \rightarrow X_c l$ for the whole region of electron energy from 2.06 fb^{-1} of (4S) data, which is shown in Fig. 1, with much smaller uncertainties compared to the previously measured results of ARGUS [12]. Now we compare the whole region of experimental electron energy spectrum of CLEO with the theoretical prediction of the ACCMM model, Eq. (4), to derive the value of p_F using χ^2 analysis. With p_F , m_c and m_{sp} as free parameters, for one standard deviation we obtain

$$p_F = 0.54^{+0.16}_{-0.15} \text{ GeV} ; \quad (5)$$

In Table I, we show the extracted values of p_F (in GeV) and $\chi^2_{\text{min}}/\text{d.o.f.}$ for the fixed input values of $m_{\text{sp}} = 0; 0.15 \text{ GeV}$ and $m_q = m_c = 1.4; 1.5; 1.6; 1.7 \text{ GeV}$, which are the values commonly used in experimental analyses. As can be noticed, these results are strongly dependent on the input value of m_c : if we use smaller m_c , the best fit value of p_F increases, and vice versa. In Fig. 1, we also show the theoretical ACCMM model spectrum with $p_F = 0.44; 0.51; 0.59 \text{ GeV}$ (with $m_c = 1.5 \text{ GeV}$, $m_{\text{sp}} = 0.0 \text{ GeV}$), corresponding to dashed-, full-, dotted-line, respectively.

The experimental data and the theoretical predictions are all normalized to the semileptonic branching ratio, $BR(B \rightarrow X_{cl}) = 10.49\%$, following the result of CLEO [11]. Previously, we extracted similarly p_F by comparing the theoretical prediction with the experimental spectrum of ARGUS [12], and we obtained [13] $p_F = 0.27^{+0.22}_{-0.27}$ GeV for the fixed input values of $m_c = 1.5$ GeV and $m_{sp} = 0.15$ GeV. As can be seen from Table I, if we fix $m_c = 1.5$ GeV and $m_{sp} = 0.15$ GeV, then we obtain from the new CLEO spectrum [11] $p_F = 0.55^{+0.09}_{-0.07}$ GeV with the minimum χ^2 being about 1.0. We note that two results are apart each other within one standard deviation, but the new result from CLEO has much smaller uncertainties. In Sections 3 and 4, we give in detail the related physics of this unexpected large value of the parameter p_F .

3. Average Kinetic Energy of Heavy Quark inside Heavy Meson

Recently considerable progresses have been achieved on the relation of the ACCMM model with QCD [14{16]. Especially Bigi et al. [14] derived an inequality between the expectation value of the kinetic energy of the heavy quark inside the hadron and that of the chromomagnetic operator, which gives

$$\langle p^2 \rangle \geq \frac{3}{4} (M_V^2 - M_P^2) : \quad (6)$$

The experimental value of the right hand side of Eq. (6) is 0.36 GeV^2 for B-meson system [18]. This bound corresponds to $p_F = 0.49$ GeV for B-meson, because in the ACCMM model the average kinetic energy, $\langle p^2 \rangle$, can be calculated from

$$\langle p^2 \rangle = \int_0^1 dp p^2 \rho(p; p_F) = \frac{3}{2} p_F^2 : \quad (7)$$

This theoretical lower bound could be significantly weakened, as shown in [17], with inclusion of the α_s corrections as well as $1/M_Q$ corrections.

This relation (7) was obtained solely from the fact that the Gaussian momentum probability distribution was taken in the ACCMM model, and therefore the lower bound $p_F = 0.49 \text{ GeV}$ is independent of any other input parameter values of the ACCMM model, and is much larger than the commonly used value $p_F = 0.3 \text{ GeV}$. Ballet al: [16] also calculated $\langle p^2 \rangle$ using the QCD sum rule approach, and obtained $\langle p^2 \rangle = 0.50 \pm 0.10 \text{ GeV}^2$ for B-meson, corresponding to $p_F = 0.58 \pm 0.06 \text{ GeV}$ from Eq. (7). We note that the heavy quark inside the hadron possesses more kinetic energy than the value one might expect naively from the nonrelativistic consideration. We also note that the Fermi momentum parameter p_F of the ACCMM model is not a truly free parameter, but is closely related to the average kinetic energy of heavy quark, which is theoretically calculable in principle.

We consider the relativistic potential model with the quantum mechanical variational technique to theoretically calculate the average kinetic energy of b-quark inside B-meson, and to compare the results with the predictions of the HQET. The potential model has been successful to describe the physics of ψ and χ families with the nonrelativistic Hamiltonian [19,20]. However, for B-meson it has been difficult to apply the nonrelativistic potential model because of the relativistic motion of the light quark inside B-meson. In this work, we study B-meson system with a realistic Hamiltonian, which is relativistic for the light quark and nonrelativistic for the heavy quark, and adopt the variational method to solve it. We take the Gaussian function as the trial wave function, and obtain the ground state energy and wave function by minimizing the expectation value of the Hamiltonian.

For the B-meson system we start with the Hamiltonian

$$H = M + \frac{p^2}{2M} + \frac{q}{p^2 + m^2} + V(r); \quad (8)$$

where $M = m_b$ is the heavy quark mass and $m = m_{sp}$ is the u- or d-quark mass (which corresponds to the spectator light quark mass in the ACCMM model). We apply the variational method to the Hamiltonian (8) with the trial wave function

$$\psi(r) = \left(\frac{1}{\zeta}\right)^{3/2} e^{-\frac{1}{2\zeta} r^2}; \quad (9)$$

where the parameter ζ is a variational parameter. The Fourier transform of $\psi(r)$ gives the momentum space wave function $\psi(p)$, which is also Gaussian,

$$\psi(p) = \frac{1}{(\zeta\pi)^{3/2}} e^{-\frac{1}{2\zeta} p^2}; \quad (10)$$

We note here that the Gaussian momentum probability distribution of the ACCMM model equals $\langle p; p_f \rangle = 4 \int \langle p; j \rangle^2$. See Eqs. (3) and (10). The ground state is given by minimizing the expectation value of H ,

$$\frac{d}{d\zeta} \langle H \rangle = 0; \quad \frac{d}{d\zeta} \langle H \rangle = 0 \text{ at } \zeta = \zeta_f; \quad (11)$$

and then the value $E = \langle H \rangle$ approximates the B-meson mass M_B , and at the same time we get p_f , the Fermi momentum parameter in the ACCMM model. As is well known, the value of ζ or p_f corresponds to the measure of the radius of the two body bound state, as can be seen from the relation, $\langle r^2 \rangle = 3 \langle p^2 \rangle^{-1}$ or $\langle r^2 \rangle^{1/2} = 3 \langle p^2 \rangle^{-1/2}$.

We now take in Eq. (8) the Cornell potential, which is composed of the Coulomb and linear potentials with a constant term,

$$V(r) = -\frac{c}{r} + K r + V_0 = -\frac{4}{3} \frac{s}{r} + K r + V_0; \quad (12)$$

The additive constant V_0 , which is related to the regularization concerned with the linear confining potential [21], is usually known as flavor dependent: $V_0 = 0$ for heavy-heavy meson system, $V_0 = -0.2 \text{ GeV}$ for B-meson system [22]. We use the value of $K = 0.19 \text{ GeV}^2$ [23] for the string tension, and for the parameter $c = (\frac{4}{3} s)$ we will consider two values $s = 0.35$ and 0.24 separately. The first choice $s = 0.35$ is the value which has been determined by the best fit of (cc) and (cb) bound state spectra [23], and $s = 0.24$ is that given by the running coupling constant for the QCD scale at M_B .

With the Gaussian trial wave functions, (9) and (10), the expectation value of each term of the Hamiltonian (8) is given as follows:

$$\begin{aligned} \langle \frac{p^2}{2M} \rangle &= \langle \psi | \frac{p^2}{2M} | \psi \rangle = \frac{3}{4M} \alpha^2 ; \\ \langle \frac{1}{p^2 + m^2} \rangle &= \langle \psi | \frac{1}{p^2 + m^2} | \psi \rangle = \frac{4}{\pi} \int_0^{\infty} e^{-x^2} \frac{1}{x^2 + (m/\alpha)^2} x^2 dx ; \\ \langle V(r) \rangle &= \langle \psi | -\frac{c}{r} + K r + V_0 | \psi \rangle = \frac{2}{\pi} (-c \alpha + K \alpha) + V_0 : \quad (13) \end{aligned}$$

Then we have

$$\begin{aligned} E(\alpha) &= \langle H \rangle \\ &= M + \frac{1}{2M} \frac{3}{2} \alpha^2 + \frac{2}{\pi} (-c \alpha + K \alpha) + V_0 + \frac{4}{\pi} \int_0^{\infty} e^{-x^2} \frac{1}{x^2 + (m/\alpha)^2} x^2 dx : \end{aligned} \quad (14)$$

In our previous study [24], we obtained the last integral in Eq. (14) as a power series of $(m/\alpha)^2$. And when we write up to the order of $(m/\alpha)^4$, we now get

$$\begin{aligned} E(\alpha) &= M + \frac{3}{4M} \alpha^2 + \frac{2}{\pi} (-c \alpha + K \alpha) + V_0 \\ &\quad + \frac{2}{\pi} \left(1 + \frac{1}{2} (m/\alpha)^2 + \frac{5}{32} - 2c_1 (m/\alpha)^4 + \frac{1}{4} (m/\alpha)^4 \ln(m/\alpha) \right. \\ &\quad \left. + O((m/\alpha)^6) \right) ; \end{aligned} \quad (15)$$

where $c_1 = 0.0975$. Up to the order of $(m/\alpha)^2$, $E(\alpha)$ becomes

$$E(\alpha) = M + \frac{3}{4M} \alpha^2 + \frac{2}{\pi} ((1 - c) \alpha + (K + \frac{1}{2} m^2) \alpha) + V_0 ; \quad (16)$$

and the next order terms ($O((m/\alpha)^4)$) contribute only less than 1%. Then, we find the minimum value of $E(\alpha)$ in (16) by the variational method, and the minimum point is given by

$$\frac{\partial}{\partial \alpha} E(\alpha) = \frac{3}{2M} \alpha + \frac{2}{\pi} ((1 - c) + (K + \frac{1}{2} m^2)) = 0 ; \quad (17)$$

where

$$(1 - c) = 1 - \frac{4}{3} s ; \quad \text{and} \quad K + \frac{1}{2} m^2 : \quad (18)$$

We rewrite Eq. (17) as

$$(1 - c) + \frac{b}{M} \alpha^3 = 0 ; \quad (19)$$

where $b = 3^{\frac{p}{2}} = 4$ is a constant. Then, we expand ψ , which satisfies Eq. (19), as a power series of $1/M$,

$$\psi = a_0 + a_1 \frac{1}{M} + a_2 \frac{1}{M^2} + \dots; \quad (20)$$

and by matching the order by the order in (19), we get

$$a_0 = \frac{s}{2}; \quad a_1 = \frac{b}{2} \frac{s}{2}; \quad a_2 = \frac{5b^2}{8} \frac{s}{2} - \frac{s}{3}; \quad \dots; \quad (21)$$

As can be easily seen, since $bM \ll 1$, Eq. (19) has an approximate solution $\psi \approx a_0$.

Using Eqs. (20) and (21), we can obtain the numerical values of the coefficients a_0, a_1, a_2 , and that of ψ which minimizes $E(\psi)$ in Eq. (16), for $s = 0.35$ and 0.24 separately. We also considered three different values of the light quark mass $m(m_{sp}) = 0.00; 0.15; 0.30 \text{ GeV}$, in order to see the dependence of the results on the light quark mass. As we can see from (17) and (18), the effect of m comes in only through the little modification of K , because $K + m^2 = 2K$. The results of this calculation for a_0, a_1, a_2 and ψ with the input values of s and the light quark mass $m(m_{sp})$ are presented in Table II. As previously explained, we fixed $K = 0.19 \text{ GeV}^2$ and $V_0 = 0.2 \text{ GeV}$. However, the exact value of V_0 is irrelevant in our calculations of ψ , (20) and (21), but it is necessary for the calculation of E in Eq. (23) below.

With ψ of (20) and (21), we can get the following expectation values of the terms in the Hamiltonian (8):

$$\frac{T}{2M} = \frac{\hbar^2 \langle \psi | \nabla^2 | \psi \rangle}{2M} = \frac{3}{4M}$$

^yThe numerical value of ψ is fairly insensitive to the potential we choose. In Ref. [25], ψ has been calculated numerically from six different potential models, and found to be $\psi = 0.56 \pm 0.02 \text{ GeV}$, where the error is only the statistical error of the six different results.

$$= \frac{1}{2M} \left[\frac{3}{2} - \frac{3b^2}{2} - \frac{1}{2} \frac{1}{M} + \frac{9b^2}{4} - \frac{1}{4} \frac{1}{M^2} \right] + O\left(\frac{1}{M^3}\right); \quad (22)$$

$$\begin{aligned} \langle \frac{1}{h^2} \frac{1}{p^2 + m^2 + V(r)} \rangle &= \frac{1}{2M} \left(\frac{1}{2} + \frac{1}{2} \right) + V_0 \\ &= V_0 + \frac{1}{2} \left(\frac{1}{M} + \frac{b^2}{4} - \frac{1}{2} \frac{1}{M^2} \right) + O\left(\frac{1}{M^3}\right); \end{aligned} \quad (23)$$

Finally, $E(\mu)$ in (16) is expressed as a power series in $1/M$,

$$\begin{aligned} E(\mu) &= M + \frac{T}{2M} \\ &+ V_0 + \frac{1}{2} \left(\frac{1}{M} + \frac{b^2}{4} - \frac{1}{2} \frac{1}{M^2} \right) + O\left(\frac{1}{M^3}\right); \end{aligned} \quad (24)$$

In Eq. (24), the M -independent terms come from $\langle \frac{1}{h^2} \frac{1}{p^2 + m^2 + V(r)} \rangle$, which can be considered as the contributions from the light degrees of freedom. The term of the order of $1/M$ is from the heavy quark momentum squared $\langle p^2 \rangle$, that is, from the average kinetic energy of the heavy quark inside the heavy-light meson. Both $\langle \frac{1}{h^2} \frac{1}{p^2 + m^2 + V(r)} \rangle$ and $\langle p^2 \rangle$ contribute to the term of the order of $1/M^2$. In the HQET, the mass of a heavy-light meson is represented [26] by

$$M_M = M + \frac{1}{2M} (T + \mu) + O\left(\frac{1}{M^2}\right); \quad (25)$$

where $\lim_{M \rightarrow \infty} (M_M - M)$ is the contribution from the light degrees of freedom, for which Neubert obtained [26] $\mu = 0.57 \pm 0.07$ GeV. $T = \langle p^2 \rangle$ is the expectation value of the kinetic energy of the heavy quark (up to $2M$) inside a heavy-light meson, and μ is the expectation value of the energy due to the chromomagnetic hyperfine interaction with $\gamma_v = 1/4$ and $\gamma_p = 3/4$. In this paper we do not consider the chromomagnetic hyperfine interaction term. We will present a detailed study on the correspondences between the relativistic quark model and the heavy quark effective theory in another forthcoming paper [27]. Here we calculated only T and μ up to the order of $1/M^2$ by using (22) and (23), and obtained the values shown in Table III. In Table III, we also show the values of the Fermi momentum

parameter p_F (; shown in Table II) of the ACCMM model using the relation (7).

Gremm et al: [28] recently extracted the average kinetic energy, $T = \langle p^2 \rangle$, by comparing the prediction of the HQET [29] with the shape of the inclusive $B \rightarrow X_l$ lepton energy spectrum [30] for $E_l = 1.5 \text{ GeV}$, in order to avoid the contamination from the secondary leptons of cascade decays of $b \rightarrow c \rightarrow s$. They obtained $\langle p^2 \rangle(T) = 0.35 \pm 0.05 \text{ GeV}^2$ for $|V_{ub}|=|V_{cb}| = 0.08$ and $\langle p^2 \rangle(T) = 0.37 \pm 0.05 \text{ GeV}^2$ for $|V_{ub}|=|V_{cb}| = 0.1$, which correspond to $p_F = 0.48 \pm 0.03 \text{ GeV}$ and $p_F = 0.50 \pm 0.03 \text{ GeV}$, respectively. Their results are remarkably close to the our value in (5) extracted from the recent model independent lepton energy spectrum of $B \rightarrow X_c l$ [11], as explained in Section 2.

We summarize Section 3 by noting that the value of the Fermi momentum parameter of the ACCMM model is $p_F = 0.5 \pm 0.6 \text{ GeV}$ and is much larger than 0.3 GeV , as can be seen from Table III, and the heavy quark inside the hadron possesses much more kinetic energy than the value one might expect naively from the nonrelativistic consideration.

4. Dependence of $|V_{cb}|$ and $|V_{ub}|=|V_{cb}|$ on the Average Kinetic Energy of Heavy Quark inside B-meson

Now we consider the dependence on the average kinetic energy of b-quark (or equivalently Fermi momentum parameter p_F of the ACCMM model) in the B-meson semileptonic decay, $\langle p^2 \rangle$, of the measurements of $|V_{cb}|$ and $|V_{ub}|=|V_{cb}|$. The B-meson inclusive branching fraction is related to the CKM matrix V_{cb} and V_{ub} by

$$\text{BR}(B \rightarrow X_l) = \frac{e_c^2 |V_{cb}|^2}{e_B} + \frac{e_u^2 |V_{ub}|^2}{e_B} = \frac{e_c^2 |V_{cb}|^2}{e_B}; \quad (26)$$

where the factors $\frac{e_q}{e_B}(B \rightarrow X_{q1})(p_F)$ must be calculated from theory. (See Eq. (1).) CLEO has extracted $|V_{cb}| = 0.040 \pm 0.001 \pm 0.004$ from their measurements [11] of

$$\text{BR}(B \rightarrow X \ell) = (10.49 \pm 0.17 \pm 0.43) \% ;$$

$$\tau_B = (1.61 \pm 0.04) \text{ psec} ; \quad (27)$$

and by assuming $\tau_c = (39 \pm 8) \text{ psec}^{-1}$. If we instead theoretically calculate τ_c in the ACCMM model by using $p_F = 0.5 \pm 0.6 \text{ GeV}$, the result of the ACCMM model becomes

$$|V_{cb}| = |V_{cb}|_{\text{CLEO}} \frac{\tau_c^{(\text{CLEO})}}{\tau_c^{(p_F = 0.5 \pm 0.6)}} = |V_{cb}|_{\text{CLEO}} \cdot 1.1 = 0.044 \pm 0.001 \pm 0.004 ; \quad (28)$$

We can easily understand this large correction ($\sim 10\%$) in $|V_{cb}|$ due to the change in p_F , because within ACCMM model from Eqs. (1,2)

$$\tau_c / m_b^5 = W^5 (m_B^2 - 2m_B p_F)^{5/2} ;$$

and therefore $\tau_c^{(p_F = 0.3)} = \tau_c^{(p_F = 0.5)} \cdot 1.25 ; \quad (29)$

The ACCMM model also provides an inclusive lepton energy spectrum of the B-meson semileptonic decay to obtain the value of $|V_{ub}| = |V_{cb}|$. The lepton energy spectrum is useful in separating $b \rightarrow u$ transitions from $b \rightarrow c$, since the end-point region of the spectrum is completely composed of $b \rightarrow u$ decays. In applying this method one integrates (4) in the range $2.3 \text{ GeV} < E_\ell$ at the B-meson rest frame, where only $b \rightarrow u$ transitions exist [31]. So we theoretically calculate²

$$\tau_c^{(p_F)} = \int_{2.3}^Z dE_\ell \frac{d\Gamma_B}{dE_\ell}(p_F; m_{sp}; m_q; m_B) ; \quad (30)$$

In (30) we specified only p_F dependence explicitly in the left-hand side. Then one compares the theoretically calculated $\tau_c^{(p_F)}$ with the experimentally measured

²We note that the dependences of the lepton energy spectrum on perturbative and non-perturbative QCD corrections [8,29] as well as on the unavoidable specific model parameters (e.g. the parameter p_F of the ACCMM model [10]) are strongest at the end-point region of the inclusive $d\Gamma/dE_\ell$ distribution. Therefore, Eq. (30) may have very limited validity for the determination of $|V_{ub}| = |V_{cb}|$ as shown in [32].

width e_{exp} in the region $2.3 \text{ GeV} < E_1$, to extract the value of $|V_{ub}|$ from the relation

$$e_{\text{exp}} = |V_{ub}|^2 e(p_F) : \quad (31)$$

In the real experimental situations [3,4,12,31], the only measured quantity is the number of events in this region of high E_1 compared to the total semileptonic events number, i.e. the branching-fraction $e_{\text{exp}} = \frac{e_{\text{total}}}{s.l.}$. Since the value $\frac{e_{\text{total}}}{s.l.}$ is proportional to $|V_{cb}|^2$, only the combination $|V_{ub}=V_{cb}|^2$ is extracted.

We now consider the possible dependence of $|V_{ub}=V_{cb}|^2$ as a function of the parameter p_F from the following relation

$$\frac{e_{\text{exp}}}{e_{\text{total}}^{s.l.}} / \frac{|V_{ub}|^2}{|V_{cb}|^2}_{p_F=p_F} e(p_F) = \frac{|V_{ub}|^2}{|V_{cb}|^2}_{p_F=0.3} e(p_F=0.3) ; \quad (32)$$

where $|V_{ub}=V_{cb}|^2_{p_F=p_F}$ is determined with an arbitrary value of the Fermi momentum parameter p_F . In the right-hand side we used $p_F = 0.3 \text{ GeV}$ because this value is commonly used in the experimental determination of $|V_{ub}=V_{cb}|$. Then one can get a relation

$$\frac{|V_{ub}|}{|V_{cb}|}_{p_F=p_F} = \frac{|V_{ub}|}{|V_{cb}|}_{p_F=0.3} \sqrt{\frac{e(0.3)}{e(p_F)}} : \quad (33)$$

We numerically calculated theoretical ratio $e(0.3)=e(p_F)$ by using (4) and (30) with $m_{sp} = 0.15 \text{ GeV}$, $m_q = m_u = 0.15 \text{ GeV}$, which are the values commonly used by experimentalists, and $m_B = 5.28 \text{ GeV}$. We show the values of $|V_{ub}(p_F)=V_{ub}(p_F=0.3)|$ as a function of p_F in Fig. 2. If we use $p_F = 0.5 - 0.6 \text{ GeV}$, instead of $p_F = 0.3 \text{ GeV}$, in the experimental analysis of the end-point region of lepton energy spectrum, the value of $|V_{ub}=V_{cb}|$ becomes significantly changed.

Previously the CLEO [31] analyzed with $p_F = 0.3 \text{ GeV}$ the end-point lepton energy spectrum to get

$$\begin{aligned} |V_{ub}=V_{cb}| &= 0.76 - 0.08 \text{ (ACMM with } p_F = 0.3 \text{ [31])} ; \\ &= 1.01 - 0.10 \text{ (Isgur et al. (ISGW) [33])} : \end{aligned} \quad (34)$$

As can be seen, those values differ by two standard deviations^{*}. However, if we use $p_F = 0.5 - 0.6$ GeV, the result of the ACCMM model becomes

$$10 \leq |V_{ub}=V_{cb}| \leq 1.07 - 0.11 \text{ (ACCMM with } p_F = 0.5 - 0.6); \quad (35)$$

and these two models are in a good agreement for the value of $|V_{ub}=V_{cb}|$.

We note here that the dependence of $|V_{ub}=V_{cb}|$ on the parameter p_F is much stronger compared to that of $|V_{cb}|$. This is because the p_F dependence of the inclusive distribution $d\Gamma/dE_1$ is particularly sensitive if we restrict ourselves only in the limited region of end-point, as shown in Eq. (30). We would like to emphasize again that the measurements of the hadronic invariant mass spectrum [5,6] in the inclusive $B \rightarrow X_{c(u)} l$ decays can be much more useful in extracting $|V_{ub}|$ with better theoretical understandings, where we can use almost the whole region of decay spectrum: i.e. in the forthcoming asymmetric B -experiments with microvertex detectors, BABAR and BELLE, the total separation of $b \rightarrow u$ semileptonic decays from the dominant $b \rightarrow c$ semileptonic decays would be experimentally viable using the measurement of inclusive hadronic invariant mass distributions. And we could determine the ratio of CKM matrix elements $|V_{ub}=V_{cb}|$ from the ratio of those measured total integrated decay rates [6], which is theoretically described by the phase space factor and the well-known perturbative QCD correction only.

5. Conclusions

The value of the Fermi momentum parameter of the ACCMM model $p_F = 0.3$

^{*} There now exists an improved version of ISGW model, so-called ISGW 2 [34], which gives a considerably harder end-point spectrum than that of ISGW. Therefore, it seems clear that the prediction of ISGW on $|V_{ub}=V_{cb}|$, Eq. (34), will decrease when re-analyzed by experimentalists, even though the changes would be small [34].

GeV, which has been commonly used in experimental analyses, has no theoretical or experimental clear justification. Therefore, it is strongly recommended to determine the value of p_F more reliably and independently, when we think of the importance of its role in experimental analyses. It is particularly important in the determination of the value of $\langle V_{ub}=V_{cb} \rangle$. We note that the dependence of $\langle V_{ub}=V_{cb} \rangle$ on the parameter p_F is very strong, because the inclusive lepton energy distribution is particularly sensitive to the variation of p_F if we restrict ourselves only in the limited region of end-point. A better determination of p_F is also interesting theoretically since it has its own physical correspondence related to the average kinetic energy $\langle p^2 \rangle$ of the heavy quark inside B-meson. Within the ACCMM model the average kinetic energy is calculated as $\langle p^2 \rangle = \frac{3}{2} p_F^2$, solely from the fact that the Gaussian momentum probability distribution has been taken in the ACCMM model. Therefore, the Fermi momentum parameter p_F of the ACCMM model is not a truly free parameter, but is closely related to the average kinetic energy of heavy quark, which is theoretically calculable in principle.

In this context we theoretically calculated the value of p_F in the relativistic quark model using quantum mechanical variational method. It turns out that $p_F = 0.5 - 0.6$ GeV, which is consistent with the value of p_F determined by comparing the ACCMM model prediction and the model independent lepton energy spectrum of the CLEO measurement, $p_F = 0.54^{+0.16}_{-0.15}$ GeV. We note that the value of the Fermi momentum parameter of the ACCMM model is much larger than 0.3 GeV, and the heavy quark inside the hadron possesses much more kinetic energy than the value one might expect naively from the nonrelativistic consideration. We also found the correspondences between the relativistic quark model and the heavy quark effective theory by the $1/M_Q$ expansion, and the result shows that they are consistent with each other.

If we use $p_F = 0.5 - 0.6$ GeV, instead of $p_F = 0.3$ GeV, in the experimental analysis of the end-point region of lepton energy spectrum, the value of $\langle V_{ub}=V_{cb} \rangle$

is increased by the factor of 1:3 ~ 1:5 compared with the case of $p_e = 0.3 \text{ GeV}$. Here we would like to emphasize that the measurements of the hadronic invariant mass spectrum in the inclusive $B \rightarrow X_{c(u)} l$ decays can be much more useful in extracting $|V_{ub}|$ with better theoretical understandings. In future asymmetric B factories with vertex detector, the hadronic invariant mass spectrum will offer alternative ways [5,6] to select $b \rightarrow u$ transitions that are much more efficient than selecting the upper end of the lepton energy spectrum, with much less theoretical uncertainties.

Acknowledgements

The work was supported in part by the Korean Science and Engineering Foundation, Project No. 951-0207-008-2, in part by Non-Directed-Research-Fund, Korea Research Foundation 1993, in part by the CTP, Seoul National University, in part by Yonsei University Faculty Research Grant, in part by Daeyang Foundation at Sejong University, in part by the Basic Science Research Institute Program, Ministry of Education 1997, Project No. BSR I-97-2425, and in part by COE fellowship of the Japanese Ministry of Education, Science and Culture.

REFERENCES

- [1] M . Kobayashi and T . Maskawa, Prog. Theor. Phys. 49, 652 (1973).
- [2] For example, see C . S . Kim , J . L . Rosner and C . P . Yuan, Phys. Rev. D 42, 96 (1990); H . Quinn, Nucl. Phys. B (Proc. Suppl.) 37A , 21 (1994).
- [3] CLEO Collab., R . Fulton et al., Phys. Rev. Lett. 64, 16 (1990).
- [4] ARGUS Collab., H . Albrecht et al., Phys. Lett. B 234, 409 (1990); *ibid*: B 241, 278 (1990); *ibid*: B 255, 297 (1991).
- [5] V . Barger, C . S . Kim and R . J . N . Phillips, Phys. Lett. B 235, 187 (1990); *ibid*: B 251, 629 (1990); C . S . Kim , D . S . Hwang, P . Ko and W . Namgung, Nucl. Phys. B (Proc. Suppl.) 37A , 69 (1994); Phys. Rev. D 50, 5762 (1994).
- [6] C . S . Kim , hep-ph/9605201 (1996).
- [7] A . O . Bouzas and D . Zappala, Phys. Lett. B 333, 215 (1994).
- [8] N . Cabibbo, G . Corbo and L . M aiani, Nucl. Phys. B 155, 93 (1979); G . Corbo, *ibid*. B 212, 99 (1983); M . Jezabek and J . H . Kuhn, *ibid*: B 320, 20 (1989).
- [9] C . S . Kim and A . D . Martin, Phys. Lett. B 225, 186 (1989).
- [10] G . Altarelli, N . Cabibbo, G . Corbo, L . M aiani and G . Martinelli, Nucl. Phys. B 208, 365 (1982).
- [11] CLEO Collab., B . Barish et al., Phys. Rev. Lett. 76, 1570 (1996).
- [12] ARGUS Collab., H . Albrecht et al., Phys. Lett. B 318, 397 (1993).
- [13] Dae Sung Hwang, C . S . Kim and W uk Namgung, Phys. Rev. D 53, 4951 (1996).
- [14] I . I . Bigi, M . A . Shifman, N . G . Uraltsev and A . I . Vainshtein, Int. J. of Mod. Phys. A 9, 2467 (1994); Phys. Lett. B 328, 431 (1994); Phys. Rev. D 52, 196 (1995).

- [15] C. Csaki and L. Randall, Phys. Lett. B 324, 451 (1994).
- [16] P. Ball and V M. Braun, Phys. Rev. D 49, 2472 (1994); E. Bagan, P. Ball, V M. Braun and P. Gosdzinsky, Phys. Lett. B 342, 362 (1995).
- [17] A. Kapustin, Z. Ligeti, M B. Wise and B. Grinstein, hep-ph/9602262 (1996).
- [18] Particle Data Group: K. Hikasa et al., Phys. Rev. D 50, 1173 (1994).
- [19] E. Eichten, K. Gottfried, T. Kinoshita, K D. Lane and T M. Yan, Phys. Rev. D 17, 3090 (1978); *ibid*: D 21, 313 (E) (1980); *ibid*: D 21, 203 (1980).
- [20] C. Quigg and J.L. Rosner, Phys. Rept. 56, 167 (1979); E J. Eichten and C. Quigg, Phys. Rev. D 49, 5845 (1994); *ibid*: D 52, 1726 (1995).
- [21] W. Lucha, F F. Schoberl and D. Gromes, Phys. Rept. 200, 127 (1991).
- [22] L P. Fulcher, Phys. Rev. D 50, 447 (1994).
- [23] K. Hagiwara, A D. Martin and A W. Peacock, Z. Phys. C 33, 135 (1986).
- [24] D S. Hwang, C S. Kim and W. Namgung, Z. Phys. C 69, 107 (1995).
- [25] D S. Hwang and G .H. Kim, Phys. Rev. D 53, 3659 (1996).
- [26] M. Neubert, Phys. Rev. D 46, 1076 (1992); Phys. Rept. 245, 259 (1994).
- [27] D S. Hwang, C S. Kim and W. Namgung, in preparation.
- [28] M. Gremm, A. Kapustin, Z. Ligeti and M B. Wise, hep-ph/9603314 (1996).
- [29] I. I. Bigi, M A. Shifman, N G. Uraltsev and A. I. Vainshtein, Phys. Rev. Lett. 71, 496 (1993); A V. Manohar and M B. Wise, Phys. Rev. D 49, 1310 (1994); B. Blok, L. Koyrakh, M. Shifman and A. I. Vainshtein, Phys. Rev. D 49, 3356 (1994); A. Falk, M. Luke and M. Savage, Phys. Rev. D 49, 3367 (1994); T. Mannel, Nucl. Phys. B 423, 396 (1994); M. Neubert, Phys. Rev. D 49, 3392 (1994).

- [30] CLEO Collab., J. Bartelt et al.; CLEO/CONF 93-19 (1993).
- [31] CLEO Collab., J. Bartelt et al.; Phys. Rev. Lett. 71, 4111 (1993).
- [32] R. Akhouri and I.Z. Rothstein, hep-ph/9512303 (1995).
- [33] N. Isgur, D. Scora, B. Grinstein and M. B. Wise, Phys. Rev. D 39, 799 (1989).
- [34] D. Scora and N. Isgur, Phys. Rev. D 52, 2783 (1995).

TABLES

TABLE I. The values of p_F (in GeV) and $\chi^2_{\text{min}}/\text{dof}$ for the fixed input parameter values m_{sp} and m_c (in GeV). We derived the values using χ^2 analysis by comparing the whole region of experimental electron energy spectrum of CLEO [11], which is shown in Fig. 1, with the theoretical prediction of ACCMM model, Eq. (4) using p_F as a free parameter.

m _{sp} = 0.00					m _{sp} = 0.15					
m _c = 1.4 1.5 1.6 1.7					m _c = 1.4 1.5 1.6 1.7					
p _F	0.64	0.09	0.51 ^{+0.08} _{0.07}	0.40 ^{+0.07} _{0.05}	0.29 ^{+0.07} _{0.06}	0.69	0.10	0.55 ^{+0.09} _{0.07}	0.44 ^{+0.09} _{0.06}	0.32 ^{+0.08} _{0.03}
χ ² _{m in}	1.09		1.00	1.41	2.05	1.44		1.05	1.09	1.47

TABLE II. The numerical values of the coefficients a_0, a_1, a_2 in the $1/M$ expansion of \mathcal{E} , Eq. (20), and the values of α which minimizes $\mathcal{E}(\alpha)$ in Eq. (16). We varied $\alpha_s = 0.35, 0.24$ and the light quark mass $m(\bar{m}_{sp}) = 0.00, 0.15, 0.30$ GeV.

		a_0	a_1	a_2	
$\alpha_s = 0.35$	$m_{sp} = 0.00$	0.60	0.60	1.50	0.54
	$m_{sp} = 0.15$	0.61	0.63	1.62	0.54
	$m_{sp} = 0.30$	0.67	0.76	2.13	0.61
$\alpha_s = 0.24$	$m_{sp} = 0.00$	0.53	0.36	0.63	0.49
	$m_{sp} = 0.15$	0.54	0.38	0.68	0.49
	$m_{sp} = 0.30$	0.59	0.46	0.89	0.54

TABLE III. The average kinetic energy T (up to $2M$) of the heavy quark, the contribution of the light degrees of freedom ϵ , and the Fermi momentum parameter p_F of B -meson system, for $s = 0.35, 0.24$ and $m_c(m_{sp}) = 0.00, 0.15, 0.30$ GeV. The results obtained by the χ^2 analysis of the recent CLEO lepton energy spectrum, and those from the HQET and the QCD sum rule approaches are also presented.

		T	p_F ()	
$s = 0.35$	$m_{sp} = 0.00$	0.45	0.44	0.54
	$m_{sp} = 0.15$	0.47	0.46	0.54
	$m_{sp} = 0.30$	0.57	0.52	0.61
$s = 0.24$	$m_{sp} = 0.00$	0.36	0.52	0.49
	$m_{sp} = 0.15$	0.38	0.54	0.49
	$m_{sp} = 0.30$	0.45	0.61	0.54
from CLEO data [11]				$0.54^{+0.16}_{-0.15}$
Bigi et al: [14]		0.36		0.49
Ball et al: [16]		$0.50^{+0.10}_{-0.10}$		$0.58^{+0.06}_{-0.06}$
Neubert [26]			$0.57^{+0.07}_{-0.07}$	
Gremm et al: [28]	$V_{ub}=V_{cb}= 0.08$	$0.35^{+0.05}_{-0.05}$		$0.48^{+0.03}_{-0.03}$
Gremm et al: [28]	$V_{ub}=V_{cb}= 0.10$	$0.37^{+0.05}_{-0.05}$		$0.50^{+0.03}_{-0.03}$




Fig. 1 The normalized lepton energy spectrum of $B \rightarrow X_c l$ for the whole region of electron energy from the recent CLEO measurement [11]. Also shown are the theoretical ACCMM model predictions, Eq. (4), using $p_F = 0.44; 0.51; 0.59$ GeV, corresponding to dashed-, full-, dotted-line, respectively. The minimum χ^2 equals to 1.00 with $p_F = 0.51$ GeV. We fixed $m_{sp} = 0.0$ GeV and $m_q = m_c = 1.5$ GeV.




Fig. 2 The ratio $|V_{ub}(p_F)/V_{ub}(p_F = 0.3)|$ as a function of p_F .