

Large Transverse Momentum Jet Production and DIS Distributions of the Proton

M. Klasen, G. Kramer
II. Institut für Theoretische Physik*
Universität Hamburg
D - 22761 Hamburg, Germany

Abstract

We have calculated the single jet inclusive cross section as measured at Fermilab in next-to-leading order QCD using recent parton distributions of the CTEQ collaboration. We studied the scheme dependence of the jet cross section by employing the $\overline{\text{MS}}$ and DIS factorization schemes consistently. For $E_T > 200$ GeV, we find that the cross section in the DIS scheme is larger than in the $\overline{\text{MS}}$ scheme yielding a satisfactory description of the CDF data over the whole E_T range in the DIS scheme.

Recently, the CDF collaboration presented a precise measurement of the inclusive differential cross section for jet production in $p\bar{p}$ collisions at 1.8 TeV [1]. The measurement was compared to next-to-leading order (NLO) perturbative QCD predictions [2] for jet transverse energies E_T in the range of 15 to 440 GeV in the central pseudorapidity region $0.1 \leq |\eta| \leq 0.7$, using a selection of commonly used parton distributions. The experimental results for $E_T > 200$ GeV show evidence of a possible deviation as compared to the NLO prediction based on the current sets of parton distributions, which are obtained from global analyses [3, 4, 5, 6] of deep inelastic lepton-nucleon scattering and related data. Before explanations for this deviation based on new physics [7] are seriously considered, it is crucial to study possible explanations within the Standard Model.

Clearly, the perturbative predictions of the jet cross section depend on the parton distributions. Then the question arises whether the parton distributions can be adjusted to accommodate the jet measurements while still having a good description of the data sets as used in previous global analyses. This vital question has been addressed by two groups. Huston et al. [8] have carried out a new global QCD analysis incorporating the CDF inclusive jet data with

*Supported by Bundesministerium für Forschung und Technologie, Bonn, Germany under Contract 05 6HH93P(5) and EEC Program "Human Capital and Mobility" through Network "Physics at High Energy Colliders" under Contract CHRX-CT93-0357 (DG12 COMA)

$E_T > 75$ GeV (the very precise data in the low E_T range are excluded due to potential theoretical and experimental problems) and reached the conclusion, that there is enough flexibility in NLO global analyses to enhance the high E_T inclusive jet cross section by 25 – 35% above previous calculations. A similar analysis was done by Glover et al. [9]. They also incorporated the CDF single jet inclusive measurement in a global NLO parton analysis of the available deep inelastic and related data. In particular, they included the jet data in the region $E_T > 50$ GeV in their fit. They find that it is impossible to accommodate both the jet data over the complete E_T range and the deep inelastic structure function data. However, the CDF data for $E_T < 200$ GeV and the deep inelastic data were reasonably well fitted simultaneously. A fit over the full E_T range yields quarks which are completely incompatible with the large- x structure function data, a value for α_s , which is larger than the current world average, and a renormalization of the CDF jet data, which is barely compatible with the allowed range [1]. So from these two studies it still seems to be unclear, whether the difference between the CDF inclusive jet cross section data and the NLO predictions can be attributed to insufficient knowledge of the parton distributions.

It is well known that there are other uncertainties of the perturbative calculations which should be considered. First, we have the dependence on the renormalization and factorization scales. Second, changes in the strong coupling α_s resulting in changes of Λ_{QCD} must be considered. These uncertainties have been discussed in [8] and mainly affect the renormalization and not the shape of the inclusive jet cross section. Third, there exists the dependence of the perturbative predictions on the factorization scheme. To our knowledge, this dependence of the single jet cross section has not been investigated yet.

Beyond leading order, the parton distribution functions $f^a(x, \mu^2)$ depend on the factorization scheme. In applications to physical processes, the scheme chosen for the parton distributions must match that for the hard scattering cross section in the parton model formula [10]. The same parton distributions in two different schemes differ because of two effects. First, the scheme influences the evolution kernel. Second, the functional dependence at the starting scale $\mu = Q_0$ differs since the parton distributions are fitted to the same data with the evolution kernel for the appropriate scheme. Thus, they are functionally equivalent in the sense that they yield the same physical cross sections for the data included in the analysis. The $\overline{\text{MS}}$ parton distributions are guaranteed to satisfy the momentum sum rule. The DIS scheme [11] was defined specifically to make the relation between the parton distributions and F_2^γ as simple as possible by absorbing all the NLO terms into the definition of $f_{DIS}^q(x, \mu^2)$. This does not yet define the DIS prescription for the gluon distribution. Here, the convention has been adopted to require the momentum sum rule to be preserved in the DIS scheme as well. However, this requirement fixes only the second moment of the gluon distribution, so that one completes the definition by requiring the condition on the second moment to be valid for all moments. This convention was first introduced by Diemoz et al. [12] and is now generally applied for parton distribution function analyses in the DIS scheme.

Functionally equivalent parton distributions in the $\overline{\text{MS}}$ and the DIS schemes have been constructed by the CTEQ collaboration [4, 5]. The most recent ones are the sets CTEQ3M ($\overline{\text{MS}}$) and CTEQ3D (DIS), respectively [5]. These two sets are obtained from independent fits to the same data sets under the same assumptions except for calculating the evolution kernel. The total χ^2 for the two sets is supposed to be very similar. Unfortunately, this is not stated

explicitly in [5]. However, this fact was reported for the earlier sets CTEQ1M and CTEQ1D [4]. Thus, the two sets describe totally equivalent physics as defined by the data sets selected in the analysis. Now we can ask ourselves whether they also yield the same single jet inclusive cross sections or not.

There also exist DIS parton distributions from the other collaborations [3, 6]. In particular, MRS(D0') and MRS(D-') sets in the DIS scheme are available [3]. They are obtained from the corresponding sets in $\overline{\text{MS}}$, which were fitted to the deep inelastic and other data, by applying the $O(\alpha_s)$ perturbative transformation formula between the two schemes. It is known that this may be unreliable in situations where the NLO terms involving, for example, a large gluon contribution, are of comparable size as the LO term involving small sea quarks [5]. Furthermore, the DIS version of MRS is not equivalent to the Diemoz et al. description, since only the quark distribution functions are transformed. The GRV collaboration [6] also constructed a DIS set in their recent 1994 analysis (GRV(94)) by fitting $\overline{\text{MS}}$ and DIS parton distributions independently to the deep inelastic and other data. Unfortunately, this analysis is restricted to three flavors with the charm contribution to the deep inelastic structure function generated perturbatively. This makes this set less useful for calculations of other hard scattering cross sections. Therefore, we shall restrict ourselves to the CTEQ3 sets in the following.

Our calculation uses the NLO parton level program JETSAM [13] which originally was designed for the calculation of inclusive single jet cross sections in resolved photoproduction. This program was developed on the basis of the theory of Aversa et al. [15] which uses the phase space slicing method to cancel collinear divergences. The cuts and the algorithm for defining jets out of up to three partons are modelled as closely as possible to the experimental set-up. The jets are defined according to the Snowmass algorithm [14] with a jet cone size of $R = 0.7$ and lie in the pseudorapidity range between $0.1 \leq |\eta| \leq 0.7$ as in the CDF analysis. The factorization and renormalization scales are chosen to be $\mu_F = \mu_R = E_T/2$. The Monte Carlo JETSAM contains the necessary modifications of the NLO hard scattering cross sections to run it in the $\overline{\text{MS}}$ and the DIS scheme as defined above.

First, we compare the CDF jet data with the NLO prediction obtained from the CTEQ3M set of partons. The fractional difference between the data and the theoretical prediction is plotted in Fig. 1. The data are taken from Table 2 in [1]. Over the observed range of E_T , the experimental cross section decreases by more than a factor 10^8 and the quadratic sum of the *correlated* systematic uncertainties grows from $\pm 18\%$ at $E_T = 50$ GeV to $\pm 36\%$ at $E_T = 400$ GeV. In Fig. 1, the agreement of the data with the theory for the CTEQ3M set is measured by the distance of the data points from the horizontal line at zero. The normalization shown is absolute. These results show excellent agreement in shape and normalization for $E_T < 200$ GeV, where the cross section falls by six orders of magnitude. Above $E_T = 200$ GeV, the CDF cross section is significantly higher than the NLO cross section. This agrees with the statements made in ref. [1, 8, 9], where other parton sets in the $\overline{\text{MS}}$ scheme obtained from recent global analyses had been selected for the NLO calculation of the jet cross section. Thus, the agreement below 200 GeV and disagreement above 200 GeV seems to be a unique feature of all $\overline{\text{MS}}$ parton sets from recent global analyses which include both fixed-target and HERA deep inelastic data.




Figure 1: Fractional differences of the CDF inclusive jet data, the CTEQ3D prediction, and the CTEQ3D prediction without compensation terms from the CTEQ3M NLO QCD predictions.

In Fig. 1, we have also plotted the prediction for the CTEQ3D parton distributions. It is given by the dashed line which presents the fractional difference of the CTEQ3D cross section from the CTEQ3M cross section, i.e. $(\text{CTEQ3D}-\text{CTEQ3M})/\text{CTEQ3M}$. As we can see, this line deviates very little from the horizontal line (the CTEQ3M result) for $E_T < 150$ GeV, showing that the single inclusive jet cross section does not depend on the chosen factorization scheme in this E_T range. Above $E_T = 150$ GeV, the DIS prediction starts to deviate from the horizontal line and increases up to 0.4 at $E_T = 400$ GeV. Thus, in the large E_T range the jet cross section depends on the factorization scheme. For $E_T > 150$ GeV, the DIS cross section is larger than the $\overline{\text{MS}}$ cross section so that it agrees with all the large E_T data points except the last one which, however, has a large statistical error. In addition, we must take into account the systematic error of the experimental data [1].

In Fig. 1, we also show a second curve (dotted). This presents the fractional difference to the CTEQ3M prediction, obtained also with the DIS parton distribution set CTEQ3D, but without the appropriate modification of the hard scattering cross section to the DIS scheme, i.e. this cross section is left as calculated in the $\overline{\text{MS}}$ scheme. Of course, this is an inconsistent procedure. But it shows that the deviation of the DIS cross section from the $\overline{\text{MS}}$ cross section at $E_T = 400$ GeV, which is 40%, comes to a large extent from the modified parton distributions (25%) and to a lesser extent from the modified subtraction terms in the hard scattering cross section (15%). At the larger values of E_T , these two effects do not compensate each other as one might expect. At the smaller $E_T < 150$ GeV, this compensation occurs, the dashed curve is nearer to the horizontal line than the dotted one, although the difference between the two curves is very small.

We also performed the same calculations for the older sets CTEQ2M and CTEQ2D [5], respectively. The results are similar. The agreement between the data and the predictions with either the $\overline{\text{MS}}$ or DIS parton distributions in the $E_T < 150$ GeV is less satisfactory. The fractional difference of the CTEQ2D cross section from the CTEQ2M cross section shows the same increase with increasing E_T up to the value 0.5 at $E_T = 400$ GeV.

Fig. 2 compares the quark and gluon distributions at $Q^2 = 10$ and 10^4 GeV² of the two sets CTEQ3M and CTEQ3D. We plot the singlet quark distribution $\sum_a(q_a + \bar{q}_a)$ and the gluon dis-




Figure 2: The CTEQ3M and CTEQ3D parton distributions $\sum_a(q_a + \bar{q}_a)$ and g at $Q^2 = 10$ and 10^4 GeV².

tribution g multiplied by x^2 , so that the area under the curves is the total momentum fraction carried by the quarks and gluons. We see that for both Q^2 values, the DIS quarks are larger

(smaller) than the $\overline{\text{MS}}$ quarks for $x \geq 0.3$ ($x \leq 0.3$), whereas the DIS gluons are larger than the $\overline{\text{MS}}$ gluons over the whole x range of $0.01 < x < 1$. At smaller E_T , where the jet cross section receives contributions mainly from qg - and gg -initiated subprocesses [9], the compensation between the change of the parton distributions from $\overline{\text{MS}}$ to DIS and the modification of the hard scattering subprocesses seems to work reasonably well (see Fig. 1). For $E_T > 200$ GeV, the jet production is dominated by $q\bar{q}$ initiated subprocesses [9]. Since centrally produced jets of transverse energy E_T sample partons at $x \simeq x_T = \frac{2E_T}{\sqrt{s}}$, we can understand that the increased quark distribution for $x > 0.3$ in the DIS scheme leads to an increased jet cross section for $E_T > 250$ GeV in case that in this region the compensation is distorted due to the singling out of a particular subprocess. The increase of the gluon distribution in this region is small, so that the $q\bar{q}$ process is mainly responsible for the increase of the jet cross section. For smaller E_T , the effect of the larger gluon is compensated by a smaller quark distribution for $x < 0.3$.

In summary, we have studied the dependence of the inclusive jet cross section on the factorization scheme. We have considered two schemes, $\overline{\text{MS}}$ and DIS, for which parton distributions have been constructed in a global analysis by the CTEQ collaboration. We find that the inclusive jet cross section for $E_T > 200$ GeV in the DIS scheme is up to 40% larger than in the $\overline{\text{MS}}$ scheme and accounts well for the CDF inclusive jet cross section data in this E_T region as well as in the low E_T range.

Of course, the significant scheme dependence of the high E_T jet cross section is an artifact of finite order perturbation theory. In infinite order, this dependence is absent. From this point of view, our results indicate that the NLO theory is inadequate and one must go to the NNLO in order at least to diminish the scheme dependence and to obtain more reliable results. However, since parton distribution functions are involved, which are determined from other processes, the problem of the scheme dependence might have another solution. It is conceivable, that the parton distributions in the DIS scheme are the right ones with enough flexibility built in to describe all deep inelastic data together with the CDF jet data. Then, the task would be to find the corresponding $\overline{\text{MS}}$ set which describes the same data by perhaps changing the input parametrization of the distribution functions, in particular that of the gluon, which is much less constrained by deep inelastic data than the quark distribution. Such an investigation is beyond the intention of this work. We therefore conclude that it might still be possible that the difference between the CDF inclusive data and the NLO QCD predictions can be due to a deficiency in the way parton distributions in the $\overline{\text{MS}}$ scheme are constructed in the usual global analyses work as performed by the CTEQ, MRS, and GRV collaborations.

References

- [1] F. Abe et al., CDF collaboration, Fermilab preprint, FERMILAB-PUB-96/020-E (1996).
- [2] S. Ellis, Z. Kunszt, D. Soper, Phys. Rev. Lett. 62 (1989) 2188, ibid. 64 (1990) 2121.
- [3] A.D. Martin, R.G. Roberts, W.J. Stirling, Phys. Rev. D50 (1994) 6734, Phys. Lett. B354 (1995) 155.
- [4] J. Botts et al., CTEQ collaboration, Phys. Lett. B304 (1993) 159.
- [5] H.L. Lai et al., CTEQ collaboration, Phys. Rev. D51 (1995) 4763.

- [6] M. Glück, E. Reya, A. Vogt, *Z. Phys.* C67 (1995) 433.
- [7] V. Barger, M.S. Berger, R.J.N. Phillips, Madison preprint, MADPH-95-920 (1995), G. Altarelli, N. Di Bartolomeo, F. Fernglis, R. Gatto, M. Magnano, CERN preprint, CERN-TH-96-20 (1996), P. Chiappetta, J. Layssac, F.M. Renard, C. Verzegnassi, Marseille preprint, CPT-96-P-3304 (1996), M. Bander, UC Irvine preprint, UCI-TR-96-76 (1996).
- [8] J. Huston et al., CTEQ collaboration, Michigan State preprint, MSU-HEP-508/12, FSU-HEP-951031, CTEQ-512 (1995).
- [9] E.W.N. Glover, A.D. Martin, R.G. Roberts, W.J. Stirling, Durham preprint, DTP/96/22, RAL-TR-96-019 (1996).
- [10] For a recent review and original references, see: J. Collins, D. Soper, G. Sterman in: *Perturbative Quantum Chromodynamics*, A. Mueller (ed.), World Scientific, Singapore, 1989.
- [11] G. Altarelli, R.K. Ellis, G. Martinelli, *Nucl. Phys.* B143 (1978) 521, *ibid.* B157 (1979) 461. For an alternative DIS scheme, see [15].
- [12] M. Diemoz, F. Ferroni, E. Longo, G. Martinelli, *Z. Phys.* C39 (1988) 21.
- [13] S.G. Salesch, DESY 93-196 (1993), H1-05/95-441 (1995).
- [14] J.E. Huth et al. in: *Proceedings of the 1990 DPF Summer Study on High Energy Physics*, Snowmass, Colorado, E.L. Berger (ed.), World Scientific, Singapore, 1992, p. 134.
- [15] F. Aversa, M. Greco, P. Chiappetta, J.-Ph. Guillet, *Nucl. Phys.* B327 (1989) 205, *Z. Phys.* C46 (1990) 253.