HIGH ENERGY PROTON ACCELERATION IN DISSOLVING PLASMA CLOTS

V. DAVIDOVSKY and SH. SHVARTSMAN¹

Physics Department Case Western Reserve University Cleveland, OH 44106, USA

Abstract

We consider the acceleration of particles (protons) in a plasma clot by an induced field which appears due to the dissolving of the clot. The mechanism of this acceleration is based on a model where the clot consists of double flat moving current layers. It is assumed that the currents flow in opposite directions, while the clot moves in a perpendicular one. It is shown that protons can reach the highest observed energy (of the order of $10^{11} - 10^{12}$ GeV), providing that the size of the clot is large enough.

¹e-mail: sms25@po.cwru.edu

There is a well known problem in high energy cosmic rays physics of how one can explain the existence in the universe of protons with the energy in the range of $10^{11} - 10^{12}$ GeV. Unfortunately all existing models do not successfully describe this phenomena.

In this paper we are going to discuss a mechanism that might explain the existence of particles which have such high energies. This mechanism assumes that somewhere in the universe there exist very large and sufficiently dense plasma clots. Such objects recently were discovered in space, indeed these are the SS433 objects [1] and the jets from the active galactic nuclei [2]. We also assume that inside the clots there exists an internal current as it will be described below.

Plasma clots usually posses collective electromagnetic field. This field may be produced by either a collective electric charge or a current inside the clot. The collision of the clot's particles with the medium, as well as the internal recombination in the cooling clot, are responsible for dissolving of the moving clot. Therefore, the collective electromagnetic field decreases, and the energy and the momentum of the vanishing field can be distributed among the rest of the clot particles. This means that the energy of the collective field can be transferred to a comparatively small number of particles, and those particles can reach high energy. This effect might explain the origin of cosmic ray protons with the highest observed energies. Recently, some events were discovered where the energies are larger than 10^{11} GeV [3, 4, 5, 6].

As the first step toward the justification of the above proposed acceleration mechanism, we shall show that the energy spectrum of accelerating particles is close to the observed one. For this purpose it is natural to assume that the number of particles, say N_p , in the clot decreases exponentially in time

$$N_p = N_{0\,p} \exp(-\alpha t) \tag{1}$$

where N_{0p} is the initial number of particles in the clot and α is a slowly varying function of time. Then, during the time interval dt the loss in the clot energy, dE, can be presented as

$$dE = \epsilon dN_p \tag{2}$$

where ϵ is the sum of the kinetic energy of a particle and a portion of the collective field energy per one particle at a given instant of time. A part of energy dE is transferred to the rest of the clot particles, such that each of them receives the energy portion $d\epsilon$

$$d\epsilon = -k\frac{dE}{N_p} = k\alpha\epsilon dt \tag{3}$$

where k is a proper fraction. This gives

$$\epsilon = \epsilon_0 \exp(\alpha kt) \tag{4}$$

Then a simple relation holds

$$\frac{N_p}{N_0 p} = \left(\frac{\epsilon}{\epsilon_0}\right)^{-\frac{1}{k}} \tag{5}$$

This spectrum is close to the observed one [7] if the parameter k is of the order of one-half to two-thirds. This shows that a k-portion $(k \simeq 0.5)$ of the energy of the clot's collective field per one particle can be transferred to the rest of the particles inside the clot.

To estimate the maximum of the energy the particles (protons) can gain, let us consider a simple model as follows. We assume that the collective electromagnetic field is produced by a current inside the clot. This current consists of the two double flat current layers flowing in opposite directions, say x-direction. The current inside the single layer is composed by both the electrons that have the velocity $v_x^{(e)}$ and protons that have the velocity $v_x^{(p)}$. We also suppose that both layers as whole system is moving in the perpendicular direction, say the z-one, with the velocity v_z relative to the medium.

Due to the above proposed model the current density in the reference frame K' where the current layers are at rest can be expressed in the following form

$$j'_{x} = \begin{cases} j', & \text{if } 0 < z' < L' \\ -j', & \text{if } -L' < z' < 0 \\ 0, & \text{if } |z'| > L' \end{cases}$$
(6)

where L' is the effective width of the layer.

The corresponding vector-potential $\vec{A'}$ and magnetic field $\vec{B'}$ are (we present only the none zero components)

$$A'_{x} = \begin{cases} \mu_{0}j'z'(L'-z'/2), & \text{if } 0 < z' < L' \\ \mu_{0}j'z'(L'+z'/2), & \text{if } -L' < z' < 0 \\ \mu_{0}j'L'^{2}/2, & \text{if } z' > L' \\ -\mu_{0}j'L'^{2}/2, & \text{if } z' < -L' \end{cases}$$

$$B'_{y} = \begin{cases} \mu_{0}j'(L'-z'), & \text{if } 0 < z' < L' \\ \mu_{0}j'(L'+z'), & \text{if } -L' < z' < 0 \\ 0, & \text{if } |z'| > L' \end{cases}$$

$$(7)$$

Due to the collisions with the medium particles part of the electrons and protons inside the current layers live the clot. Because the plasma clot remains quasineutrality both number of particles that leave the clot are the same. Therefore current density j'_x decreases. In the reference frame K' medium particles have the velocity v_z directed opposite to the z-axis. This velocity v_z is the one of the current layer in the laboratory reference frame K where the medium is at rest. In the reference frame K' the concentration of the clot electrons n'inside the layer can be written in the following form

$$n' = n'_0 \exp(-\sigma N' v_z t') \tag{9}$$

where n'_0 is the initial concentration of the clot electrons in the layer, N' is the concentration of the particles in the medium, and σ is the scattering cross section of an electron inside the clot by the medium particles. Here we neglect the thermal motion of the electrons inside the clot.

Let us now pass to the laboratory reference frame K. The component A'_x of the vector potential (7) does not transform because it is the transverse component of a 4-vector (as

well as the current density j), although the variables it depends upon, do. Then this vector potential in the reference frame K has the form

$$A_x^{\pm} = \frac{\mu_0}{2} j(z - v_z t) [2L \mp (z - v_z t)] \gamma_z^3, \tag{10}$$

$$j = en'_0 v_x \exp\{-\sigma N v_z (t - \frac{v_z}{c^2} z) \gamma_z^2\}, \qquad (11)$$

$$\gamma_z = (1 - \frac{v_z^2}{c^2})^{-\frac{1}{2}} \tag{12}$$

where the sign + corresponds to the region $0 < z - v_z t < L$, while the sign - corresponds to the region $-L < z - v_z t < 0$. The corresponding electric and magnetic fields are (non zero components only)

$$B_{y}^{\pm} = B_{ya}^{\pm} + B_{yc}^{\pm}$$

$$B_{yc}^{\pm} = \mu_{0}j[L \mp (z - v_{z}t)]\gamma_{z}^{3}, \quad B_{ya}^{\pm} = j\mu_{0}\frac{v_{z}^{2}}{2c^{2}}\sigma N(z - v_{z}t)[2L \pm (z - v_{z}t)]\gamma_{z}^{5},$$

$$E_{x}^{\pm} = E_{xa}^{\pm} + E_{xc}^{\pm},$$

$$E_{xc}^{\pm} = v_{z}B_{yc}^{\pm}, \quad E_{xa}^{\pm} = \frac{c^{2}}{v_{z}}B_{ya}^{\pm}$$
(13)

The fields E_{xc}^{\pm} and B_{yc}^{\pm} are the usual fields produced by the moving current, while the fields E_{xa}^{\pm} and B_{ya}^{\pm} are the ones which appear due to the dissolving of the clot. The acceleration effect we consider here (acceleration of the clot particles) is due to the fields E_{xa}^{\pm} and B_{ya}^{\pm} .

We estimate the maximum of the energy the particles can gain due to the described above acceleration. Notice, that the potential (10) does not depend on the transverse coordinate x. Therefore there exists an integral of motion

$$p_x + eA_x = p_{0x} + eA_{0x} = \text{const.}$$
 (15)

where the index 0 labels the initial condition. As $j \rightarrow 0$, the potential A_x vanishes as well. This allows one to find the maximum of the transverse momentum

$$(p_x)_{\max} = p_{0x} + eA_{0x} \tag{16}$$

Let us consider now the longitudinal motion of the particles. The magnetic field B_{ya}^{\pm} produces the Lorentz force

$$F_z = ev_x B_{ya}^{\pm} \tag{17}$$

that acts on the electrons as well as on the protons in the positive z-direction in both regions $z - v_z t > 0$ and $z - v_z t < 0$. It should be noticed that the field B_{ya}^{\pm} is responsible for the acceleration of the clot as the whole system in z-direction. Some evidence of a such acceleration were mentioned in [8].

Notice that the longitudinal force which is acting on the electrons is greater than the one acting on the protons. This is because the transverse velocity v_x of the electrons is

greater than the one for the protons. Due to the quasineutrality of the clot, there appears a longitudinal electric field E_z that decelerates the electrons and accelerates the protons.

The longitudinal equations of motion for both electrons (e) and protons (p) are

$$\frac{dp_z^e}{dt} = -eE_z^{\pm} + ev_x^e B_{ya}^{\pm},$$

$$\frac{dp_z^p}{dt} = eE_z^{\pm} + ev_x^p B_{ya}^{\pm}$$
(18)

As a result, one obtains

$$\frac{d(p_z^e + p_z^p)}{dt} = e(v_x^e + v_x^p)B_{ya}^{\pm}$$
(19)

The longitudinal force in eq.(19) is $v_z(v_x^e + v_x^p)/2c^2$ times less than the force $2eE_z^{\pm}$ which is acting on the e^-p pair in the current direction. Therefore the acceleration in the *x*-direction is at least not less than the one in the *z*-direction. This allows one to estimate the maximum of the proton energy

$$\epsilon_{\max} \approx (p_x)_{\max} c = c(p_{x0} + eA_{x0}) \tag{20}$$

If the increase in the energy is much larger than the initial energy, then one finds

$$\epsilon_{\max} \approx ecA_{x0} \approx \frac{\mu_0}{2} ecj_0(z_0 - v_{z0})(2L_0 \mp (z_0 - v_{z0}t))\gamma_{z0}^3$$
(21)

where j_0 is the initial current density.

One can accept an estimation

$$(z_0 - v_{z0}t)(2L_0 \mp (z_0 - v_{z0}t))\gamma_{z0}^2 \approx L_0^{\prime 2}$$
(22)

This leads to an estimation for the maximum energy

$$\epsilon_{\max} \approx \frac{\mu_0}{2} e c j_0 L_0^{\prime 2} \gamma_{z0} \tag{23}$$

Thus we can write an estimation for the maximum gained energy

$$\epsilon_{\max} \approx 2j_0 L_0^{\prime 2} \gamma_{z \, 0} 10^{-7} (\text{Gev})$$
 (24)

The relatively new evidences [9] and models [10] show that the values

$$L'_0 \approx (0.01 - 0.1) \text{pc}$$

do not appear to be unrealistic. Accepting this estimation we can find the current density j_0 which is necessary for the protons to gain the maximum energy of the order of 10^{11} Gev. This gives

$$j_0 \approx 5(10^{-11} - 10^{-12}) \text{A/m}^2$$
 (25)

Estimate now the initial magnetic field which corresponds to the current density (25). Assuming that the relativistic factor $\gamma_{z\,0}$ is of the order of one we get from (13) the following relation

$$B_0 \approx \mu_0 j_0 L_0 \tag{26}$$

This yields

$$B_0 L_0 \approx \frac{\epsilon_{\max}}{ec} \tag{27}$$

For $\epsilon_{\text{max}} \approx 10^{11}$ Gev and $L_0 \approx 0.1$ pc the necessary value of the magnetic field is about 3Gs.

These estimations do not depend on the model. In order to prove this, we consider two different models

1) Thin long coil. The vector potential which describes the current at the clot particles location has the form

$$A_{\psi} = \frac{\mu_0}{2} jar \tag{28}$$

where r is the radius of the coil, and a is its thickness.

2) Thin round current loop. The vector potential describing the current at the clot particles location has the form

$$A_{\psi} = \frac{\mu_0}{2} j a^2 \ln(8R/a) \tag{29}$$

where a is the radius of current string, and R the radius of the loop.

It can be shown that in these models the result (23) for the maximum energy has a similar form if one makes the following substitutions

$$L_0^{'2} \rightarrow ar$$
 for the model (28)
 $L_0^{'2} \rightarrow a^2 \ln(8R/a)$ for the model (29)

Similar effects appear when the collective field inside the clot is produced by an effective electric charge in the clot instead of the current.

ACKNOWLEDGMENT

We would like to thank Boris V. Vayner, Lawrence Krauss and the members of particles/astrophysics seminar at CWRU for critical and useful comments. We also acknowledge the industrial problem solving group at CWRU for support of this work. We thank Hiro Fujita for careful reading of the manuscript and critical comments.

References

- [1] C.B. Stephenson and N. Sanduleak, Astrophys. J. Suppl. 33, 459, (1977)
- [2] T.J. Pearson, et al., Nature **290**, 365, (1981)
- [3] D.J. Bird, et al., Phys. Rev. Lett. **71**, 3401, (1993)
- [4] D.J. Bird, et al., Astrophys. J. **424**, 491, (1994)
- [5] G. Brooke, et al., Proc. of 19-th Int. Cosmic Ray Conf. (La Jolla), 2, 150, (1985)
- [6] N.N. Efimov, et al., in Astrophysical Aspects of the Most Energetic Cosmic Rays, ed. M. Nagano & F. Takahara (World Scientific, Singapore-1991)
- [7] A.M. Hillas, Ann. Rev. Astron. Astrophys. 22, 425, (1984)
- [8] W. Jaffe, H.C. Ford and Ferrarese, Nature, **364**, 213, (1993)
- [9] T. Zwiter et al., Fund. of Cosmic Phys. **13**, 310, (1989)
- [10] J.L. Gomez, et al., Astronomy and Astrophys. **292**, 33, (1994)