SIN P-TNP/96-07

TIFR/TH/96-22

April 1996

Matter induced ! m ixing

Abhee K.Dutt-Mazum der¹, Binayak Dutta-Roy

Saha Institute of Nuclear Physics, 1/AF Bidhan Nagar

Calcutta - 700 064, India

and

Anirban Kundu²

Tata Institute of Fundam ental Research, Hom i Bhabha Road,

M um bai - 400 005, India.

A bstract

A novel mechanism for ! mixing induced by neutron-proton asymmetry in nuclear matter is uncovered and the variation of the mixing angle with the extent of asymmetry is presented.

PACS numbers: 12.38 M h, 11.30 R d, 12.40 V v, 21.65.+ f

¹E-m ail: abhee@ tnp saha emet.in

²E-m ailakundu@ theory.tifr.res.in

The mixing of and ! m esons has attracted considerable attention for quite some time. The phenomenon of the needle (the narrow !) in the haystack (the broad resonance) was one of the amusing consequences of this idea. This has also been invoked in the context of the observed charge symmetry violation in mirror nuclei | commonly known as 0 kam ato-N olen-Schier anomaly [1, 2]. Several authors have investigated the problem and have attempted to explain the e ect by constructing the Charge Symmetry V iolating (CSV) potential, taking 0 ! m ixing into account [3]. Controversies in this eld have sparked o interest in alternate sources of isospin symmetry violation. In ref. [4], ! mixing has been studied within the fram ework of Vector M eson D om inance where the electrom agnetic m ixing is considered, while the same has been investigated by G ao et al [5] using an extended Nambu-Jona-Lasinio model. Several authors also take an isospin violating vertex factor as a source of the mixing [6] and others have im plicated the u-d m ass di erence in the quark loop as a contributor to the mixing [7].

However, the present treatment is somewhat dierent both in motivation and genesis from the earlier works, the main focus being the study of the mixing arising from the isospin symmetry breaking stemming from the nuclear medium through which the hadron propagates. Here signi cant mixing results, as we shall show, when the p \$ n symmetry of the nuclear matter is broken. The interaction Lagrangian involving the nucleons and the vector mesons may be written as

 $L_{int} = g [p p n n]_{0} + \frac{p}{2}g [p n^{+} + hc] + g[n n + p p]!$ (1)

which is isospin conserving. It may be noted that the coupling of the isovector 0 with the neutron and the proton has opposite sign in contrast to what

obtains for the isoscalar !. As a consequence, when we evaluate the selfenergy diagram s relevant for ⁰ ! m ixing, the neutron and proton loops contribute with opposite signs. As long as the ground state is symmetric under p \$ n interchange their contributions cancel exactly (as is the case for vacuum or for symmetric nuclear matter) as must be the case to the extent that isospin symmetry is respected not only by the Lagrangian but also by the medium. On the other hand if the ground state symmetry is broken i.e. number of neutron and proton in nuclear matter dier, then the corresponding Ferm im omenta, k_F^p and k_F^n , will be unequal whereby the cancellation does not take place and ⁰ ! mixing occurs.

The form of the nucleon propagator in nuclear matter may be expressed as

$$G(k) = G_F(k) + G_D(k)$$
 (2)

where

$$G_{\rm F}(k) = (k + M) \left[\frac{1}{k^2 - M^2 + i}\right]$$
 (3)

and

$$G_{D}(\mathbf{k}) = (\mathbf{k} + \mathbf{M}) \left[\frac{\mathbf{i}}{\mathbf{E}(\mathbf{k})} (\mathbf{k}_{0} - \mathbf{E}(\mathbf{k})) (\mathbf{k}_{F} - \mathbf{j}\mathbf{\tilde{k}}\mathbf{j}) \right]$$
(4)

with E $(\mathbf{\tilde{x}}) = \mathbf{\tilde{x}}\mathbf{j} + \mathbf{M}^2$, M being the elective mass of the nucleon in nuclear matter [8]. This density dependent shift of the nucleon mass (M ! M) in nuclear matter is estimated, for the present purpose, by solving the self-consistent equation within the framework of the W alecka M odel [8, 9]. The rst term in G (k), namely G_F (k), is the same as the free propagator of a spin $\frac{1}{2}$ ferm ion, except for the fact that the elective mass of the nucleon in nuclear matter is to be used, while the second part, G_D (k), involving (k_F $\mathbf{\tilde{x}}$), arising from Pauliblocking, describes them odi cations of the same in the nuclear matter at zero tem perature [8], as it deletes the on m ass-shellpropagation of the nucleon in nuclearm atterw ith m om enta below the Ferm i m om entum. Finite tem perature e ects m ay be incorporated by replacing the H eaviside -function by the Ferm i distribution, but the m ain features to be exposed in this letter are not m odi ed signi cantly. In general, the vector-boson induced polarization of the m edium, to one-loop, m ay be written as

$$lm (q^{2}) = \frac{i}{(2)^{4}}^{Z} d^{4}k \operatorname{Tr}[i^{1}iG (k + q)i^{m}iG (k)]$$
(5)

where represents the (!)NN vertex factor and l;m are the isospin indices. Just like the nucleon propagator, the polarization function can also be expressed as a sum of two parts,

$$(q^2) = F(q^2) + D(q^2)$$
 (6)

where the free (D irac) part ^F com es from the product $G^{F}(k + q)G^{F}(k)$, and the dense part ^D com es from $[G^{F}(k + q)G^{D}(k) + G^{D}(k + q)G^{F}(k)]$. The fourth combination, $G^{D}(k + q)G^{D}(k)$, does not contribute in the region of stability which concerns us here [10]. Form ulating the suitable D yson equations for the dressed propagators, one arrives [10] at the condition for nding their poles:

$$(q) = 0$$
 (7)

where the dielectric function (q) is the product of transverse and longitudinal dielectric functions:

$$(q) = {}_{L} (q) {}_{T}^{2} (q)$$
 (8)

and individual zeros for $_{\rm L}$ or $_{\rm T}$ gives the poles for the longitudinal or the transverse modes respectively. In our notation, taking $q = (q_0; j_1; 0; 0)$,

(;) = (0;1) pertains to the longitudinal and (;) = (2;3) to the transverse m ode.

For the unm ixed case

$$T_{(L)} = (1 - \frac{T_{(L)}}{q^2 - m_v^2})$$
(9)

where v = (;!) is the relevant vector m eson. In the static lim it (jqj=0), the elective m ass of the vector m eson is given by

$$q_0^2 = m_v^2 + m_v^2 = m_v^2 + L_v^L$$
: (10)

where ${}^{\rm L}_{\rm vv} = {}^{(v)}_{00} + {}^{(v)}_{11}$ and ${}^{\rm T}_{\rm vv} = {}^{(v)}_{22} = {}^{(v)}_{33}$.

W hen mixing takes place between and !, the polarization tensor becom es a 8 8 m atrix, with nonzero [!]. However, one can put it in a block diagonal form corresponding to longitudinal and transverse sectors. Because of the isotropy of nuclear matter, the transverse sector can further be split into a direct sum of two 2 2 m atrices, with $_{22} = _{33}$. We are interested only in the static limit; therefore, it will su ce to work with the simpler transverse part only, putting jqj = 0. In this case the dielectric function is given by the determ inant of

 $(\ _{!} = \ _{!})$, and the shifted m asses are found from the zeros of $_{T}$, i.e. from the following equation:

$$(q_0^2 m^2 m^2) (q_0^2 m_1^2 m_1^2) (T_1 m_1^2) = 0$$
 (12)

One notes that the solutions of eqn. (12) must be identical with those obtained for the longitudinal mode, since at the static limit there cannot be any preferred direction. We will omit the superscript T in our future

discussions, keeping in m ind that at the static lim it, $_{00} = 0$ and $_{11} = _{22}$. In eq. (12), the left-hand side is nothing but the determ inant of the m atrix $q_0^2 1$ M, where M is the one-loop m odi ed m ass m atrix

$$M = 0 + m^{2} + m^{2} + m^{2} + m^{2}$$

$$M = 0 + m^{2} + m^{2} + m^{2}$$

$$M = 0 + m^{2} + m^{2$$

The mixing angle () between 0 and ! elds can be obtained from

$$\tan 2 = \frac{2}{(m_1^2 - m_2^2) + (1)}$$
(14)

As has already been m entioned, the polarization tensor consists of two terms, one coming from the dense part describing the polarization of the Ferm i sea, while the other arises out of the nucleon-antinucleon excitation, i.e the e ect of the D irac Vacuum. The evaluation of D (q²) is straightforward as there is a nite upper limit involved in the loop integral [11]. However, the free part F (q²), needs to be regularized. We use the m ethod of dimensional regularization and employ the following subtraction scheme [12]

$$\mathbb{Q}^{n} \quad F \quad (q^2) = \mathbb{Q} \quad (q^2)^n \, \mathbf{j}_{M-1} \quad \mathbf{y}_{q^2 = m_v^2} = 0 \quad (n = 0; 1; 2:::; 1):$$
(15)

It may be noted, from (1) that whereas the !pp and !nn vertex factors com e with the same sign, the signs of the ⁰nn and ⁰pp vertex factors di er. This ensures the complete cancellation of !mixing amplitudes coming from the proton and the neutron bops for symmetric nuclearmatter (as well as for vacuum). However, [!] (q²) acquires a non-zero value whenever the ground state is not symmetric under p \$ n as has already been mentioned earlier. This comies solely from the bop integrals relevant for the densitydependent part, as the integrals have dimensional erent upper limits, viz, k_F^n and k_F^p , corresponding to neutron and proton loops, leading to a partial cancellation only.

W e present our results in Figs. 1 and 2. W e have taken M $_{n} = M_{p} = 938$ M eV, and norm alnuclear matter density ($_{0} = 0.17$ fm⁻³). The values of the coupling constants g and g are taken to be 2.71 and 10.1 respectively as estimated from Vector M eson D om inance (VMD) ts [13, 14].

In Fig.(1) we observe that in symmetric matter, mixing does not take place, while with increasing asymmetry the mixing becomes appreciable. We present here the results pertaining to the role of mixing on the masses at three times the normal nuclear matter density in order to make the elect stand out in clear relief.

It m ay be noted that the extent of m ixing, as expected, varies appreciably with asymmetry parameter (= $\frac{N_n N_p}{N_p + N_n}$). While the mass for 0 , taking m ixing into account, increases with , that of ! exhibits an opposite trend as is expected for the eigen values of a m assm atrix. A lso one should appreciate the fact that the mass splitting of ! in matter is larger by an order of magnitude compared to what is observed in vacuum. In addition an overall decrease of ! m eson m asses are observed. As far as the m ixing is concerned, the e ect of asymmetry is more pronounced for the case of $^{\rm 0}$ than for ! . No m ixing, however, takes place in the symmetric lim it for reasons explained in the text already as we are not including electrom agnetic or other such sources of mixing in order to focus attention on the mechanism at hand. In norm al nuclear density also such an e ect on masses due to m ixing is observed but on a lower scale. However, even at norm al nuclear matter density, we nd a signi cant value of the mixing angle () which, starting from zero, increases alm ost linearly with asymmetry parameter ()

as shown in Fig. 2.

The value of the mixing amplitude here is ten times larger, even at norm alnuclearm atter density, than what one estimates for electrom agnetic mixing [4]. In matter, even though the mass dierence of and ! meson (giving the energy denominator) increases by an order of magnitude the mixing amplitude is also quite large, therefore we do arrive at mixing angles comparable to those obtained from the more conventional sources.

The qualitative and dram atic manifestation of the mixing of and ! shows up in the two pion decay channel. The mixing results in principle to a narrow spike arising from the component in the mixed ! (which has a sm all width). In nuclear matter, how ever, this mode is not useful as the pions, being strongly interacting, would rescatter and lose the initial information. How ever, the dilepton mode or the decay channel + could possibly reveal the e ect of mixing.

To conclude and summarize, we stress that the mixing of ⁰ ! may also arise from the isospin asymmetric nuclear matter which is completely a matter induced e ect arising from the two dimensions (D (q²)) viz. the respective Ferm imomenta $k_{\rm F}^{\rm p}$ and $k_{\rm F}^{\rm n}$ in the proton and neutron loop. In the limit of complete asymmetry, and at densities available in the core of neutron stars as well as in relativistic heavy ion collision, this shift can be nearly 50 M ev for ⁰ and 25-M ev for !, from their unmixed values. It may be noted that even in the absence of mixing, the mass of ! and ⁰ gets shifted in the medium. The mixing amplitude is an order of magnitude larger than that for electrom agnetic mixing, even at normal nuclear matter density, which makes the phenomenon interesting and worth investigating.

8

References

- [1] K.Okam oto, Phys Lett 11, 150 (1964).
- [2] J.A.Nolen, Jr. and J.P. Schier, Ann. Rev Nucl. Sci19, 471 (1969).
- [3] L.A.Barreiro, A.P.Gateao, G.Krein, Phys Lett B 358, 7 (1995).
- [4] H.B.O'Connel, B.C. Pearce, A.W. Thom as and A.G.W illiam s, Phys Lett B 354, 14 (1995).
- [5] Shun-fu-Gao, C.M. Shakin and Wei-Dang Sun Phys. Rev. C 53, 1374 (1996)
- [6] S.Gardner, C.J.Horow itz, J.Piekarewicz, nucl-th/9508035
- [7] T.D.Cohen, G.A.M iller, Phys. Rev C 52, 3428 (1995)
- [8] B D. Serot and JD. W alecka, Adv. Nucl. Phys, (1986), vol. 16
- [9] J.D.W alecka, Ann. Phys. 83 491 (1974)
- [10] SA.Chin, Ann.Phys. 108, 301 (1977)
- [11] A K.Dutt-Mazum der, B.Dutta-Roy, A.Kundu and T.De, Phys. ReV C 53 790 (1996)
- [12] A.K.Dutt-Mazum der, A.Kundu, T.De, B.Dutta-Roy, Phys. Lett B, in press.
- [13] J.J. Sakurai, Currents and Mesons (Univ. of Chicago Press, Chicago, 1969).
- [14] H. Shiom i and T. Hatsuda, Phys. Lett. B 334 281 (1994)

Figure captions

- Fig. 1: Variation of vector m eson e ective m asses against asymmetry parameter () at three times norm all nuclear m atter density. Solid line and dotted line represent m asses with and without m ixing respectively.
- 2. Fig. 2: Variation of mixing angle ($^{0})$ against asymmetry parameter at norm alnuclearm atter density.



