

hep-ph/mm nnyyy
 IISc-C T S-9/96
 LNF-96/018 (P)

Eikonalized mini-jet cross-sections in collisions¹

A .Corsetti

INFN , Physics Department, University of Rome La Sapienza, Rome, Italy

R .M .Godbole²

Center for Theoretical Studies, Indian Institute of Sciences

Bangalore 560 012, India

and

G .Pancheri

INFN -Laboratori Nazionali di Frascati, I00044 Frascati

Abstract

In this note we assess the validity and uncertainties in the predictions of the eikonalized mini-jet model for $\gamma\gamma$. We are able to find a choice of parameters where the predictions are compatible with the current data. Even for this restricted range of parameters the predictions at the high \sqrt{s} energies, which can be reached at the TeV energy e^+e^- colliders, differ by about 25%. LEP 2 data can help pinpoint these parameters and hence reduce the uncertainties in the predictions.

¹ To appear in the proceedings of the workshop on e^+e^- 2000 GeV Linear Colliders Annecy, G rand Sasso, Hamburg (1995).

² On leave of absence from Dept. of Physics, Uni. of Bombay, Bombay, India.

Eikonalized mini-jet cross-sections

A. Corsetti¹, R.M. Godbole², G. Pancheri³

¹ INFN, Univ. La Sapienza, Roma, Italy

² CTS, IISc, Bangalore, India

³ INFN, Frascati, Italy

In this note we wish to assess the validity and uncertainties of the eikonalized mini-jet model in predicting σ_{ab}^{inel} and further to ascertain whether measurements at LEP-200 and HERA can constrain various parameters of the model. In its simplest formulation, the eikonalized mini-jet cross-section is given by

$$\sigma_{ab}^{inel} = \sigma_{ab}^{had} \int_0^Z d^2b [1 - e^{-n(b;s)}] \quad (1)$$

where the average number of collisions at a given impact parameter b is obtained from

$$n(b;s) = A_{ab}(b) \left(\sigma_{ab}^{soft} + \frac{1}{\sigma_{ab}^{had}} \sigma_{ab}^{jet} \right) \quad (2)$$

with $A_{ab}(b)$ the normalized transverse overlap of the partons in the two projectiles and σ_{ab}^{had} to give the probability that both colliding particles a, b be in a hadronic state. σ_{ab}^{soft} is the non-perturbative part of the cross-section from which the factor of σ_{ab}^{had} has already been factored out and σ_{ab}^{jet} is the hard part of the cross-section. The rise in σ_{ab}^{jet} drives the rise of σ_{ab}^{inel} with energy [1]. We have also assumed the factorization property

$$\sigma_p^{had} = \sigma^{had}; \quad \sigma^{had} = (\sigma_p^{had})^2:$$

The predictions of the eikonalized mini-jet model [2] for photon induced processes [3] depend on 1) the assumption of one or more eikonals, 2) the hard jet cross-section $\sigma_{ab}^{jet} = \int_{p_{tmin}}^{\infty} \frac{d^2p_t}{dp_t^2} dp_t^2$ which in turn depends on the minimum p_t above which one can expect perturbative QCD to hold, viz. p_{tmin} , and the parton densities in the colliding particles a and b , 3) the soft cross-section σ_{ab}^{soft} , 4) the overlap function $A_{ab}(b)$, defined as

$$A_{ab}(b) = \frac{1}{(2\pi)^2} \int_0^Z d^2q F_a(q) F_b(q) e^{iq \cdot b} \quad (3)$$

where F is the Fourier transform of the b -distribution of partons in the colliding particles and 5) last but not the least σ_{ab}^{had} .

In this note we shall restrict ourselves to a single eikonal. The hard jet cross-sections have been evaluated in LO perturbative QCD. The dependence of σ_{ab}^{jet} on p_{tmin} is strongly correlated with the parton densities used. Here we show the results using GRV densities [4] (see ref. [5] for the results using the DG densities [6]). For the purposes of this note, we determine σ_{ab}^{soft} from σ_p^{soft} which is obtained by a fit to the photoproduction data. We use the Quark Parton Model suggestion $\sigma_{ab}^{soft} = \frac{2}{3} \sigma_p^{soft}$.

In the original use of the eikonal model, the overlap function $A_{ab}(b)$ of eq. (3) is obtained using for F the electromagnetic form factors and thus, for photons, a number of authors [7, 8] have assumed for F the pole expression used for the pion electromagnetic form factor, on the basis of Vector Meson Dominance (VMD). We shall investigate here another possibility, i.e. that the b -space distribution of partons in the photon is the Fourier transform of their intrinsic transverse momentum distributions. This will correspond to use the functional expression expected for the perturbative part [9]

$$\frac{dN}{dk_t^2} = \frac{1}{k_t^2 + k_0^2} \quad (4)$$

Recently this expression was confirmed by the ZEUS [10] Collaboration, with $k_0 = 0.66 \pm 0.22$ GeV. For e^+e^- collisions, the overlap function is now simply given by

$$A(b) = \frac{1}{4} k_0^3 K_1(k_0 b) \quad (5)$$

with K_1 the Bessel function of the third kind. It is interesting to notice that for photon-photon collisions the overlap function will have the same analytic expression for both our ansatz: the VMD inspired pion form factor or the intrinsic transverse momentum; the only difference being that the former corresponds to a fixed value of $k_0 = 0.735$ GeV whereas the latter allows us to vary the value of the parameter k_0 . Thus both possibilities can be easily studied by simply changing k_0 appropriately. Notice that the region most important to this calculation is for large values of the parameter b , where the overlap function changes trend, and is larger for smaller k_0 values.

As for P^{had} , this is clearly expected to be $O(\alpha_{\text{em}})$ and from VMD one would expect $1=250$. From phenomenological considerations [8] and fits to HERA data, one finds a value $1=200$, which indicates at these energies a non-VMD component of $\sim 20\%$. It should be noticed that the eikonalised minijet cross sections do not depend on A and P^{had} separately, but depend only on the ratio of the two [11, 12].

Having thus established the range of variability of the quantities involved in the calculation of total photonic cross sections, we now proceed to calculate and compare with existing data the eikonalized minijet cross-section for e^+e^- collisions. We use GRV (LO) densities and values of p_{min} deduced from a best fit to photoproduction. As discussed in [15], it is possible to include the high energy points in photoproduction using GRV densities and $p_{\text{min}} = 2$ GeV, but the low energy region would be better described by a smaller p_{min} . This is the region where the rise, according to some authors, notably within the framework of the Dual Parton Model, is attributed to the so-called soft Pomeron. For our studies here we use $p_{\text{min}} = 2.0$ GeV. We also use $P^{\text{had}} = 1=204$ and $A(b)$ from eq.(5) with different values of k_0 . One choice for k_0 is the pole parameter value in the photon b -distribution expression, which includes both the intrinsic transverse momentum option 0.66 ± 0.22 GeV as well as the pion form factor value, 0.735 GeV. The other value, 1 GeV, is a possible choice which appears to fit the present data better than everything else. Our predictions are shown in Fig.(1). A comparison with existing data shows that all of

Figure 1: Total inelastic photon-photon cross-section for $p_{\text{min}} = 2: \text{GeV}$ and different parton b-distribution in the photon. The solid line corresponds to $k_0 = 1: \text{GeV}$.

our choices are compatible with the data within the present experimental errors. At high energies, however, like the ones reachable with the proposed linear photon colliders, these predictions vary by about 25%. Reducing the error in the LEP 1 region and adding new data points in the cm. region attainable at LEP 2, can help pinpoint and restrict the choices. Were the LEP 1 and LEP 2 data to confirm the present values, we believe that the best representation of the present data is obtained with the higher k_0 value.

Acknowledgements

R. M. G. wishes to acknowledge support from C.S.I.R. (India) under grant no. 03(0745)/94/EMR-II. This research is supported in part by the EEC program "Human Capital and Mobility", contract CT 92-0026 (DG 12 COMA).

References

- [1] D.Cline et al., Phys. Rev. Lett. 31 (1973) 491, T.Gaisser and F.Halzen, Phys. Rev. Lett. 54 (1985) 1754, G.Pancheri and Y.N.Srivastava, Phys. Lett. B 182 (1985).
- [2] A.Capella and J.Tran Thanh Van, Z.Phys.C 23 (1984)168, P.L'Heureux, B.Margolis and P.Valin, Phys. Rev. D 32 (1985) 1681, L.Durand and H.Pi, Phys. Rev. Lett. 58 (1987) 58.
- [3] J.C.Collins and G.A.Ladinsky, Phys. Rev. D 43 (1991) 2847.
- [4] M.Gluck, E.Reya and A.Vogt, Phys. Rev. D 46 (1992) 1973.
- [5] M.Drees and R.M.Godbole, Z.Phys.C 59 (1993) 591.
- [6] M.Drees and K.Grassie, Z.Phys.C 28 (1985) 451.
- [7] K.Honjo et al., Phys. Rev. D 48 (1993) 1048.
- [8] R.S.Fletcher, T.K.Gaisser and F.Halzen, Phys. Rev. D 45 (1992) 377; erratum Phys. Rev. D 45 (1992) 3279.
- [9] J.Field, E.Pietarinen and K.Kajantie, Nucl. Phys. B 171 (1980) 377; M.Drees, In the proceedings of the 23rd International Symposium on Multiparticle Dynamics, Aspen, Colo., Sep. 1993, Eds. M.M.Black and A.R.White.
- [10] M.Derrick et al., ZEUS Collaboration, Phys. Lett. B 354 (1995) 163.
- [11] M.Drees, Univ. Wisconsin report MADPH-95-867, Proceedings of the 4th workshop on TRISTAN physics at High Luminosities, KEK, Tsukuba, Japan, Nov. 1994.
- [12] M.Drees and R.M.Godbole, Journal Phys. G., G 21 (1995) 1559.
- [13] H.Abramowicz, K.Charchula and A.Levy, Phys. Lett. B 269 (1991) 458.
- [14] K.Hagiwara, M.Tanaka, I.Watanabe and T.Izubuchi, Phys. Rev. D 51 (1995) 3197.
- [15] C.Corsetti, R.M.Godbole and G.Pancheri, in preparation.