

May, 1996

YCTP-P 9-96

Chiral Perturbation Theory, Large- N_c and the π^0 Mass

Nick Evans^{*}, Stephen D. H. Hsu^y and Mykola Schwetz^z

Slane Physics Laboratory, Yale University, New Haven, CT 06511

Abstract

Using chiral perturbation theory and the large- N_c expansion, we obtain expressions for the π^0 mass and π^0 mixing in terms of low-energy chiral Lagrangian parameters. This is accomplished through an intermediate step of 'matching' the topological susceptibility in the large- N_c and chiral Lagrangian descriptions. By inserting the values of well-measured parameters we obtain predictions involving the second order parameters L_6, L_7 and L_8 . The prediction for L_6 is quite restrictive even after allowing for $1/N_c$ corrections.

^{*}nick@zen.physics.yale.edu

^yhsu@hsunext.physics.yale.edu

^zm.s@genesis2.physics.yale.edu

1 Introduction

Chiral perturbation theory (CHPT) provides a useful description of low-energy QCD in terms of the relevant light degrees of freedom [1]. These light degrees of freedom are simply the $N_f^2 - 1$ pseudo-Goldstone bosons (PGB's) which result from the breaking of chiral symmetry:

$$SU(N_f)_L \times SU(N_f)_R \rightarrow SU(N_f)_V :$$

We do not include the $U(1)_A$ symmetry in the above as it is violated by the axial anomaly [2]. The axial anomaly has physical effects due to the presence in QCD of field configurations with non-trivial winding number. Were this not the case the chiral Lagrangian would have to be modified to accommodate an additional (ninth) pseudo-Goldstone boson, which has the quantum numbers of the η' . It is well-known that this is what occurs in the large N_c limit { the physical effects of the anomaly are suppressed [3] *. In the large- N_c limit the low-energy effective Lagrangian contains the η' as well as the pions [5].

It might be regarded as a failure of the large- N_c expansion that the η' is so heavy { $m_{\eta'} = 958 \text{ MeV}$ { relative to the other strange PGB's such as the K^* ; K^0 ; K^0 and η . However, we will argue here that this is a consequence of the smallness of the light quark masses and need not imply a breakdown in large- N_c thinking. The point is that if one formally takes the limit $N_c \rightarrow \infty$ with $m_{u,d,s}$ fixed, it is easy to see that the η' becomes roughly degenerate with the strange PGB's (see section 2). However, it is more appropriate for the real-world to consider a double limit in which $N_c \rightarrow \infty$ and $m_{u,d,s} \rightarrow 0$ simultaneously, in some fixed ratio. Indeed, the breaking of the chiral $SU(3) \times SU(3)$ symmetry associated with the strange quark mass $m_s = 4$ to 15% is still smaller than $1/N_c$ ($N_c = 3$) $\sim 30\%$. As we will discuss below, after taking this into account it becomes clear that the η' is heavy relative to the PGB's because of the smallness of m_s rather than a failure of large- N_c . Using a simultaneous scaling of N_c and m_q we can contemplate a large- N_c limit with the following hierarchy of masses:

$$m_{\text{baryon}} \gg m_{\eta'} \gg m_{K^*}; \quad (1)$$

We note that a similar point of view was taken by Leutwyler in [6], where a bound on the light quark mass ratios was derived using a double expansion in large- N_c and the quark masses.

We will take the viewpoint that large- N_c gives a qualitatively correct picture of QCD dynamics, and exploit the tension between the standard CHPT effective Lagrangian and the one resulting from large- N_c . In particular, we will utilize the special status of the η' in

*Despite this, the η' -angle may still have physical consequences in the large- N_c limit [3]. Also see [4] for a discussion of the possible large- N_c behavior of instantons.

determining a quantity known as the topological susceptibility:

$$\frac{\langle \chi^2 \rangle}{V} = \frac{1}{V Z} \left. \frac{\partial^2 Z}{\partial \theta^2} \right|_{\theta=0}; \quad (2)$$

where Z is the QCD partition function, V the volume of spacetime and θ is the winding number given by

$$\theta = \frac{1}{32\pi^2} \int_V d^4x \text{Tr} G \tilde{G}; \quad (3)$$

In the large volume limit, Z can be expressed in terms of the vacuum energy density ϵ_0 :

$$Z = \exp(-V \epsilon_0); \quad (4)$$

The problem is thus reduced to that of computing the ϵ_0 dependence of ϵ_0 . This can be done either in CHPT, which is equivalent to an expansion in the light quark masses m_q , or in the large- N_c effective Lagrangian, which involves a double expansion in $1/N_c$ and m_q . Equating the two results for the susceptibility will yield a relationship at leading order in N_c between m_0 and low-energy parameters appearing in CHPT. An alternative way of viewing our calculation is in terms of two different effective Lagrangians valid at the two energy scales $\mu_1 = m_0$ and $\mu_2 = m_K$. At the lower scale the η^0 has been ‘integrated out’. Requiring that both Lagrangians (one with the η^0 , the other without) yield the same susceptibility implies a non-trivial relation between their parameters.

In the following sections we will compute the topological susceptibility in both CHPT and Large- N_c CHPT to obtain the desired relation. In the final section we will present our results and some final comments.

2 Large- N_c Effective Lagrangian

As previously mentioned, at large- N_c the effective Lagrangian must be modified to include an additional light meson, the η^0 , which plays the role of an additional PGB associated with $U(1)_A$ [3, 5]. It is convenient to describe all N_f^2 PGBs in terms of a $U(N_f)$ matrix field $U(x)$. The η^0 is related to the phase of the determinant of $U(x)$:

$$\det U(x) = e^{i\theta(x)} \quad (5)$$

with

$$\theta(x) = \frac{q}{F} \frac{\text{Tr} U(x)}{N_f} \quad (6)$$

The effective Lagrangian at large N_c is

$$L_N = \frac{F^2}{4} \text{Tr} (\partial_\mu U^\dagger \partial^\mu U) + R \text{Tr} (M U^\dagger) - \frac{1}{2} (\partial_\mu \theta)^2; \quad (7)$$

where the pion decay constant $F = O(N_c^{1/2})$, the chiral symmetry breaking scale $\Lambda = O(N_c)$ (related to the quark condensate in the chiral limit) and $\theta = O(1)$ (related to the topological susceptibility of pure gluodynamics [3]). Note that the θ angle has been absorbed from the mass matrix M into a shift of ϕ . L_N represents the leading order terms in the effective Lagrangian in the formal counting scheme where one takes the quark masses m_q in the mass matrix M to be $O(N_c^{-1})$ and derivatives $\partial = O(N_c^{1/2})$ m .

Taking the quark mass matrix to be (we now restrict ourselves to $N_f = 3$ and the SU(2) isospin limit $m_u = m_d = m$)

$$M = \begin{pmatrix} 0 & 1 \\ m & 0 & 0 \\ 0 & m & 0 \\ 0 & 0 & m_s \end{pmatrix} \quad (8)$$

we obtain the following mass relations:

$$\begin{aligned} m^2 &= \frac{2m}{F^2} \\ m^2 &= \frac{2(m + 2m_s)}{3F^2} \\ m_{\pi^0}^2 &= \frac{2(2m + m_s)}{3F^2} + \frac{6}{F^2} \\ m_{\pi^0}^2 &= \frac{2 \frac{p}{2} (m - m_s)}{3F^2} : \end{aligned} \quad (9)$$

The first two equations are identical to the results of CHPT, and the last equation describes η mixing. Note that both the anomalous and CHPT contributions to $m_{\pi^0}^2$ are formally $O(1/N_c)$ in N_c counting. As mentioned previously it is possible to take the quark masses $m_q = O(N_c^{-1}) \rightarrow 0$ as $N_c \rightarrow \infty$ such that the anomalous term continues to dominate (i.e., $\gg m_q$). This is the limit that we will consider. In real QCD where the $N_c \rightarrow \infty$ limit is only approximate the anomalous contribution to m_{π^0} is presumably not suppressed below the CHPT contribution which is small because of the quark masses. We argue therefore that the large N_c form of (7) is appropriate to QCD.

The parameter θ can be extracted at leading order by requiring that the eigenvalues of the η mass matrix correspond to the phenomenologically observed values of 547 and 958 MeV respectively. If we allow CHPT and $1/N_c$ uncertainty of 50% in the mass relations (9) (from, e.g., uncertainties in F , and subleading interactions not in (7) which shift the mass relations (9) etc.), we obtain the result

$$(137 \text{ MeV})^4 < \theta < (202 \text{ MeV})^4 : \quad (10)$$

In our calculations, we have used $F = 93 \text{ MeV}$, the allowed range of light quark masses $m_u = m_d = 7 \text{ MeV}$ and measured pion and kaon masses to conclude that $m_s = 150 \text{ MeV}$ and $\theta = O(200 \text{ MeV})^3$.

Note that the above results are consistent with the claim that m_s mainly due to the smallness of m_s .

The next step is to compute the topological susceptibility using L_N . This requires minimizing the potential energy at small angle and differentiating twice with respect to i . The $SU(N_f)$ isospin limit ($m = m_s$) of this problem was analyzed previously by Leutwyler and Smilga [7]. In this case the global minimum of the potential occurs when U is a multiple of the unit matrix $U = e^{i\pi/N_f}$. The vacuum energy is of the form

$$E_0 = -N_f m + \frac{1}{2} \frac{m^2}{N_f + m} + O(m^4); \quad (11)$$

and the topological susceptibility at $\theta = 0$ is

$$\frac{\langle \chi^2 \rangle}{V} = \frac{m}{N_f + m}; \quad (12)$$

In the limit we are considering, the denominator is dominated by m , so

$$\frac{\langle \chi^2 \rangle}{V} \approx \frac{m}{N_f} \left(1 - \frac{m}{N_f}\right); \quad (13)$$

The second term is $O(m^2)$, and one may wonder whether it is consistent to retain it given the neglect of terms of higher order in (M/U) in L_N . The point is that due to an additional power of θ the term in (13) is larger by a factor of N_c than the subleading corrections we have neglected. When we match topological susceptibilities with the true low-energy theory described by the $SU(N_f)$ chiral Lagrangian (no θ), this term will fix a combination of $L_{6;7;8}$ at leading order in N_c .

It is important to note that the η' and the PGB's play a unique role in determining $\frac{\langle \chi^2 \rangle}{V}$. The other particles not described by $U(x)$, such as the meson, or the baryons p, n, \dots are expected to have zero expectation value in the vacuum, and hence do not contribute to the susceptibility even if their interactions have some θ -dependence. This observation is independent of large- N_c , which only guarantees that (12) is accurate to leading order.

The extension of (12) to the case of a heavy strange quark is straightforward, and we obtain

$$\frac{\langle \chi^2 \rangle}{V} = \frac{M}{1 + M}; \quad (14)$$

where $M = (1/m_u + 1/m_d + 1/m_s)^{-1} = m m_s / (m + 2m_s)$. Note that this reduces to m/N_f where $N_f = 3; 2$ for $m_s \rightarrow m$ and $m_s \rightarrow 1$ respectively. M is approximately m , which is determined by the pion mass relation in (9).

The range in θ given in (10) translates to the following range in the topological susceptibility

$$3.7 \cdot 10^7 < \frac{\langle \chi^2 \rangle}{V} < 4.05 \cdot 10^7 \quad (15)$$

The uncertainty in χ does not lead to a large variation in $\langle h^2 \rangle = V$. There are, however, additional $1/N_c$ corrections to the relation (14) which we have not yet examined in detail. As mentioned previously, the uncertainties represented in (15) are due to $1/N_c$ corrections to the mass relations (9). However, subleading interactions which do not appear in (7) may also alter the calculation of the susceptibility.

There are two types of subleading corrections, both formally of order $1/N_c$, which we must consider.

Higher orders in (M/U) which induce corrections suppressed by $O(m_q)$. The most important of these are of the form of the $L_{6,7,8}$ interactions in (19) below. In large- N_c counting $L_{6,7}$ are suppressed by Zweig's rule so it is L_8 which is most important[†].

$O(1/N_c)$ interactions of the ϕ in L_N which violate the $U(N_f) \times U(N_f)$ flavor symmetries but which preserve $SU(N_f)$ symmetries. These corrections represent the fact that the ϕ is only a PGB in the large- N_c limit. The most important of these is of the form

$$(\bar{\psi} \psi)^2 \text{Re tr}(M e^{i \frac{1}{N_f} U^Y}) ; \quad (16)$$

where the coefficient is $O(1/N_c)$.

We can estimate the size of the uncertainties introduced in the large- N_c susceptibility by corrections of the above type. Naively, the $L_{6,7,8}$ interactions will lead to a correction of order $O(m_s)$ which would be of order 15%. However, the issue is somewhat more complicated than this. The structure of L_8 is such that its effect on the calculation of the susceptibility is equivalent to a shift in the quark masses in (14): $m_i \rightarrow m_i(1 + O(m_i))$. Hence only m_s receives a non-negligible correction. However, the strange quark mass has only a very small effect on the susceptibility because it is so much more massive than the u, d quarks. The parameter $M = m(1 + O(m = m_s))$ in (14) depends only weakly on m_s . The $L_{6,7}$ operators, although suppressed relative to L_8 by a power of $1/N_c$, have a larger effect on the susceptibility. An explicit calculation shows that $L_{6,7}$ shift the susceptibility by $O(m_s 1/N_c)$, leading to a larger uncertainty than is given by the range (15). This is the dominant subleading correction. We will associate an additional error of 5% to this correction, leading to the following range of susceptibility (representing roughly 10% error):

$$3.5 \cdot 10^7 < \frac{\langle h^2 \rangle}{V} < 4.3 \cdot 10^7 ; \quad (17)$$

[†] In the real world the central values of $L_{6,7}$ are roughly $(1/N_c)L_8$, which is at least consistent with the $1/N_c$ expansion approximation.

The second type of correction leads to a only a small uncertainty in the susceptibility. Let ϕ_0 be the value of ϕ which minimizes the vacuum energy ϕ_0 in (11) at non-zero θ :

$$\phi_0 = \frac{1}{(1 + M^2)^{1/2}} : \quad (18)$$

We can then estimate the shift in the susceptibility by substituting ϕ_0 into (16) and differentiating twice with respect to θ . This leads to a correction which is suppressed by a factor of $O(N_c^{-1} M^{-2})$, making it less than 5% of (14).

3 Chiral Lagrangian

We next compute the topological susceptibility using the standard $SU(N_f)$ chiral Lagrangian appropriate below the ϕ_0 mass. The heavy ϕ_0 has been 'integrated-out' and its effects are exhibited only in the chiral coefficients. The susceptibility only depends on the potential of the chiral Lagrangian. Terms with derivatives are irrelevant.

At $O(m^2)$ the potential is [1]

$$V = -\text{Re tr}(M U^\dagger) - L_6 \frac{16}{F^4} (\text{Re tr} M U^\dagger)^2 - L_7 \frac{16}{F^4} (\text{Im tr} M U^\dagger)^2 - L_8 \frac{8}{F^4} \text{Re tr}(M U^\dagger M U^\dagger) ; \quad (19)$$

where the mass matrix M now incorporates the θ -angle, $M = M \exp i \theta / N_f$. Here the $U(x)$ matrix field is restricted to unit determinant and is parameterized as

$$U(x) = \exp(i 2^{-1} \pi(x) T^a / F) : \quad (20)$$

The topological susceptibility is again found by minimizing the potential for U and taking the second derivative of Z with respect to (i) at the minimum. We find

$$\frac{\chi^2}{V} = \frac{2}{9} \left[\frac{(2m + m_s)}{2} + \frac{16}{F^4} ((L_6 + L_7)(2m + m_s)^2 + L_8(2m^2 + m_s^2)) \right. \\ \left. - \frac{4(m_s - m)^2 \left(\frac{1}{2} + \frac{16}{F^4} ((L_6 + L_7)(2m + m_s) + L_8(m + m_s)) \right)^2}{(m + 2m_s) + \frac{16}{F^4} (2L_6(2m + m_s)(m + 2m_s) + 4L_7(m_s - m)^2 + 2L_8(m^2 + 2m_s^2))} \right] ; \quad (21)$$

By equating this result, which is a function of the second order parameters of the chiral Lagrangian (19), with the range (15) determined from the large N_c expansion we can impose constraints on the values of the parameters L_6 ; L_7 and L_8 in the chiral Lagrangian (19).

The expression (21) has some interesting properties. The leading order behavior is

$$\frac{\chi^2}{V} = M^2 + \dots ; \quad (22)$$

which agrees with the leading order large- N_c result (14) for $\mu \gg M$. Expanding (21) in powers of the light quark masses, we can see that its leading dependence on L_6 is $L_6(m)(m_s=F^4)$, whereas the leading dependence in $L_{7,8}$ is $L_{7,8}(m)(m=F^4)$. This implies that the susceptibility is much more sensitive to changes in L_6 than $L_{7,8}$. When we perform the matching of susceptibilities below, we will find that L_6 is more restricted than $L_{7,8}$.

We can derive further constraints on the $L_{6;7,8}$ by matching higher derivatives of the vacuum energy with respect to i in the large- N_c and chiral descriptions. This corresponds to matching $h^4 i$, $h^6 i$, etc. Unfortunately, we have checked that in the chiral Lagrangian description $h^4 i$ is still only weakly dependent on $L_{7,8}$. The leading dependence on L_6 is again larger by a factor of $O(m_s=m)$ than the dependence on $L_{7,8}$. Thus we do not believe that additional constraints on $L_{7,8}$ will be obtained by further matching.

4 Matching

As to the parameters $L_6; L_7$ and L_8 from low energy data yields the following one-sign a range of values [1]:

$$\begin{aligned} L_6 &= \quad \quad .0002 \quad .0003 \\ L_7 &= \quad \quad .0004 \quad .0002 \\ L_8 &= \quad + .0009 \quad .0003 \quad : \end{aligned} \tag{23}$$

Upon matching the results for the topological susceptibility in the large N_c (14) and standard chiral lagrangian (21) we obtain a more restrictive prediction for the volume in $L_6; L_7$ and L_8 space consistent with the observed meson masses upto the errors described above in (17). The combined constraints are shown in Fig. 1. The parameter L_6 is well determined by the fit. L_7 and L_8 are not so well constrained. In Fig 2 and Fig 3 we project our limits onto the $L_6 - L_8$ and $L_6 - L_7$ planes. Figure 2 shows the allowed regions of $L_6 - L_8$ with L_7 unconstrained and Figure 3 shows the allowed regions of $L_6 - L_7$ with L_8 unconstrained.




Fig.1: The region of $L_{6,7,8}$ space consistent with the observed meson masses.




Fig.2: Projection of allowed region of parameter space onto L_6 - L_8 plane.




Fig.3: Projection of allowed region of parameter space onto L_6 - L_7 plane.

5 Discussion

The η' has certain special properties in QCD as it is 'almost' the Goldstone boson associated with spontaneously broken $U(1)_A$ symmetry. Within the large- N_c expansion this 'almost' can be made quantitative and relations derived (see (14)) between the η' and η masses and the topological susceptibility. The latter quantity measures the sensitivity of the QCD vacuum energy to small changes in θ at $\theta = 0$. We can also compute the susceptibility directly in chiral perturbation theory. The leading order result in CHPT is precisely that obtained from the large- N_c Lagrangian if the η' is taken to be very massive ($m_{\eta'} \gg M_\pi$). However, the subleading corrections in CHPT come from $L_{6,7,8}$ whereas in the large- N_c expansion they depend on the ratio $M_\pi = f_\pi$. We have exploited this observation to derive what are essentially large- N_c predictions for the second order CHPT coefficients (particularly L_6) in terms of the η' mass which is taken as an input. Our prediction for L_6 is restrictive even when $1/N_c$ corrections are included. The central value we predict is within the 1σ region previously extracted from low-energy data (also using large- N_c) [1].

A simple way of understanding our calculation is in terms of effective Lagrangians valid at two scales, $\mu_1 \sim m_\rho$ and $\mu_2 \sim m_K$; . At the lower scale the η' has been 'integrated out'. The effect of virtual η' s appears in the coefficients $L_{6,7,8}$ and is determined at leading order in large- N_c . In this sense our calculation is similar to that of [8], which estimated the parameters L_i from resonance exchange.

Acknowledgements

This work was supported under DOE contract DE-AC02-ERU3075.

References

- [1] J. Gasser and H. Leutwyler, *Ann. Phys.* 158, 142 (1984); For a recent review, see J. Bijnens, G. Ecker and J. Gasser, hep-ph/9411232. To be published in 2nd Edition of DAPHNE Physics Handbook.
- [2] For a review, see, e.g., *Aspects of Symmetry*, S. Coleman, Cambridge, 1985.
- [3] E. Witten, *Nucl. Phys. B* 156, 269 (1979).
- [4] E. V. Shuryak, hep-ph/9503467
- [5] E. Witten, *Ann. Phys.* 128, 363 (1980).
P. Di Vecchia and G. Veneziano, *Nucl. Phys. B* 171, 253 (1980).
P. Di Vecchia, F. Nicodemi, R. Pettorino and G. Veneziano, *Nucl. Phys. B* 181, 318 (1981).
- [6] H. Leutwyler, hep-ph/9601234.
- [7] H. Leutwyler and A. Smilga, *Phys. Rev. D* 46, 5607 (1992).
- [8] G. Ecker, J. Gasser, A. Pich and E. de Rafael, *Nucl. Phys. B* 321, 311 (1989).