

Perturbative QCD approach and particle energy spectra¹

SERG IO LUP IA

Max-Planck-Institut für Physik
(Werner-Heisenberg-Institut)
Fohringer Ring 6, D -80805 München, Germany
E-mail: lupia@mppmu.mpg.de

Abstract

The Modified Leading Log Approximation of QCD and the local parton hadron duality picture are shown to give a fully satisfactory description of data on charged particle energy spectra in e^+e^- annihilation. The analysis of data in terms of moments points out the running of α_s both at very low cms energy ($O(3 \text{ GeV})$) and at very low particle energy ($O(200\text{--}500 \text{ MeV})$). The study of the universal soft limit of energy spectra at low particle energy is proposed as a test of QCD coherence. The extension of this approach at identified particles' spectra is discussed.

¹to be published in the Proceedings of the XXXIst Rencontre de Moriond, "QCD and High Energy Interactions", Les Arcs, March 23rd–30th, 1996, Ed. J. Tran Thanh Vân, Editions Frontieres, Paris-CEDEX

The Modified Leading Log Approximation (MLLA)¹⁾ plus the Local Parton Hadron Duality (LPHD)²⁾ as hadronization prescription give a quite successful description of production process of soft and semihard particles (for a recent discussion see e.g.³⁾). Predictions on the single particle inclusive momentum spectrum, the well-known "hump-backed plateau", is one of the best successes of this approach. However, the study of the spectrum alone does not allow to see to what extent experimental data reveals peculiar features of the theory, like for instance coherence and the running of the coupling. In this short note, new alternative analyses more sensitive than the spectrum itself to the different aspects of the theory are discussed. Due to the limited amount of space, I just give the flavour of our studies and I refer to the original literature⁴⁾⁽⁶⁾ for more detailed presentations.

MLLA predictions⁷⁾ for the moments of single particle inclusive energy spectrum have been compared in⁴⁾ with moments extracted from experimental spectra. Good agreement between experimental data and theoretical predictions of MLLA with running coupling has been found in a wide cms energy range from 3 up to LEP energy. The model with fixed coupling can reproduce the multiplicity and the slope of the first moment only. The two models are particularly different at low cms energies, where the effect of the running coupling is large. Notice that the model with fixed coupling cannot describe the data even if one relaxes the normalization at thresholds.⁶⁾ The sensitivity of moments to the running of the QCD coupling has then been established; the validity of the description at low cms energy is particularly interesting. In this respect, it is important to point out that further tests of this picture could be performed at HERA (in the Breit frame), where many data points at different Q^2 could be obtained within the same experiments. Going to larger cms energies, the extrapolation of theoretical predictions have been shown to be successful at LEP-1.5 cms energies;⁵⁾ the same analysis could be repeated at the forthcoming LEP-2.

Let us now look at the invariant cross section $E \frac{dn}{d^3p}$ at particle energy E of few hundreds MeV. QCD coherence predicts that very soft particles do not multiply;¹⁾ since they have a long wavelength, they should see the total initial charge only. Particle production should then be independent of cms energy for low energy particles.⁸⁾ In addition, $E \frac{dn}{d^3p}$ should go to a finite limit for $E \rightarrow m_h$; both the scaling with cms energy and the existence of a soft limit can be derived analytically in Double Log Approximation.⁴⁾ As shown in Figure 1a, both effects are seen in experimental data for charged particle spectra in a wide range of cms energies; the approximate scaling with cms energy coincides with the scaling observed by the OPAL Collaboration⁹⁾ in the variable $\ln p$ at small momenta. Let us also notice that MLLA predictions (here not shown, see^{4),6)} not only show a similar behavior, but can reproduce also quantitatively the experimental data down to very low particle energies. On the other hand, by switching off the running of the QCD coupling, MLLA predictions show a flatter slope far away from experimental behavior. Thus, the running of the QCD coupling is visible in the particle spectra even in the energy range of a few hundred MeV. In addition, the approximate scaling law and the existence of a finite soft limit strongly support QCD coherence. As shown in




Figure 1: a): Invariant distribution $E \frac{dn}{d^3p}$ for charged particle as a function of the particle energy E with $Q_0 = 270 \text{ MeV}$. Data extracted from ^{10;11}). b): same as in a) but for charged pions with $Q_0 = m$; data from ^{12;11}).

Figure 1b, the approximate scaling and the finite limit are seen also in inclusive pion spectra at different cms energies; more detailed studies on π^0 , K and p inclusive spectra are in progress.⁶⁾

In conclusion, good agreement of experimental data on moments of inclusive energy spectra in e^+e^- collisions with predictions of MLLA with running α_s plus LPHD (2 free parameters) for P_{thr} ranging from about 3 GeV to LEP-1.5 energy has been found. MLLA with fixed α_s is unable to reproduce moments both with and without absolute normalization at threshold. Data on the invariant cross section $E \frac{dn}{d^3p}$ give evidence for the existence of a finite soft limit and approximate scaling at different cms energies at particle energy of few hundreds MeV, as suggested by QCD coherence. MLLA with running α_s reproduces this effect. Direct evidence for running α_s effects in hadronic spectra, test of coherence effects and further evidence for LPHD down to low cms energies and to low particle energies have been achieved.

I thank Wolfgang Ochs and Valery A. Khoze for discussion and collaboration on the subjects of this talk. I thank J. Tran Thanh Vân, A. Capella and the Organizing Committee for the nice and stimulating atmosphere created at the Conference.

References

- [1] Yu.L. Dokshitzer, V.A. Khoze, A.H. Mueller, S.I. Troyan, "Basics of Perturbative QCD", Editions Frontieres, Gif-sur-Yvette CEDEX France (1991).
- [2] Ya. I. Azimov, Yu. L. Dokshitzer, V.A. Khoze, S.I. Troyan, Z.Phys. C 27 (1985) 65; Z.Phys. C 31 (1986) 213.
- [3] V.A. Khoze and W. Ochs, "Perturbative QCD approach to multiparticle production", preprint Durham DTP/96/36, MPIPhT-96/29 (1996).
- [4] S. Lupia and W. Ochs, Phys. Lett. B 365 (1996) 339.
- [5] V.A. Khoze, S. Lupia and W. Ochs, "Perturbative description of particle spectra at LEP-1.5", preprint Durham DTP/96/38, MPIPhT-96/28, hep-ph/9604410 (April 1996).
- [6] S. Lupia and W. Ochs, paper in preparation.
- [7] Yu. L. Dokshitzer, V.A. Khoze and S. I. Troyan, Int. J. Mod. Phys. A 7 (1992) 1875.
- [8] V.A. Khoze, in "Z⁰ Physics", Cargèse 1990, Eds. M. Levy, J-L. Basdevant, M. Jacob, D. Speiser, J. Weyers and R. Gastmans, NATO ASI, Series B: Physics Vol 261, p. 419.
- [9] MARK I Coll., J. L. Siegrist et al., Phys. Rev. D 26 (1982) 969;
 MARK II Coll., J. F. Patrick et al., Phys. Rev. Lett. 49 (1982) 1232;
 TASSO Coll., W. Braunschweig et al., Z.Phys. C 47 (1990) 187;
 L3 Coll., B. Adeva et al., Phys. Lett. B 259 (1991) 199;
 ALEPH Coll., D. Buskulic et al., CERN-PPE/96-43, March 1996;
 DELPHI Coll., P. Abreu et al., Phys. Lett. B 372 (1996) 172;
 OPAL Coll., M. Z. Akrawy et al., CERN-PPE/96-47, March 1996.
- [10] TOPAZ Coll., R. Itoh et al., Phys. Lett. B 345 (1995) 335;
 ALEPH Coll., D. Buskulic et al., Z.Phys. C 55 (1992) 209;
 DELPHI Coll., P. Abreu et al., Phys. Lett. B 347 (1995) 447;
 OPAL Coll., M. Z. Akrawy et al., Phys. Lett. B 247 (1990) 617.
- [11] ADONE Coll., B. Esposito et al., Lett. Nuovo Cimento 19 (1977) 21;
 DASP Coll., R. Brandelik et al., Phys. Lett. B 67 (1977) 358;
 ARGUS Collaboration, S. Behrends et al., Phys. Rev. D 31 (1985) 2161;
 TPC/2 Collaboration, H. Aihara et al., Phys. Rev. Lett. 61 (1988) 1263.