

Tadpole Method and Supersymmetric $O(N)$ Sigma Model

Tomohiro Matsuda

Department of Physics, University of Tokyo
Bunkyo-ku, Tokyo 113, Japan

Abstract

We examine the phase structures of the supersymmetric $O(N)$ sigma model in two and three dimensions by using the tadpole method. Using this simple method, the calculation is largely simplified and the characteristics of this theory become clear. We also examine the problem of the fictitious negative energy state.

1 Introduction

Many years ago, Gross and Neveu [1] have shown that dynamical symmetry breaking is possible in asymptotically free field theories. They obtained an expansion in powers of $1/N$ that is non-perturbative in g^2 . This leads to a massive fermion and to a bound state at threshold.

Polyakov [2] has pointed out that the $O(N)$ sigma model is asymptotically free and that the fundamental particle acquires a mass for $N > 2$.

Witten [3] has constructed a supersymmetric version of the two-dimensional $O(N)$ sigma model. This is a hybridization of the non-linear sigma model and Gross-Neveu model with Majorana fermions.

There comes a natural question: What is the difference between non-supersymmetric models and supersymmetric ones? If there is any difference, how is it realized? Many

authors tried to answer this question [4, 5], but some questionable arguments are still left. The problem of the negative energy state is one of them [6, 7]. To maintain the positivity of the vacuum energy, inclusion of the chiral condensation effect was crucial in ref[7]. However in the three dimensional model there is a weak coupling phase where the chiral condensation vanishes but the bosonic condensation is still possible.

The purpose of this paper is to clarify these ambiguities and present a systematic treatment of this model. To show explicitly what is going on, we are not going to use the equation of motion for supersymmetric auxiliary fields at the first stage. If we eliminate these fields, it becomes difficult to find what relations we are dealing with.

2 Review of the non-linear sigma model

In this and the next section we are going to review well-known results on the $O(N)$ non-linear sigma model and the four-fermion model for the convenience of checking the notations. If readers feel boring, please skip to section 4.

The Lagrangian for the $O(N)$ sigma model is defined by

$$L = \frac{1}{2} n_j \partial^2 n_j \quad (2.1)$$

with the local non-linear constraint

$$n_j n_j = \frac{N}{g^2} : \quad (2.2)$$

The sum over the flavor index j runs from 1 to N . This constraint can be implemented by introducing a Lagrange multiplier φ .

Let us consider the Euclidean functional integral in the form :

$$\begin{aligned} Z &= \int \mathcal{D}n \int \mathcal{D}\varphi \exp \left[-\frac{1}{2} \int d^D x (\partial n)^2 - \int d^D x \left(\varphi \left(n^2 - \frac{N}{g^2} \right) \right) \right] \\ &= \int \mathcal{D}\varphi \int \mathcal{D}n \exp \left[-\frac{1}{2} \int d^D x (\partial n)^2 - \int d^D x \varphi \left(n^2 - \frac{N}{g^2} \right) \right] \end{aligned} \quad (2.3)$$

The integral over n is Gaussian and can be performed in a standard fashion. We have:

$$Z = \int \mathcal{D}\varphi \exp \left[-\frac{1}{2} \int d^D x (\partial \varphi)^2 - \frac{N}{2} \int d^D x \ln \left(\varphi^2 + \frac{N}{g^2} \right) \right] \quad (2.4)$$

Let us first compute the variation of the action with respect to ϕ . We get [8]:

$$\begin{aligned}\frac{N}{2g^2} &= \frac{N}{2} \text{tr} \ln (\partial^2 + m^2) \\ &= \frac{N}{2} G(x; x; 0)\end{aligned}\quad (2.5)$$

Here we have introduced the Green function:

$$G(x; y; 0) = \langle y | (\partial^2 + m^2)^{-1} | x \rangle \quad (2.6)$$

The meaning of the above equation becomes transparent if we notice that

$$\begin{aligned}\langle n_i(x) n_j(y) \rangle &= \frac{1}{Z} \int \mathcal{D}\phi \exp \left[-\frac{1}{2} \int d^D x (\partial \phi)^2 - \frac{N}{g^2} \int d^D x \phi^2 \right] \\ &= \frac{\int \mathcal{D}\phi \phi_i(x) \phi_j(y) e^{-W}}{\int \mathcal{D}\phi e^{-W}} \\ W &= \frac{N}{2g^2} \int d^D x \phi^2 - \frac{N}{2} \text{tr} \ln (\partial^2 + m^2): \end{aligned}\quad (2.7)$$

If integration is to be approximated by the saddle point ϕ_0 , we obtain

$$\langle n_i(x) n_j(y) \rangle = \phi_0(x) \phi_0(y) \quad (2.8)$$

These equations show that eq.(2.5) is nothing but the condition $\langle \phi^2 \rangle = \frac{N}{g^2}$. This is the main idea of the tadpole method [9]. Let us now solve eq.(2.5). Passing to the momentum representation,

$$\begin{aligned}G(x; y; 0) &= \int \frac{d^D p}{(2\pi)^D} \frac{e^{ip(x-y)}}{p^2 + m^2} \\ \frac{N}{2g^2} &= \frac{N}{2} G(x; x; 0) \\ &= \frac{N}{2} \int \frac{d^D p}{(2\pi)^D} \frac{1}{p^2 + m^2}: \end{aligned}\quad (2.9)$$

For $D=2$ we obtain:

$$\begin{aligned}1 &= \frac{g^2}{4} \log \frac{4}{m^2} \\ m^2 &= \frac{4}{g^2} \exp \left(-\frac{4}{g^2} \right)\end{aligned}\quad (2.10)$$

For $D=3$, the situation is slightly different. We have a critical coupling g_{cr}^2 defined by

$$1 = g_{cr}^2 \int \frac{d^3 p}{(2\pi)^3} \frac{1}{p^2}: \quad (2.11)$$

If $g^2 > g_{cr}^2$ then the equation has a non-trivial solution $\phi_0 \neq 0$. Using g_{cr} , we can rewrite (2.9) as:

$$\begin{aligned} 1 &= g^2 \int \frac{d^3 p}{(2\pi)^3 p^2} - g^2 \int \frac{d^3 p}{(2\pi)^3} \frac{1}{p^2} \frac{1}{p^2 + \phi_0^2} \\ &= \frac{g^2}{g_{cr}^2} - g^2 \int \frac{d^3 p}{(2\pi)^3} \frac{\phi_0^2}{p^2 (p^2 + \phi_0^2)} \end{aligned} \quad (2.12)$$

The integral in (2.12) is convergent and proportional to $\frac{3}{2} \phi_0^{-1} = \frac{1}{\phi_0}$. Therefore, we have:

$$m_n^2 \phi_0 = \text{const} : \frac{g^2}{g_{cr}^2} \phi_0^2 \quad (2.13)$$

If we take $g^2 < g_{cr}^2$ something goes wrong with (2.12). It does not have any solution, so the constraint $\langle n^2 \rangle = \frac{N}{g^2}$ cannot be satisfied.

We should also consider the possibility of spontaneous breaking of $O(N)$ symmetry. In above discussions, we have implicitly assumed that the vacuum expectation value of n would vanish. Let us consider what may happen if n itself gets non-zero vacuum expectation value. Because of $O(N)$ symmetry, the vacuum expectation value of n ($n_1; n_2; \dots; n_N$) may be written as

$$\langle n \rangle = (0; 0; \dots; \frac{1}{\sqrt{N}} v = g) : \quad (2.14)$$

So that the constraint equation (2.5) becomes

$$\begin{aligned} \langle n^2 \rangle &= \langle n \rangle^2 + \langle 1 \text{ loop} \rangle \\ &= N \frac{v^2}{g^2} + \int \frac{d^3 p}{(2\pi)^3} \frac{1}{p^2 + \phi_0^2} \\ &= \frac{N}{g^2} : \end{aligned} \quad (2.15)$$

Of course, in two dimensions we cannot expect n to get any expectation value. For $D = 3$, we have another critical coupling g_{cr}^0 :

$$\frac{1}{g_{cr}^0} \frac{v^2}{g^2} = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{p^2} \quad (2.16)$$

If g is smaller than g_{cr} , then v grows. As a result, the constraint equation has a solution in the weak coupling region ($g_{cr}^0 < g < g_{cr}$) in a sense that not eq.(2.9) but eq.(2.15) is satisfied by some ϕ_0 . As far as we are dealing with the non-supersymmetric sigma model, we have no primary reason to believe that the vacuum expectation value of the field $v = \langle n_j \rangle$ would not obtain a non-vanishing value in the strong coupling region in three dimensions.

3 Review of the four-fermion model

The four-fermion model is described by the Lagrangian

$$\mathcal{L} = \frac{i}{2} \sum_j \bar{\psi}_j \not{\partial} \psi_j + \frac{g^2}{8N} (\sum_j \bar{\psi}_j \psi_j)^2 \quad (3.1)$$

where the sum of the flavor index j runs from 1 to N and we require that g remains constant as N goes to infinity. By introducing a scalar auxiliary field we may rewrite (3.1) as

$$\mathcal{L} = \frac{i}{2} \sum_j \bar{\psi}_j \not{\partial} \psi_j + \frac{1}{2} \sum_{j,j'} \bar{\psi}_j \psi_j \frac{N}{2g^2} \psi_{j'} \bar{\psi}_{j'} \quad (3.2)$$

Let us consider the functional integral in the form :

$$Z = \int \prod_j D\psi_j D\bar{\psi}_j \exp \int d^D x \left[\frac{1}{2} \sum_j \bar{\psi}_j (i \not{\partial} + \phi) \psi_j - \frac{N}{2g^2} (\sum_j \bar{\psi}_j \psi_j)^2 \right] \quad (3.3)$$

Integrating over the field ψ_j we get an effective action for the field ϕ :

$$Z = \int D\phi \exp \int d^D x \left[-\frac{N}{2g^2} \phi^2 + \frac{N}{2} \text{Tr} \ln (i \not{\partial} + \phi) \right] \quad (3.4)$$

We impose the stationary condition which gives the gap equation.

$$\frac{N}{g^2} \langle \phi \rangle = \frac{N}{2} \int \frac{d^D p}{(2\pi)^D} \text{tr} \frac{1}{\not{p} + \langle \phi \rangle} = 0 \quad (3.5)$$

As is in the non-linear sigma model discussed in the previous section, this represents the condition

$$\frac{N}{g^2} \langle \phi \rangle = \frac{1}{2} \langle \sum_j \bar{\psi}_j \psi_j \rangle \equiv \chi = \langle \phi \rangle : \quad (3.6)$$

For $D=2$ we obtain :

$$\begin{aligned} \frac{1}{g^2} &= \int \frac{d^2 p}{(2\pi)^2} \frac{1}{p^2 + \langle \phi \rangle^2} \\ \langle \phi \rangle^2 &= \int \frac{d^2 p}{(2\pi)^2} \frac{1}{p^2 + \langle \phi \rangle^2} \end{aligned} \quad (3.7)$$

For $D=3$, we have a critical coupling constant. The saddle point exists only within the branch

$$0 < \frac{1}{g^2} < \frac{1}{g_{cr}^2} \quad (3.8)$$

where

$$\frac{1}{g_{cr}^2} = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{p^2} : \quad (3.9)$$

4 Phases in the Supersymmetric Non-Linear Sigma Model

The supersymmetric non-linear sigma model is usually defined by the Lagrangian

$$L = \frac{1}{2} \sum_j \dot{x}_j^2 + \frac{1}{2} \sum_j D_j^2 \quad (4.1)$$

with the non-linear constraint

$$\sum_j x_j x_j = \frac{N}{g^2} \quad (4.2)$$

where the sum of the flavor index j runs from 1 to N . The superfields x_j may be expanded out in components

$$x_j = n_j + i \sigma_j + \frac{1}{2} F_j \quad (4.3)$$

and the supercovariant derivative is

$$D = \frac{\partial}{\partial \theta} - i \sigma \quad (4.4)$$

In order to express the constraint (4.2) as a function, we introduce a Lagrange multiplier superfield λ .

$$\lambda = \lambda + i \sigma + \frac{1}{2} F \quad (4.5)$$

We thus arrive at the manifestly supersymmetric action for the supersymmetric sigma model.

$$S = \int d^4x \left[\frac{1}{2} \sum_j \dot{x}_j^2 + \frac{1}{2} \sum_j D_j^2 + \frac{N}{g^2} \lambda \right] \quad (4.6)$$

In component form, the Lagrangian from (4.6) is

$$L = \frac{1}{2} \sum_j \dot{n}_j^2 + \frac{1}{2} \sum_j \dot{\sigma}_j^2 + \frac{1}{2} \sum_j F_j^2 - \sum_j n_j F_j - \frac{1}{2} \sum_j n_j^2 + \frac{1}{2} \sum_j \sigma_j^2 + \frac{N}{2g^2} \lambda \quad (4.7)$$

We can see that λ , σ_j and F_j are the respective Lagrange multiplier for the constraints:

$$\begin{aligned} n_j n_j &= \frac{N}{g^2} \\ n_j \sigma_j &= 0 \\ n_j F_j &= \frac{1}{2} \sum_j \sigma_j^2 \end{aligned} \quad (4.8)$$

The second and the third constraints of (4.8) are supersymmetric transformations of the first. We must not include kinetic terms for the field and so as to keep these constraints manifest. We can examine these constraints in a way that we did in the previous section.

(1) Scalar part

$$\langle n_j n_j \rangle = \langle n_n = \langle \phi^2 \rangle + \langle \phi'^2 \rangle = \frac{N}{g^2} \quad (4.9)$$

In two dimensions, this relation induces nonzero value to the mass term of the field n .

$$\begin{aligned} m_n &= \langle \phi^2 \rangle + \langle \phi'^2 \rangle \\ &= \frac{4}{g^2} \exp \left(-\frac{1}{g^2} \right) \end{aligned} \quad (4.10)$$

When $D=3$, m_n is nonzero in the region $g > g_{cr}$. The critical coupling is defined by

$$\frac{1}{g_{cr}^2} = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{p^2} : \quad (4.11)$$

$O(N)$ symmetry is expected to be spontaneously broken by non-zero value of v in the region $g < g_{cr}$. And when $g = g_{cr}^0$, m_n would vanish.

(2) Fermionic part

$$\langle n_j F_j \rangle = \frac{1}{2} \langle \bar{\psi}_j \psi_j \rangle \quad (4.12)$$

This relation includes auxiliary field F_j , to be eliminated by equation of motion. After substituting F_j by $\bar{\psi}_j$, we obtain at one-loop level:

$$\begin{aligned} \langle n_j F_j \rangle &= \langle n_j \bar{\psi}_j \psi_j \rangle \\ &= \langle \phi^2 \rangle \langle n_j \bar{\psi}_j \psi_j \rangle \\ &= \frac{1}{2} \langle \bar{\psi}_j \psi_j \rangle \end{aligned} \quad (4.13)$$

If we impose the $O(N)$ symmetric constraint $\langle n^2 \rangle = \frac{N}{g^2}$, we have

$$\begin{aligned} \frac{N}{g^2} \langle \phi^2 \rangle &= \frac{1}{2} \langle \bar{\psi}_j \psi_j \rangle \langle n^2 \rangle \\ \frac{N}{g^2} &= N \int \frac{d^D p}{(2\pi)^D} \frac{1}{p^2 + \langle \phi^2 \rangle} : \end{aligned} \quad (4.14)$$

For $D=2$, the solution is

$$\langle \phi^2 \rangle = \frac{4}{g^2} \exp \left(-\frac{1}{g^2} \right) : \quad (4.15)$$

Substituting $\langle \phi \rangle$ in the first constraint (4.9) with (4.15), we can find that $\langle \phi \rangle$ must vanish. (in this point our result is different from [5]) This means that the field gains the same mass as n , and simultaneously supersymmetric order parameter $\langle \phi \rangle$ vanishes. We can say that the supersymmetry is not broken in two dimensions as is predicted by Witten [10]. Moreover, we can examine the assumption of vanishing v as follows. We can show that the following relation can exist for the effective potential [6].

$$\begin{aligned} \frac{\partial V}{\partial v} &= N v (\phi_0 + \langle \phi \rangle^2) \\ &= 0 \end{aligned} \quad (4.16)$$

This means that v must vanish if chiral condensation occurs.

For $D = 3$, we have a critical coupling constant. As far as $g < g_{cr}$, we have nothing to worry about. In the strong coupling region, both supersymmetry and $O(N)$ symmetry are preserved in a fashion like two dimensions. However, in the weak coupling region, something goes wrong. There is no non-trivial solution for constraint (4.12) and there is no fermionic condensation (This means that the only possible solution is $\langle \phi \rangle = 0$). Thus we can see from eq.(4.16) that v can be non-zero in this weak coupling region. This is supported by the constraint (4.9) because this does not have any solution in the weak coupling region unless we allow v not to vanish. Eq.(2.16) suggests:

$$v^2 = 1 - g_{cr}^2 \int \frac{d^3 p}{(2\pi)^2} \frac{1}{p^2} \quad (4.17)$$

Naive consideration also supports this analysis. In general, we can expect that quantum effects in correlation functions like $\langle n_j n_j \rangle$ or $\langle \bar{\psi} \psi \rangle$ would vanish in the weak coupling limit. But we have an $O(N)$ symmetric constraint. It is natural to think that the field n itself gains expectation value to complement quantum effects. This simply means that classical effects become dominant in the weak coupling region, therefore the $O(N)$ symmetric constraint is satisfied classically. (i.e. in the weak coupling limit $g \rightarrow 0$ we obtain $v = 1$. This is a classical solution of the constraint.) As a result, in the weak coupling region, $O(N)$ symmetry is spontaneously broken by non-zero value of v .

We should also note that there is a possible solution of non-zero ϕ_0 . (We neglect eq.(4.16) for a while because ϕ_0 may become a function of v .) It induces a supersymmetry

breaking term to the Lagrangian:

$$L_{\text{break}} = -\frac{1}{2} \left(\frac{N}{g^2} \right)^2 \quad (4.18)$$

On the constrained phase space $(\mathbf{n})^2 - \frac{N}{g^2} = 0$, vacuum energy also seems to vanish for non-zero v_0 as long as the constraint (4.9) is satisfied. Does it mean there is a flat direction along v_0 ? Of course this statement is unnatural. After including effective kinetic term $(\partial_\mu v)^2$, we can find positive vacuum energy for the supersymmetry breaking phase. Therefore in the supersymmetric model, v is not a free parameter but fixed by the requirement of vanishing v_0 . This means g_{cr}^0 should be adjusted to g , and v is fixed:

$$v^2 = 1 - g^2 \int \frac{d^3 p}{(2\pi)^3} \frac{1}{p^2} \quad (4.19)$$

So we can conclude:

(1) In two dimensions, both supersymmetry and $O(N)$ symmetry are not broken. This means that v and v_0 remain zero for all value of g .

(2) In three dimensions, both supersymmetry and $O(N)$ symmetry are not broken (i.e. v and v_0 remain zero) in the strong coupling region. $O(N)$ symmetry can be broken in the weak coupling region, but supersymmetry is kept unbroken in both phases.

5 Negative Energy

In this section, we will reconsider whether negative energy states in supersymmetric theories [7, 11] can exist or not. One may wonder why such a state appears, but it is really a confusing matter. Because we have not enough space, we refer [7] in which detailed analysis on this topic can be found. In ref. [7], two dimensional supersymmetric non-linear sigma model and supersymmetric Yang-Mills model are analyzed.

For us, the main problem is the value of v . Naively calculated 1-loop effective potential shows that it has negative energy state at $v \neq 0$. For example in $D = 3$ [6],

$$V = \frac{N}{2} \left(v^2 - \frac{1}{g^2} \right)^2 + v^2 \int \frac{d^3 k}{(2\pi)^3} \ln(k^2 + v^2) + \int \frac{d^3 k}{(2\pi)^3} \text{tr} \ln(k^2 + v^2) \quad (5.1)$$

We can think that this problem comes from the instability of the tree level potential $V = (n^2 - N/g^2)$ along the direction of ϕ . In general we have to set $\phi = 0$ by hand, but it should be determined by considering some effects.

First, we are going to examine two dimensional non-linear sigma model from a different point of view.

We can calculate an effective potential for ϕ in two dimensional $O(N)$ supersymmetric non-linear sigma model using trace anomaly equation [7]. As a result, we can obtain:

$$V(\phi) = -\frac{N}{8} \ln \frac{\phi^2}{\Lambda^2} + \text{const} \quad (5.2)$$

This potential has unnatural characteristics like negative energy or unstable vacuum.

This term can appear in the effective action at 1-loop level (we should note the meaning of ϕ is somewhat different from eq.(5.2)). Not yielding to a trace anomaly equation, after integrating over n we can obtain:

$$Z = \int \mathcal{D}\phi \mathcal{D}n \exp \int d^2x \left[\frac{i}{2} \phi_j \partial_\mu \phi_j + \frac{1}{2} \partial_\mu \phi_j \partial_\mu \phi_j - \frac{1}{2} \phi_j^2 + \frac{N}{2g^2} \right] \frac{N}{2} \text{tr} \ln (\partial^2 + \phi_j^2) \quad (5.3)$$

Integration can be done for the last term and we can obtain the same result as (5.2) except for ϕ^2 which appeared in the mass term.

But there are some problems. First, the effective action we derived does not include fermionic loop corrections that leads to the chiral condensation. Including the fermionic loop corrections, we can reach at the result we have obtained in the previous section. The vacuum state is supersymmetric and there is no negative vacuum energy. To simplify the argument, it is very useful to separate every constraint and discuss each property as we have done in section 4.

Second problem is the treatment of the effective action. Usually we think that after integrating out n fields the integration over ϕ cannot be done exactly so we always consider a stationary phase approximation. To actually determine the stationary point, we vary with respect to the constant value of ϕ . The resulting equation is the gap equation:

$$\frac{N}{g^2} = N \int \frac{d^2p}{(2\pi)^2} \frac{1}{p^2 + \phi_0^2 + \langle \phi^2 \rangle} \quad (5.4)$$

But there is a problem. ϕ is a Lagrange multiplier so its tree level potential is not stable for $\phi = 0$. Naively calculating the 1-loop potential, we will find (continuous) negative

energy state. In general supersymmetric non-gauge theories, $F_j F_j$ type term in the kinetic term $(-1/2) D^2 \phi_j$ is responsible for the positivity of the vacuum energy. With this term, scalar potential is always written as $V = \frac{1}{2} W_{ij}^2$. In our model, this term will appear in the effective kinetic term and is responsible for the positivity of the vacuum energy. Of course, there is a possibility that the kinetic term would be a non-trivial (special) function of ϕ . Then, the positivity of the vacuum energy is not manifest and the argument of negative energy would be trustworthy. (But in our approximation such a term does not appear.) Moreover, as we have shown in the previous section, the stationary point ϕ_0 is exactly determined by fermionic constraint in two dimensions and resulting effective potential $V^{\text{eff}}(\phi_j)$ vanishes in the stationary phase approximation.

Can we apply the same argument to the three dimensional model? Naively calculating the 1-loop effective potential, a negative energy state appears in the wrong vacuum $\phi \neq 0$ even if we consider the fermionic condensation. In this case, we must also consider the effective kinetic term that yields effective term. Including this, we can expect that the scalar potential is always positive.

6 Conclusion

Some authors claimed that in supersymmetric models, there can be a supersymmetry breaking accompanied by negative energy and negative norm states that lead to other instabilities. It is true that we cannot ignore such a possibility in general but we can make sure of the absence of such a vacuum at least in $O(N)$ sigma model in two and three dimensions.

Merely adding the Lagrange multiplier fields and taking it as a scalar potential, we would be led to unnatural arguments. If relating the Lagrange multiplier to the potential is necessary, we should have considered about the effective kinetic terms. Of course, we must be careful not to forget to include both fermionic and bosonic loops [7, 12]. The same can be said for the analysis of supersymmetric Yang-Mills or supersymmetric QCD theories. Decomposed in component fields, these theories look like ordinary QCD with Majorana fermions or that with Higgs fields. So we tend to forget their origin and analyze these theories in usual way of QCD.

We have analyzed the phase structures of $O(N)$ supersymmetric sigma model in two and three dimensions by using the tadpole method.

We have shown that after including fermionic constraint and a effective kinetic term, is determined as $\mu = 0$ and the supersymmetry breaking vacuum has positive energy. There is no fear of negative energy states at least in $O(N)$ sigma model discussed above.

Acknowledgment

We thank K. Fujikawa, Y. Kazama, S. Iso and K. Hori for many helpful discussions.

References

- [1] D. J. Gross and A. Neveu, Phys. Rev. D 10 (1974) 3235
- [2] A. M. Polyakov, Phys. Lett. 59B (1975) 79
- [3] E. Witten, Phys. Rev. D 16 (1977) 2991
- [4] A. C. Davis, P. Salamonson and J. W. Holten, Nucl. Phys. B 208 (1982) 484
- [5] O. Alvarez, Phys. Rev. D 17 (1978) 1123
- [6] V. G. Koubes and K. T. Mahanthappa, Phys. Rev. D 43 (1991) 3428
- [7] H. Hata and Y. Kazama, Nucl. Phys. B 232 (1984) 85
A. Higuchi and Y. Kazama, Nucl. Phys. B 206 (1982) 152
- [8] A. M. Polyakov Gauge Fields and Strings, Harwood Academic Publishing (1987)
- [9] R. Miller, Phys. Lett. 124B (1983) 59, Nucl. Phys. B 241 (1984) 535
S. Weinberg Phys. Rev. D 7 (1973) 2887
- [10] E. Witten, Nucl. Phys. B 202 (1982) 253
- [11] Ulrich Ellwanger Z. Phys. C 18 (1983) 81
R. K. Kaul and L. M. Izrachi, J. Phys. G 15 (1989) 1633
- [12] T. Matsuda, J. Phys. A 28 (1995) 3809