

The Lowest Order Hadronic Contribution to the Muon $g - 2$ Value with Systematic Error Correlations

D.H. Brown and W.A. Worstell

Physics Department, Boston University, 590 Commonwealth Avenue, Boston, MA 02215
(11 June 1996)

We have performed a new evaluation of the hadronic contribution to $a_\mu = (g - 2)/2$ of the muon with explicit correlations of systematic errors among the experimental data on $\sigma(e^+e^- \rightarrow \text{hadrons})$. Our result for the lowest order hadronic vacuum polarization contribution is $a_\mu^{\text{had}} = 702.6(7.8)(14.0) \times 10^{-10}$ where the first error is statistical and the second is systematic. The total systematic error contributions from below and above $\sqrt{s} = 1.4$ GeV are $(13.1) \times 10^{-10}$ and $(5.1) \times 10^{-10}$ respectively, and are hence dominated by the low energy region. Therefore, new measurements on $\sigma(e^+e^- \rightarrow \text{hadrons})$ below 1.4 GeV can significantly reduce the total error on a_μ^{had} . In particular, the effect on the total errors of new hypothetical data with 3 % statistical and 0.5 - 1.0 % systematic errors is presented.

I. ROLE OF HADRONIC CONTRIBUTION IN $G - 2$

A new measurement of the anomalous magnetic moment of the muon, $a_\mu \equiv (g-2)/2$, to an absolute accuracy of $\sigma_{exp}^{a_\mu} \sim \pm 4.0 \times 10^{-10}$ is proposed by the AGS E821 Collaboration at Brookhaven National Laboratory [1] [2]. The theoretical value of the muon $g-2$ value consists of at least the three standard model contributions: quantum electrodynamics (QED), electroweak (EW) and hadronic. The latter arise from hadronic vacuum polarization effects caused by effective photon couplings to hadrons via charged quarks and consequent quantum chromodynamics (QCD) interactions with gluons.

Any residual difference between the sum of the standard model contributions and the new experimental value a_μ^{exp} will be indicative of new physics:

$$a_\mu^{\text{residual}} = a_\mu^{\text{exp}} - a_\mu^{\text{QED}} - a_\mu^{\text{EW}} - a_\mu^{\text{had}}.$$

The new experimental value can only be sensitive to electroweak and possibly supergravity [3] [4] and muon substructure effects [5] provided the errors on the standard model contributions are known better than the experimental accuracy. The QED and EW contributions have been calculated from theory and are known an order of magnitude better than the expected experimental accuracy ($a_\mu^{\text{QED}} = 11658470.6 \pm 0.2 \times 10^{-10}$ [6] and $a_\mu^{\text{EW}} = 15.1 \pm 0.4 \times 10^{-10}$ [7]). At the same time, the lowest order hadronic contribution cannot be calculated accurately enough by QCD and hence a phenomenological procedure must be used for its calculation. While the results of the published a_μ^{had} calculations have errors which vary from just above E821 error to greater than the electroweak contribution (see Tab. I) and even differ in principle of approach, it is of interest for the interpretation of a new experimental measurement of the muon

$g - 2$ value to investigate in more detail the precise procedure used for calculation of the hadronic contribution total error.

Fortunately the hadronic contribution to the muon $g - 2$ value can be related to a dispersion integral over the experimental cross section $\sigma(e^+e^- \rightarrow \text{hadrons})$ (see Sect. II). The error on the hadronic contribution to a_μ is hence determined by the error on the experimental cross section and is by all accounts larger than the expected E821 accuracy. Therefore, the hadronic contribution error will largely determine the sensitivity level of E821 to new physics:

$$\sigma_{\text{residual}} = \sqrt{\sigma_{\text{exp}}^2 + \sigma_{\text{QED}}^2 + \sigma_{\text{EW}}^2 + \sigma_{\text{had}}^2} \sim \sqrt{\sigma_{\text{exp}}^2 + \sigma_{\text{had}}^2}.$$

This is the motivation for several recent evaluations of the lowest order hadronic contribution to a_μ which are presented in Tab. I. These calculations may roughly be placed into two categories of approach whether they are based primarily upon (aggressive) model dependent techniques which yield relatively smaller errors [8] [9] [10] or on (conservative) model independent techniques (trapezoidal integration) which yield relatively larger errors [11] [12]. A typical method for evaluating systematic error is comparison of the model-dependent calculation with one based on trapezoidal integration.

While the model dependent calculations are based on theoretical innovations to represent various components of the cross section $\sigma(e^+e^- \rightarrow \text{hadrons})$ they lack the merit of a clear prescription for correlating the systematic errors of the experimental input. These errors are in some cases as large as 20 % in overall normalization of the cross section. Therefore, the χ^2 criterion used to select a set of fit parameters, although indicative of the best overall fit to the data, may perhaps not indicate the true error on the integral over experimental data points which is the hadronic contribution to muon $g - 2$. In particular, a shift of the whole curve up or down may fall

within the experimental data point errors and yet yield a variation in a_μ^{had} larger than the stated errors.

The model independent techniques for calculating a_μ^{had} are based upon trapezoidal integration over the experimental data points. As will be described below, this approach allows for a definite procedure for determining both the central value for a_μ^{had} and its errors both of which make explicit use of the statistical and systematic errors for each experimental data point input. In particular, for each experiment an error weighted fraction will be defined and used in averaging over different experiments which make measurements of the same cross sections over common energy regions.

The most recent model independent analysis [12] was the first to account for systematic error correlations among the experimental data used in the a_μ^{had} calculation. The systematic errors were correlated at each energy point separately in the process of determining an error-weighted average cross section ratio $R(s)$ (see Sect. II) *before* performing the $g-2$ integral. Special care was taken to calculate the ρ -meson contribution separately from the rest of the hadronic production cross sections and there are many significant contributions to the discussion of a_μ^{had} calculations in this important communication.

The main new aspect of the present a_μ^{had} calculation is that the systematic errors are correlated by a different error combination formula *after* performing the $g-2$ integral. Furthermore, correlations are accounted for across energy regions and between experimental measurements of the ρ -meson and of the additional hadronic cross sections which contribute to $\sigma(e^+e^- \rightarrow \text{hadrons})$. (It appears that by treating the ρ -meson separately the previous calculation did not correlate some errors completely.)

In short, the $g-2$ integral has been performed over each hadronic cross section separately without combining them into their equivalent $R(s)$ value for the present calculation. The focus is therefore upon the error-weighted average of the a_μ^{had} integral value itself (and its errors).

These innovations have yielded a result with total error remarkably equal to that obtained previously [12]. However, there is a difference obtained as to how much of the error derives from the energy region above and below 1.4 GeV which calls attention to the precise procedure of error combination employed in the calculation.

II. BACKGROUND ON A_μ^{HAD} CALCULATIONS

The hadronic contribution to a_μ consists of a dominant lowest order term, shown in Fig. 1, several higher order terms in Fig. 2 (the number above each diagram indicates how many contributing diagrams are in its class), and finally a group of hadronic light-by-light scattering terms. Detailed calculations are given elsewhere of the Figure 2 and hadronic light-by-light contributions. Since their errors are well below the expected E821 error of $\sigma_{exp}^{a_\mu} \sim \pm 4.0 \times 10^{-10}$ ($a_\mu^{Fig2} = -9.0 \pm 0.5 \times 10^{-10}$ [11] and $a_\mu^{light} = -5.2 \pm 1.8 \times 10^{-10}$ [13]) they are not discussed here. There is another hadronic light-by-light calculation with a larger error $a_\mu^{light} = -9.2 \pm 3.2 \times 10^{-10}$ [14] but it does not depend on the cross section $\sigma(e^+e^- \rightarrow \text{hadrons})$ and therefore is not of concern to the present calculation. In view of the difference between the two latest hadronic light-by-light contributions, and the consequent ambiguity over defining the *total* hadronic contribution, this paper is concerned only with the lowest order hadronic contribution to the muon $g-2$ value shown in Fig. 1.

A. Formalism of the Hadronic Contribution a_μ^{had}

The largest contribution to a_μ^{had} , shown in Fig. 1, can be related to the total Born cross section (lowest order in QED) for hadron production in electron-positron annihilations, $\sigma_{had}^0 = \sigma^0(e^+e^- \rightarrow \text{hadrons})$, by means of dispersion theory and the optical theorem [15]. Defining $\xi \equiv s/m_\mu^2$ and $\beta \equiv \sqrt{1-4/\xi}$ the result is

$$a_\mu^{had} = \frac{1}{4\pi^3} \int_{4m_\pi^2}^{\infty} ds \sigma_{had}^0 K(s) = \frac{m_\mu^2}{9} \left(\frac{\alpha}{\pi}\right)^2 \int_{4m_\pi^2}^{\infty} ds R(s) K_2(s) \frac{1}{s^2} \quad (1)$$

where the kernel function $K(s)$ in general arises from a massive photon propagator in the $g-2$ Schwinger calculation:

$$K(s) = \int_0^1 dx \frac{x^2(1-x)}{x^2 + (1-x)\xi} \xrightarrow{s \rightarrow \infty} \frac{m_\mu^2}{3s}$$

$$K(s) = \frac{1}{2\beta} \left[\frac{(1-\beta)^2}{1+\beta} \left(1 - \frac{2}{1+\beta} \ln \frac{2}{1-\beta} \right) - \frac{(1+\beta)^2}{1-\beta} \left(1 - \frac{2}{1-\beta} \ln \frac{2}{1+\beta} \right) \right] - \frac{1}{2}. \quad (2)$$

The cross section ratio $R(s) \equiv \sigma_{had}^0/\sigma_{\mu\mu}^0$ with $\sigma_{\mu\mu}^0 = 4\pi\alpha^2/3s$ and the kernel $K_2(s)$ is used in the $R(s)$ formulation of the a_μ^{had} dispersion integral:

$$K_2(s) \equiv \frac{3s}{m_\mu^2} K(s).$$

The σ_{had} formulation is useful for low energy data which are usually published as individual exclusive hadronic cross sections; the $R(s)$ form of the a_μ^{had} dispersion integral is useful for higher energies where experimental data are usually published as the inclusive ratio R .

From the a_μ^{had} dispersion integrals it is apparent that the error on the hadronic cross sections determines the error on a_μ^{had} . Therefore, the AGS E821 experimental error goal just larger than 3.5×10^{-10} on the hadronic contribution of roughly 700×10^{-10} requires a 0.5 % accuracy on the hadronic contribution calculation.

III. THE WFS A EVALUATION PROCEDURE

A. Correlation Postulates

As is implied by the $g - 2$ dispersion integral (Eq. 1) the a_μ^{had} calculation procedure consists of some combination of the following steps: 1) integration over energy, 2) weighted average over detectors and 3) sum over exclusive hadronic modes which contribute to $\sigma^0(e^+e^- \rightarrow \text{hadrons})$. The sequence of these three sums may be interchanged as for example performing the energy integration last [12]. Most calculations, however, employ the sequence (1,2,3) – the most natural one – where in the last step, the quadrature sum of errors over modes implies the assumption of zero correlation of systematic errors among experiments (a reasonable first approximation).

To go beyond the first approximation, it is necessary to survey the existing experimental data on hadronic production cross sections. The published data and recent preprints on exclusive hadron production in electron-positron annihilations used in the present calculation are listed in Appendix A where it is evident that eight detectors have measured more than one mode. Since a given detector uses the same luminosity and similar correction factor calculations (e.g. radiative corrections, efficiencies which use some of the same subroutines) it is reasonable to suppose that the cross section determinations of different exclusive hadronic modes by a single detector may in fact be correlated. The following correlation postulates are therefore intended to address this situation:

1. A single detector measuring more than one exclusive hadronic mode has 100 % correlations among systematic errors due to common luminosity and correction factor calculations;

2. Different detectors have uncorrelated errors since they do not share luminosity and correction factor calculations.

The accommodation of these correlation postulates requires a particular sequence of the three sums (1,2,3) for combination of the a_μ^{had} central values and an alternative sequence (1,3,2) for the a_μ^{had} errors. In both cases, the energy is integrated over first and separately for each detector and exclusive hadronic mode measured over energy sub-regions, where these sub-regions are defined by common energy coverage among detectors.

The former sequence (1,2,3) facilitates the need to first calculate the error weighted fractions (defined in detail below) while the latter sequence (1,3,2) is necessary for, according to postulate 1, correlating individual detector systematic errors over the modes measured by that detector. This must be done before the final uncorrelated combination of errors is made, according to postulate 2, across detectors. As this method is based on Weighted Fraction averaging with S -factor application (see Eq. 5) After a_μ^{had} integration over the energy, it is here referred to as the WFS A method.

Lastly, in the present calculation it is noted that exclusive hadronic cross sections (modes) up to 2.0 GeV have been used because this can reveal the propagation of errors from each detector and exclusive mode separately. An additional consideration, although less important, is that the exclusive hadronic mode spectra may contain interference effects (when more than one vector meson contributes to a given exclusive hadronic mode) which would otherwise require special care if individual vector meson contributions were calculated separately. Further, the uncertainty over the generation mechanism, between the e^+e^- annihilation photon and the hadronic final state, is avoided by focusing on exclusive hadronic modes themselves.

For the energy region 2.0 to 3.1 GeV, the inclusive cross section ratio $R(s)$ has been used (in the absence of exclusive data in this energy region) with the WFS A procedure for the contributing experiments. In the region above 3.1 GeV, the QCD expression has been used without the WFS A procedure since perturbative QCD is expected to be valid (see Sect. IV D).

B. Trapezoidal Integration Procedure

The usual trapezoidal integration technique takes the experimental data points pairwise: the cross sections, systematic and squared statistical errors are averaged per pair, then multiplied by the energy width of the pair, and finally they are summed over all pairs. However, it is convenient for treatment of statistical errors to expand the sum in order to remove terms which cancel. Denoting by s_k , K_k , c_{ijk} , σ_{ijk}^{stat} , and σ_{ijk}^{sys} , the energy, kernel function,

the cross section and its statistical and systematic error from the i th detector, j th exclusive mode, at the k th energy point, the integration of Eq. (1) can be represented by:

$$a_{ij} = \frac{1}{4\pi^3} \sum_{k=1}^{n-1} \left\{ \frac{1}{2} (c_{ijk} K_k + c_{ij,k+1} K_{k+1}) \right\} (s_{k+1} - s_k)$$

$$= \frac{1}{4\pi^3} \frac{1}{2} \left(A_1 + \sum_{k=2}^{n-1} A_k + A_n \right)$$

$$A_1 = c_{ij,1} K_1 (s_2 - s_1)$$

$$A_k = c_{ij,k} K_k (s_{k+1} - s_{k-1})$$

$$A_n = c_{ij,n} K_n (s_n - s_{n-1})$$

where the first and last terms in the sum are handled separately and the middle terms have an energy width across both upper and lower neighboring data points instead of across the points in pairs. This latter form is necessary for the proper treatment of the statistical errors:

$$\sigma_{ij}^{stat} = \frac{1}{4\pi^3} \frac{1}{2} \sqrt{\sigma_1^2 + \sum_{k=2}^{n-1} \sigma_k^2 + \sigma_n^2}$$

$$\sigma_1^2 = (\sigma_{ij,1}^{stat} K_1)^2 (s_2 - s_1)^2$$

$$\sigma_k^2 = (\sigma_{ij,k}^{stat} K_k)^2 (s_{k+1} - s_{k-1})^2$$

$$\sigma_n^2 = (\sigma_{ij,n}^{stat} K_n)^2 (s_n - s_{n-1})^2.$$

The systematic errors are specified by an array of values p_{ijk}^{sys} which are given as a percentage of the total cross section. Since the a_μ^{had} contribution terms above are linear in the cross section this implies that the systematic errors should be:

$$\sigma_{ij}^{sys} = p_{ij,1}^{sys} A_1 + \sum_{k=2}^{n-1} p_{ij,k}^{sys} A_k + p_{ij,n}^{sys} A_n.$$

Usually the systematic error is the same for all energy points; however, it is different for each point of the NA7 and Olya $\pi^+\pi^-$ measurements. In the cases where no systematic errors were given or readily located in the literature, values from comparable measurements with the same detector have been taken wherever possible, or as 10 % if the statistical errors dominate. In addition, a calculation with 20 % for these ambiguous values has shown that the WFSA results do not depend on the arbitrary selection of 10 % [16]. All of the systematic errors that were used are listed in Tab. IX for $\pi^+\pi^-$, Tab. X for $\pi^+\pi^-\pi^0$, Tab. XI for the higher multiplicity modes, and Tab. XII for the energy region 2.0 - 3.1 GeV.

If the desired limits of integration are inside (outside) the given data energy range, the cross section and error are linearly interpolated (extrapolated) to give the relevant pair of points. Thus our energy ranges are variable and have been set to match each of the previous evaluations for detailed comparisons [16].

C. Weighted Fraction Averaging

In order to arrive at an a_μ^{had} contribution per exclusive mode, $a_\mu^{mode} \equiv a_j$, error-weighted averages across detectors have been performed according to the PDG [17] procedure (for the i th detector and j th exclusive hadronic mode):

$$a_j \pm \sigma_j = \frac{\sum_i w_{ij} a_{ij}}{\sum_i w_{ij}} \pm \left(\frac{1}{\sum_i w_{ij}} \right)^{1/2} \quad (3)$$

where

$$w_{ij} = \frac{1}{\sigma_{ij}^2}; \quad \sigma_{ij}^2 = (\sigma_{ij}^{stat})^2 + (\sigma_{ij}^{sys})^2. \quad (4)$$

In addition, the quality of the error weighted combinations were assessed by calculating the χ^2 and PDG scale factor (i.e. χ^2 per degree of freedom) [17]:

$$\chi_j^2 = \sum_i^N w_{ij} (a_j - a_{ij})^2$$

$$S_j \equiv \sqrt{\frac{\chi_j^2}{N-1}} \quad (5)$$

where N is the number of detectors included in the average. If $S > 1$ then the errors were scaled up by this factor.

At this point the prescription for determining the error-weighted fractional contribution to the total error from a given statistical or systematic error of the i th detector and j th exclusive mode is needed. For this purpose the PDG expression above for total squared error on a weighted average (see Eq. 3) can be expanded as a sum over squared component errors where in the first step the trivial sum $\sum_{i=1}^N = N$ is used to multiply by unity:

$$\sigma_j^2 = \frac{1}{\sum_i w_{ij}} = \frac{1}{\sum_i w_{ij}} \left(\frac{1}{N} \sum_{i=1}^N \frac{\sigma_{ij}^2}{\sigma_{ij}^2} \right)$$

$$= \sum_{i=1}^N \frac{1}{N} \frac{w_{ij}}{\sum_i w_{ij}} \sigma_{ij}^2.$$

The last step makes use of the definition $w_{ij} = 1/\sigma_{ij}^2$ while the definition for the remaining σ_{ij}^2 in the numerator (see Eq. 4) leads to the separation of squared statistical from systematic terms in the sum:

$$\sigma_j^2 = \sum_{i=1}^N \left[\left(\frac{\sigma_{ij}^{stat}}{\sigma_{ij}^2} \right)^2 + \left(\frac{\sigma_{ij}^{sys}}{\sigma_{ij}^2} \right)^2 \right]$$

$$\frac{\sigma_{ij}^{stat}}{\sigma_{ij}^2} = \sqrt{\frac{1}{N} \frac{w_{ij}}{\sum_i w_{ij}}} S \sigma_{ij}^{stat} \quad (6)$$

$$\frac{\sigma_{ij}^{sys}}{\sigma_{ij}^2} = \sqrt{\frac{1}{N} \frac{w_{ij}}{\sum_i w_{ij}}} S \sigma_{ij}^{sys}. \quad (7)$$

Note that the PDG scale factor S has been inserted to emphasize the fact that the errors are to be increased if and only if the scale factor $S > 1$. (These expressions differ from those in the previous a_μ^{had} calculation [12] by the factors $1/N$ and S .)

Armed with these expressions it is possible to implement the correlation postulates for the i th detector by summing the *averaged* systematic errors (i.e. the weighted fractional systematic error contributions to the total errors in Eq. 7) linearly over contributing exclusive modes, while leaving weighted fractional statistical errors (Eq. 6) uncorrelated. The correlation postulates are simply executed by the following sums:

$$\overline{\sigma_i^{sys}} = \sum_j \overline{\sigma_{ij}^{sys}} \quad (8)$$

$$\overline{\sigma_i^{stat}} = \sqrt{\sum_j \left(\overline{\sigma_{ij}^{stat}}\right)^2}. \quad (9)$$

Note that σ_j itself is not directly used to determine the total WFSa error; to sum σ_j over j in quadrature would yield a total error which ignores correlations. This is what most previous a_μ^{had} calculations have done.

IV. APPLICATION OF THE WFSa METHOD

A. Correlations Over Energies

In general, if different detectors measure hadronic cross sections over different energy regions, there must be different error-weighted fractions and S -factors for each common energy region (i th detector, j th energy region). Such a WFSa application over energies has been applied to the dominant a_μ^{had} contributions from the hadronic modes: $\pi^+\pi^-$, $\pi^+\pi^-\pi^0$, K^+K^- , and $K_L K_S$. An example of WFSa correlations over energy is shown in Tab. V for the $R(s)$ a_μ^{had} contribution in the energy range from 2.0 to 3.1 GeV. The a_μ^{had} central values are error-weighted averaged across detectors (horizontally in the table) and summed over energies (vertically) while the detector specific statistical (systematic) errors are combined in quadrature (linearly) across the energies (vertically). The correlation of systematic errors over energies does not appear to be taken into account by the WFB method discussed in the previous a_μ^{had} calculation [12]: *Weighted Fraction averaging* (of $R(s)$ values) *Before* energy integration.

In this energy range, published inclusive $R(s)$ values have been used from the $\gamma\gamma 2$ detector [18] [19] and MARK I [20]. It is noted that the $\gamma\gamma 2$ collaboration has published values for $R_2(s)$ (two hadrons exclusively) [18] and $R_{\geq 3}(s)$ (three or more hadrons) [19] separately which are added for the present calculation. This does not appear to have been done in the previous a_μ^{had} calculations. In addition, data published as $\sigma(e^+e^- \rightarrow \text{hadrons})$ by

the BCF collaboration [21] (not apparently included previously) has been divided by $\sigma_{\mu\mu}^0$ to make them also $R(s)$ values.

B. Correlations Over Modes

If the hadronic cross sections are measured over similar energy regions, as in the case of the > 2 hadrons multiplicity cross sections shown in Tab. IV, then the WFSa method can simply be applied over the modes (i th detector, j th hadronic mode). In this case, a single error-weighted fraction and S -factor suffices for evaluation of a_μ^{had} central value and errors. As before, the a_μ^{had} central values are error-weighted averaged across detectors (horizontally in the table) and summed over modes (vertically), while the detector specific statistical (systematic) errors are combined in quadrature (linearly) across, in this case, the modes (vertically). The errors are not to be combined across detectors until all modes have been treated separately and then taken together. It is this correlation over all modes, in particular correlating the $\pi^+\pi^-$ detector total systematic errors with all the other two body and higher multiplicity modes, over all energies, which appears not to have been done before. (The previous a_μ^{had} calculation [12] appears to have accounted for these correlations only above 0.81 GeV, after the peak of the ρ -meson. Hence, it appears that the $\pi^+\pi^-$ systematic errors below 0.81 GeV were not correlated with systematic errors of measurements by the same detectors of the other modes over all energies and of the $\pi^+\pi^-$ mode above 0.81 GeV.)

C. Kaons, Narrow Resonances and Higher VMD modes

Some additional hadronic modes which contribute to the total hadronic production cross section $\sigma^0(e^+e^- \rightarrow \text{hadrons})$ require comment. In particular, experimental data in the form of total cross sections neither exists on the radiative decays of the ω and ϕ mesons ($\pi^0\gamma, \eta\gamma$) nor on the kaon pair production of the ϕ meson below certain energies. Further, there are additional contributions to hadronic vacuum polarization than from just the lowest order single vector meson dominated (VMD) amplitudes represented by the hadronic (decay) modes previously discussed.

1. Kaons and Narrow Resonances

Experimental data on the total cross sections for production of kaon pairs (charged and neutral) are limited by the fact that nuclear interactions of low momentum kaons are not well measured. Hence kaon detection efficiencies are difficult to calculate precisely, and although

existing data does span both sides of the ϕ meson it is usually a *relative* cross section useful for measurements of ϕ meson parameters. Hence a Breit-Wigner (BW) line shape with PDG 1994 parameters has been used, below the (most recent) lowest data point for kaon pair production and for the whole energy range for the radiative decays, where the errors (considered totally systematic) are evaluated by differentiating the BW formula.

In the charm and bottom threshold regions the 6 states each of the J/Ψ and Υ resonance families have been calculated separately using PDG 1994 values and the peak approximation formula for a_μ^{had} ([22] [11]). In view of the small contribution to a_μ^{had} from the b -quark (due to kernel function suppression in Eq. 1), the top quark contribution is neglected.

2. Higher Order VMD Contributions

The Vector Meson Dominance (VMD) model approximation of QCD expresses the fact that vector mesons, instead of quarks and gluons, are the relevant degrees of freedom in low energy (near threshold) QCD interactions. The VMD model approximation for hadronic vacuum polarization is depicted in Fig. 3. Most of the a_μ^{had} contributions are contained in the first term on the right of the figure. The decay modes of the vector mesons ρ , ω , ϕ and ω' account for $\sim 597 \times 10^{-10}$ (85% of a_μ^{had}), while ρ' accounts for $\sim 34.7 \times 10^{-10}$ (5% of a_μ^{had}), and ϕ' for $\sim 1.6 \times 10^{-10}$ by rough accounting. The remaining part ($\sim 10\%$) of a_μ^{had} is derived from a non- or very broadly resonant background for which it is instructive to consider some additional generation mechanisms not already included in the previous exclusive mode approach.

Namely, some portion of a_μ^{had} derives from higher order VMD interactions $V\pi$ and $V\pi\pi$ (the next terms on the right in Fig. 3) where V is a vector meson. Although these amounts are individually less than the expected E821 error ($\sim 4.0 \times 10^{-10}$) the fact that they may combine with other small contributions (whose sum may be greater than the E821 error) implies that they all should indeed be carefully considered.

The cross sections $\sigma(K^*K^\pm\pi^\mp)$, $\sigma(\omega\pi^0)$, $\sigma(\omega\pi^+\pi^-)$, $\sigma(\eta\pi^+\pi^-)$ have recently (1991,1992) been measured (ND,DM2). In particular, it has been pointed out [12] that the following exclusive hadronic modes marked with the comment “not” need to be included (B refers to the branching fraction for the decay mode in parentheses):

$$\begin{array}{llll} \omega\pi^0 \rightarrow (\pi^+\pi^-\pi^0)\pi^0 & B=0.888 & \text{is in } \pi^+\pi^-\pi^0\pi^0 \\ & \rightarrow (\pi^0\gamma)\pi^0 & B=0.085 & \text{not counted} \\ & \rightarrow (\pi^+\pi^-)\pi^0 & B=0.0221 & \text{is in } \pi^+\pi^-\pi^0 \\ \omega\pi^+\pi^- \rightarrow (\pi^+\pi^-\pi^0)\pi^+\pi^- & B=0.888 & \text{is in } \pi^+\pi^-\pi^+\pi^-\pi^0 \\ & \rightarrow (\pi^0\gamma)\pi^+\pi^- & B=0.085 & \text{not in } \pi^+\pi^-\pi^0 \\ & \rightarrow (\pi^+\pi^-)\pi^+\pi^- & B=0.0221 & \text{is in } \pi^+\pi^-\pi^+\pi^- \\ \omega\pi^0\pi^0 \rightarrow (\pi^+\pi^-\pi^0)\pi^0\pi^0 & B=0.888 & \text{is in } \pi^+\pi^-\pi^0\pi^0\pi^0 \\ & \rightarrow (\pi^0\gamma)\pi^0\pi^0 & B=0.085 & \text{not counted} \\ & \rightarrow (\pi^+\pi^-)\pi^0\pi^0 & B=0.0221 & \text{is in } \pi^+\pi^-\pi^0\pi^0. \end{array}$$

They further point out that isospin considerations [23] imply

$$\begin{aligned} 2\sigma(\pi^+\pi^-\pi^0\pi^0\pi^0) &= \sigma(\pi^+\pi^-\pi^+\pi^-\pi^0) \\ 2\sigma(\omega\pi^0\pi^0) &= \sigma(\omega\pi^+\pi^-) \end{aligned}$$

and that this must be used since $\sigma(\omega\pi^0\pi^0)$ and $\sigma(\pi^+\pi^-\pi^0\pi^0\pi^0)$ have not been measured. Therefore the $\sigma(\pi^+\pi^-\pi^+\pi^-\pi^0)$ contribution has been augmented by a factor $A_1 = 1.5$, and the contributions from $\sigma(\omega\pi^0)$ and $\sigma(\omega\pi^+\pi^-)$ will contribute with factors $B_1 = 0.085$ and $B_3 = 1.5B_1$ respectively, as noted in Tab. IV. However, the same logic applied to $\sigma(\eta\pi^+\pi^-)$ implies *not* including 100% of it as done previously [12]:

$$\begin{array}{llll} \eta\pi^+\pi^- \rightarrow (\gamma\gamma)\pi^+\pi^- & B=0.388 & \text{not in } \pi^+\pi^- \\ & \rightarrow (\pi^0\pi^0\pi^0)\pi^+\pi^- & B=0.319 & \text{is in } \pi^+\pi^-\pi^0\pi^0\pi^0 \\ & \rightarrow (\pi^+\pi^-\pi^0)\pi^+\pi^- & B=0.236 & \text{is in } \pi^+\pi^-\pi^+\pi^-\pi^0 \\ & \rightarrow (\pi^+\pi^-\gamma)\pi^+\pi^- & B=0.0488 & \text{not in } \pi^+\pi^-\pi^+\pi^-\pi^0. \end{array}$$

To avoid double counting in the $\pi^+\pi^-\pi^+\pi^-\pi^0$ and $\pi^+\pi^-\pi^0\pi^0\pi^0$ channels only the fraction $B_2 = 0.388 + 0.0488$ of the $\sigma(\eta\pi^+\pi^-)$ contribution has been included. (There is no further factor of 1/2 for inclusion of the cross section $\sigma(e^+e^- \rightarrow \eta\pi^0\pi^0)$ since it is forbidden.) The augmentation and branching factors (A_1 , B_1 , B_2 , B_3) are noted in Tab. IV where all contributions are listed in descending order.

D. The Perturbative QCD Energy Region

To test the validity of QCD for determination of the contribution to a_μ^{had} in the energy region above 3.1 GeV, a QCD parameterization [24] (including second order terms) was compared with a data-based evaluation [12]. An asymptotic kernel function [12] was used in the a_μ^{had} integral since Eq. 2 is numerically stable only up to ~ 20 GeV. The errors on the QCD contribution to a_μ^{had} were determined by its variation with $\Lambda_{\overline{MS}} \pm \Delta\Lambda_{\overline{MS}}$.

Marshall combined the results from 15 different e^+e^- annihilation experiments, fitting them to a third order QCD model with a single parameter, $\Lambda_{\overline{MS}}$. Because the fit was overconstrained, he was able to evaluate whether these experiments had overestimated their systematic errors. He then went a step further, and fitted for the absolute normalization of each of the 15 experiments independently, bounded by double their stated systematic errors ($\pm 2\sigma$). He found fitted normalizations for most ex-

periments within their stated limits, with the exception of two of the earliest experiments: Mark II and $\gamma\gamma 2$.

By using the error in his normalization constants (from the fit) rather than the stated systematic errors for each experiment, Marshall was able to significantly reduce his error on $\Lambda_{\overline{MS}}$ and hence on his overall normalization for $R(s)$. Dubnicka followed Marshall in his 1992 preprint [9], and in the present calculation both Marshall's fitted parameterization (with his errors) and trapezoidal integration [12] have been compared for the higher-energy contributions to a_μ^{had} . As the QCD and data-based results are in good agreement it is clear that the second order QCD expression is sufficient.

V. WFSA RESULTS AND PROJECTIONS

A. Summary Results

As the preceding discussion of the WFSA evaluation shows, the core a_μ^{had} problem is how to combine the correlations among the systematic errors. In the present calculation, partial a_μ^{had} integrations have been performed first so relative errors and error-weighted averaged systematic and statistical errors could be defined at an early stage of the error combination calculation. The averaged systematic errors have been correlated according to the postulates discussed in Sect. III A for all experiments measuring more than one hadronic mode; all other errors have been combined in quadrature only after such correlations have been taken into account.

All of the WFSA calculations have been collected in Tab. II. The first two lines of the table are summary results of the a_μ^{had} contributions and errors presented in Table III. In the > 2 hadrons line of Tab. III the results from Tab. IV are presented. The third line in Tab. II for $R(s)$ from 2.0 - 3.1 GeV is the summary result from Table V. All of the a_μ^{had} central values are averaged over detectors first and then summed over modes, while the errors are combined over modes first (to correlate systematics) and only combined over detectors in the end. The errors in Tab. II may be combined in quadrature since there are no correlations remaining among the categories chosen. (This feature is not completely present in the previous calculations of a_μ^{had} .)

A subtotal for a_μ^{had} contributions below 3.1 GeV is presented in Tab. II in order to show the total errors before the use of QCD. It is there seen that the QCD results do not have much influence on the final results since the error from the lower energy (< 3.1 GeV) region dominates. For comparison, if in place of the QCD calculation the data based evaluation for the region above 3.1 GeV [12] is used the final errors are only slightly higher: $a_\mu^{had} = 702.718(7.787)(14.147)$. Hence, the WFSA result is thus shown to be largely insensitive to the use of QCD or the experimental data after 3.1 GeV. (This justifies both the use of QCD down to the energy 3.1 GeV and

the neglect of third order QCD terms; as new data bring the errors from the lower energy regions downward (see Sect. V), the QCD error will become more significant however.)

It is further apparent from Table II that the error in the energy region below 1.4 GeV dominates the total error in a_μ^{had} , and this error comes mostly from the Olya $\pi^+\pi^-$ data. (One advantage of the WFSA method is that in presenting averaged errors (both statistical and systematic separately) for each detector and exclusive mode, it remains possible to identify which detector and exclusive mode is contributing the most to the overall total error.) The WFSA method most heavily weights the data with the smallest errors (see Table IX), and hence most of the stated systematic errors are indeed being added linearly for the experiments with the smallest errors.

B. Energy Region < 1.4 GeV

The overall WFSA results indicate that improved measurements below 1.4 GeV, can significantly improve upon the overall lowest order hadronic vacuum polarization uncertainty in a_μ^{had} . To show this, the WFSA procedure has been performed with an additional two experiments in view, the CMD2 experiment at VEPP-2M in Novosibirsk, Russia and a hypothetical detector at DAFNE at Frascati, Italy. The a_μ^{had} results for the energy region below 1.4 GeV are shown in Tab. VI for three cases: a) without new data, b) with expected errors (stat)(sys) = (3%)(0.5%) from the CMD2 experiment and c) CMD2 plus a second experiment with errors (stat)(sys) = (3%)(0.5%). In particular, the central values for hadronic cross sections have been chosen equal to CMD or Olya values (and hence the S factors will generally be less than one and so not applied), while the errors above have been determined for only the following exclusive hadronic modes: $\pi^+\pi^-$ (2 π), $\pi^+\pi^-\pi^0$ (3 π), $\pi^+\pi^-\pi^+\pi^-$ (4 π), $\pi^+\pi^-\pi^0\pi^0$ (4 π), $\pi^+\pi^-\pi^+\pi^-\pi^0$ (5 π), K^+K^- and K_LK_S .

The choice of 3 % statistical error is based on the CMD2 data taking assumption of 1000 pion pairs per energy point since $\pi^+\pi^-$ is the dominant contribution to a_μ^{had} . Since the luminosity collected by CMD2 has been determined by this requirement on the $\pi^+\pi^-$ cross section (and by competing requirements for other vector meson physics goals), the actual statistical errors on the non- $\pi^+\pi^-$ higher multiplicity modes will in fact be somewhat different, but this is a higher order effect here neglected.

The choice of 0.5 % systematic error is based on the fact that the limiting error on the new CMD2 cross section measurements appears to be the error on higher order corrections to the QED Bhabha cross sections used in calculations of the luminosity. While radiative corrections have been calculated to good accuracy (0.11 % [25]) for the t -channel contributions to Bhabha scattering use-

ful for the forward region luminosity monitors in use at LEP, they are not useful for CMD2; the principle luminosity is determined there by large angle Bhabha events in the barrel calorimeter where significant t - and s -channel interference terms are present. The task of calculating these interference terms to 0.5 % accuracy is well underway [26] and it is assumed that this will be the limiting error on the new CMD2 cross section measurements.

The results in Tab. VI show that without new data the errors on the contributions to $10^{10}a_\mu^{had}$ below 1.4 GeV will be $10^{10}(\text{stat})(\text{sys}) = (7.4)(13.1)$ and with particular new data they can become $10^{10}(\text{stat})(\text{sys}) = (2.1)(2.9)$ which is equivalent to a total error of 3.6×10^{-10} . Comparing this with the AGS E821 experimental error goal of $\sigma_{exp}^{a_\mu} \sim \pm 4.0 \times 10^{-10}$ it is clear that the new data is needed. In particular, if this data has errors of 3 % statistical and 0.5 % systematic on the stated modes, then they will in fact be sufficient to reduce the hadronic contribution errors in the energy range below 1.4 GeV to below the error goal of the new measurement of a_μ^{had} by the BNL AGS E821 Collaboration.

C. Energy Region > 1.4 GeV

However, the new data from BINP Novosibirsk and INFN Frascati will not reduce the errors on a_μ^{had} in the energy range above 1.4 GeV. The current error on a_μ^{had} obtained by use of the WFSA procedure in the region above 1.4 GeV is $10^{10}(\text{stat})(\text{sys}) = (2.4)(5.3)$. In this region, most of the error comes from the $\gamma\gamma 2$ detector at the former 3 GeV Adone storage ring in INFN Frascati, Italy. As this error is larger than the BNL AGS E821 experimental error goal of $\sigma_{exp}^{a_\mu} \sim \pm 4.0 \times 10^{-10}$, clearly more experiments are needed for interpretation of the new measurement of a_μ^{had} . Fortunately, it is possible to use new high statistics data on τ -decays for new measurements of multi-pion production in the energy region 1.4 - 2.0 GeV, and above 2.2 GeV there are plans to make further measurements of the cross section ratio $R(s)$ at BEPC Beijing, China [27].

1. τ Decay Data for Energy Region 1.4 - 2.0 GeV

The new high statistics data on τ -decays has already been included in the present calculation by use of the $\omega\pi^0$ cross section obtained from the ARGUS Collaboration [28] [29]. The idea is based on the fact that the coupling of $W^\pm$ bosons to quarks is related to the photon coupling to quarks by an isospin rotation (CVC relation) [30]. The effect on the total errors in the energy region 1.4 - 2.0 GeV are presented in Tab. VII where the assumed errors are $(\text{stat})(\text{sys}) = (3\%)(1\%)$ for the following hadronic modes: $\pi^+\pi^-\pi^+\pi^-$ (4 π charged), $\pi^+\pi^-\pi^0\pi^0$ (4 π neutral), $\pi^+\pi^-\pi^+\pi^-\pi^0$ (5 π) and $\pi^+\pi^-\pi^+\pi^-\pi^0\pi^0$ (6 π).

New data on the 4 π neutral, 5 π and 6 π modes can reduce the $\gamma\gamma 2$ contribution to the total errors (as shown in Row 2) while new data on the 4 π charged mode is required to reduce the DM1 and DM2 contribution to the total errors (as shown in Row 3). Therefore, if many data points ($\sim 10 - 20$) for each of the specified modes across the energy region 1.4 to 2.0 GeV with the assumed errors can be extracted from τ -decay data, then the contribution to the total errors from this energy region will be reduced significantly as shown in Tab. VII.

2. New $R(s)$ Measurements for Energy Region 2.0 - 3.1 GeV

The effect of new data on the error in the 2.0 - 3.1 GeV energy region is presented in Tab. VIII. The WFSA procedure has been performed with a new hypothetical detector measuring $R(s)$ from 2.2 - 3.1 GeV (central value of $R(s) = 3.0$ for all points so no S -factors exceed 1) with assumed errors of $(\text{stat})(\text{sys}) = (3\%)(0.5\%)$. If new $R(s)$ data in this energy region can be obtained with these errors then the $\gamma\gamma 2$ contribution to the total error can be significantly reduced.

D. Conclusion

Taking the new hypothetical data together (including τ decays and $R(s)$ measurements), the total error from the energy range above 1.4 GeV (including resonances and QCD) then becomes $10^{10}(\text{stat})(\text{sys}) = (0.6)(1.6)$. Combining the hypothetical results from above and below 1.4 GeV the total error becomes $10^{10}(\text{stat})(\text{sys}) = (2.2)(3.3)$ which is equivalent to a total error of 3.98×10^{-10} . Therefore, new measurements of $e^+e^- \rightarrow \text{hadrons}$ (including the cross section ratio $R(s)$) and τ decays with assumed errors of $(\text{stat})(\text{sys}) = (3\%)(0.5\%)$ and $(3\%)(1.0\%)$ respectively are sufficient to reduce the total error on the lowest order contribution to a_μ^{had} in all energy regions below the expected error of the new measurement of a_μ by the AGS E821 Collaboration.

Acknowledgments

The authors would like to acknowledge useful discussions with S.I. Eidelman, I. Grosse, T. Kinoshita, W.J. Marciano and V.G. Zavarzin in various aspects of this work. We would also like to thank B.I. Khazin, E.P. Solodov and V.G. Zavarzin for discussions about recent and expected CMD2 results. Comments on previous drafts by R.M. Carey, J.P. Miller and B.L. Roberts are also appreciated. Finally, a very special thanks is extended to V.A. Monich for help improving the WFSA calculation routines.

APPENDIX A: EXPERIMENTAL DATA USED

The experimental data we used for our evaluation of a_μ^{had} is listed by hadronic exclusive mode in Tab. XIII.

-
- [1] B.L. Roberts. The New Muon $g - 2$ Experiment. *Zeitschrift für Physik*, 56:S101–S108, 1992.
 - [2] V.W. Hughes. . In T. Hasegawa et al., editors, *Frontiers of High Energy Spin Physics*, pages 717–722, Tokyo, 1992. Universal Academy Press.
 - [3] Jorge L. Lopez, D.V. Nanopoulos, and Xu Wang. Large $(g - 2)_\mu$ in $SU(5) \times U(1)$ supergravity models. *Physical Review D*, 49:366, 1994.
 - [4] U. Chattopadhyay and Pran Nath. Probing Supergravity Grand Unification in the Brookhaven $g - 2$ Experiment. Nsf-itp-95-64; nub-th-3125/95; los alamos e-print archive: hep-ph, Physics Department Northeastern University, Boston, MA, 1995.
 - [5] P. Mery et al. Constraints on non-standard effects from present and future $g_\mu - 2$ measurements. *Zeitschrift für Physik C*, 46:229–252, 1990.
 - [6] T. Kinoshita. . *Physical Review Letters*, 75:4728, 1995.
 - [7] Andrzej Czarnecki, Bernd Krause, and William J. Marciano. Electroweak Fermion-loop Contributions to the Muon Anomalous Magnetic Moment. Los alamos e-print archive: hep-ph/9506256, Institut für Theoretische Teilchenphysik, Karlsruhe and BNL, 1995.
 - [8] Lev Mitrofanovich Barkov et al. (Olya and CMD) Electromagnetic pion form factor in the timelike region. *Nuclear Physics B*, 256:365–384, 1985.
 - [9] A.Z. Dubničková, S. Dubnička, and P. Stríženec. New evaluation of hadronic contributions to the anomalous magnetic moment of charged leptons. Technical Report E2-92-281, Dubna Institute of Nuclear Research, 1992.
 - [10] K. Adel and F.J.Ynduráin. Improved Evaluation of the Hadronic Vacuum Polarization Contributions to Muon $g - 2$ and $\alpha_{\overline{QED}}(M_Z)$ Using High Order QCD Calculations. Ftum 95-32; los alamos e-print archive: hep-ph 9510, Departamento de Fisica Teorica, Universidad Autonoma de Madrid, 1995.
 - [11] T. Kinoshita, B. Nižić, and Y. Okamoto. Hadronic contributions to the anomalous magnetic moment of the muon. *Physical Review D*, 31:2108, 1985.
 - [12] S. Eidelman and F. Jegerlehner. Hadronic contributions to $g - 2$ of the leptons and to the effective fine structure constant $\alpha(M_Z^2)$. *Zeitschrift für Physik C*, 67:585–602, 1995.
 - [13] M. Hayakawa, T. Kinoshita, and A.I. Sanda. Hadronic Light-by-light Scattering Effect on Muon $g - 2$. *Physical Review Letters*, 75:790, 1995.
 - [14] J. Bijnens, E. Pallante, and J. Prades. Hadronic light-by-light contributions to the muon $g - 2$ in the large N_c limit. Los alamos e-print archive: hep-ph/9511388, march 1996, NORDITA and Niels Bohr Institute, Copenhagen, Denmark, 1996.
 - [15] B.E. Lautrup, A. Peterman, and E. de Rafael. Recent developments in the comparison between theory and experiments in quantum electrodynamics. *Physics Reports*, 3:193–260, 1972.
 - [16] D.H. Brown and W.A. Worstell. Update on the Hadronic Contribution to the Muon $g - 2$ value. Technical report, Boston University Physics Department, 1995. BNL AGS E821 $g - 2$ Note # 220.
 - [17] Particle Data Group. Review of Particle Properties. *Physical Review*, 50:1173–1826, 1994.
 - [18] C. Bacci et al. $(\gamma\gamma 2)$ Total cross section for hadronic production by e^+e^- annihilation in the total center of mass energy range 1.42 - 3.09. *Physics Letters B*, 86:234–238, 1979.
 - [19] C. Bacci et al. $(\gamma\gamma 2)$ Measurement of Hadronic Exclusive Cross Sections in e^+e^- Annihilation from 1.42 to 2.20 GeV. *Nuclear Physics B*, 184:31–39, 1981.
 - [20] J.L. Siegrist et al. (Mark I) Hadron production by e^+e^- annihilation at center of mass energies between 2.6 and 7.8 GeV. *Physical Review D*, 26:969, 1982.
 - [21] M. Bernardini et al. (BCF) The Energy Dependence of $\sigma(e^+e^- \rightarrow \text{hadrons})$ in the Total Center-of-Mass Energy Range 1.2 to 3.0 GeV. *Physics Letters B*, 51:200, 1974.
 - [22] V. Barger, W.F. Long, and M.G. Olsson. New evaluation of muon $g - 2$ hadronic anomaly. *Physics Letters B*, 60:89, 1975.
 - [23] A. Pais. The Many π meson Problem. *Annals of Physics (NY)*, 9:548, 1960.
 - [24] R. Marshall. A determination of the strong coupling constant α_s from e^+e^- total cross section data. *Zeitschrift für Physik C*, 43:595–606, 1989.
 - [25] A. Arbuzov et al. The present theoretical error on the Bhabha scattering cross section in the luminosity region at LEP. Los alamos e-print archive: hep-ph 9605239, Dubna, CERN, DESY, INFN, SLAC, Poland,..., 1996.
 - [26] B.I. Khazin and E.P. Solodov, 1996. private communication.
 - [27] B. L. Roberts, 1996. private communication.
 - [28] N. Albrecht et al. (ARGUS). *Physics Letters B*, 185:223, 1987.
 - [29] S.I. Dolinsky et al. (ND) Summary of Experiments with the Neutral Detector at the e^+e^- Storage Ring VEPP-2M. *Physics Reports*, 202:99, 1991.
 - [30] F.J. Gilman and S.H. Rhie. . *Physical Review D*, 31:1066, 1985.
 - [31] L.M. Barkov et al. (Olya) Study of the Reaction $e^+e^- \rightarrow \pi^+\pi^-$ in the Energy Range from 640 to 1400 MeV. *Soviet Journal of Nuclear Physics*, 40:286, 1994.
 - [32] A. Quenzer et al. (DM1) Pion Form Factor from 480 MeV to 1100 MeV. *Physics Letters B*, 76:512, 1978.
 - [33] D. Bisello et al. (DM2) The Pion Electromagnetic Form Factor in the Time-Like Energy Range $1.35 \leq \sqrt{s} \leq 2.4$ GeV. *Physics Letters B*, 220:321, 1989.
 - [34] G. Cosme et al. (M2N) Measurement of the electron-positron annihilation cross section into $\pi^+\pi^-$ at the energies 915, 990 and 1076 MeV. Preprint LAL 1287, Laboratoire de L’Accelérateur Lineaire, Orsay, France, 1976.
 - [35] D. Bellini et al. (BCF) The Pion Electromagnetic Form Factor in the Timelike Range (1.44 - 9.0) (GeV)². *Lettre*

- al Nuovo Cimento*, 14:418, 1975.
- [36] G. Barbiellini et al. ($\mu\pi$) Hadron Pair Production by Electron-Positron Colliding Beams. *Lettere al Nuovo Cimento*, 6:557, 1973.
 - [37] B. Esposito et al. (MEA) Momentum Analysis of Kaon and Pion Pairs Produced from Time-Like Photons at 1.6 GeV Energy. *Physics Letters B*, 67:239, 1977.
 - [38] S.R. Amendolia et al. (NA7) Measurement of the Pion Form Factor in the Time-Like Region for q^2 Values between 0.1 (GeV/c)^2 and 0.18 (GeV/c)^2 . *Physics Letters B*, 138:454, 1984.
 - [39] I.B. Vasserman et al. (TOF) Measurement of the Pion Form Factor in the Reaction $e^+e^- \rightarrow \pi^+\pi^-$ in the Energy Range 0.40-0.46 GeV. *Soviet Journal of Nuclear Physics*, 33:709, 1981.
 - [40] Alexander Shekhtman. (CMD) *The Measurement of ω meson parameters on VEPP-2M with the Cryogenic Magnetic Detector*. PhD thesis, Budker Institute of Nuclear Physics, Novosibirsk, Russia, 1987. Siberian Division of Russian Academy of Sciences.
 - [41] L. M. Barkov et al. (CMD) Researches on the process $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ in the Energy Region .840 - 1.020 MeV at VEPP-2M with the Cryogenic Magnetic Detector. Preprint 89-15, Budker Institute of Nuclear Physics, Novosibirsk, Russia, 1989. in Russian.
 - [42] A. Cordier et al. (DM1) Cross Section of the Reaction $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ for C.M. Energies from 750 to 1100 MeV. *Nuclear Physics B*, 172:13, 1980.
 - [43] A. Antonelli et al. (DM2) Measurement of the $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ and $e^+e^- \rightarrow \omega\pi\pi$ Reactions in the Energy Interval 1350-2400 MeV. *Zeitschrift fur Physik C*, 56:15, 1992.
 - [44] L. M. Kurdadze et al. (Olya) The Results of Experiments with the Olya Detector at the VEPP-2M Storage Ring. Preprint 82-97, Budker Institute of Nuclear Physics, Novosibirsk, Russia, 1982.
 - [45] G.V. Anikin et al. (CMD) The Results of Experiments with CMD at the VEPP-2M Storage Ring. Preprint 83-85, Budker Institute of Nuclear Physics, Novosibirsk, Russia, 1983.
 - [46] D. Bisello et al. (DM2) Study of the Reaction $e^+e^- \rightarrow K^+K^-$ in the Energy Range $1350 \leq \sqrt{s} \leq 2400$ MeV. *Zeitschrift fur Physik C*, 39:13, 1988.
 - [47] F. Mane et al. (DM1) Study of the Reaction $e^+e^- \rightarrow K_L K_S$ in the Total Energy Range 1.4-2.18 GeV and Interpretation of the K^+ and K^0 Form Factors. *Physics Letters B*, 99:261, 1981.
 - [48] L. M. Kurdadze et al. (Olya) Production of Four Pions in e^+e^- Collisions Below 1.4 GeV. Preprint 79-69, Budker Institute of Nuclear Physics, Novosibirsk, Russia, 1979.
 - [49] L. M. Barkov et al. (CMD) Study of Multiple Pion Production Reactions at the VEPP-2M Storage Ring Using a Cryogenic Magnetic Detector. *Soviet Journal of Nuclear Physics*, 47:248, 1988.
 - [50] A. Cordier et al. (DM1) Study of the $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$ Reaction in the 1.4-2.18 GeV Energy Range. *Physics Letter B*, 109:129, 1982.
 - [51] D. Bisello et al. (DM2) DM2 Results on e^+e^- Annihilation into Multihadrons in the 1350 - 2400 MeV Energy Range. Preprint LAL 90-35, INFN Padua and Frascati, Italy and LAL Orsay, France, 1990.
 - [52] L. Kurdadze et al. (Olya) Study of the Reaction $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ at 2E up to 1.4 GeV. *JETP Letters*, 43:643, 1986.
 - [53] A. Cordier et al. (DM1) Observation of a new Isoscalar Vector Meson in $e^+e^- \rightarrow \omega\pi^+\pi^-$ annihilation at 1.65 GeV. *Physics Letters B*, 106:155, 1981.
 - [54] D. Bisello et al. (DM1) Study of the Reaction $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^+\pi^-$ in the Total Energy Range 1400-2180 MeV. *Physics Letters B*, 107:145, 1981.
 - [55] F. Mane et al. (DM1) Study of $e^+e^- \rightarrow K_S K^\pm \pi^\mp$ in the 1.4-2.18 GeV Energy Range: A New Observation of an isoscalar Vector Meson: ϕ' (1.65 GeV). *Physics Letters B*, 112:178, 1982.
 - [56] A. Cordier et al. (DM1) Study of the Reaction $e^+e^- \rightarrow K^+K^-\pi^+\pi^-$ Reaction from 1.4 to 2.18 GeV. *Physics Letters B*, 110:335, 1982.
 - [57] A. Antonelli et al. (DM2) Measurement of the Reaction $e^+e^- \rightarrow \eta\pi\pi$ in the Center of Mass Energy Interval 1350-2400 MeV. *Physics Letters B*, 212:33, 1988.
 - [58] A. Cordier et al. (DM1) Observation of a New Isoscalar Vector Meson in $e^+e^- \rightarrow \omega\pi\pi$ Annihilation at 1.65 GeV. *Physics Letters B*, 106:155, 1981.
 - [59] A. Antonelli et al. (DM2) Measurement of the $e^+e^- \rightarrow K^+K^-$ and $e^+e^- \rightarrow \omega\pi\pi$ Reactions in the Energy Interval 1350-2400 MeV. *Zeitschrift fur Physik C*, 56:15, 1992.
 - [60] B. Delcourt et al. (DM1) Study of the Reaction $e^+e^- \rightarrow p\bar{p}$ in the Total Energy Range 1925-2180 MeV. *Physics Letters B*, 86:395, 1979.
 - [61] D. Bisello et al. (DM2) Baryon Pairs Production in e^+e^- annihilation at $\sqrt{s} = 2.4$ GeV. *Nuclear Physics B*, 224:379, 1983.

TABLE I. History of a_μ^{had} calculations. The values 4 and 15×10^{-10} are the AGS E821 error goal and electroweak contribution to a_μ respectively.

Calculations based on Experimental Data			
Author(s)	Year	$10^{10} a_\mu^{had}(stat)(sys)$	ref
Budker Institute	1985	684(11)	[8]
Kinoshita, Nizic, Okamoto	1985	707(6)(17)	[11]
Dubnickova, Dubnicka, Stricinec	1992	699(4)(2) ^a	[9]
Eidelman, Jegerlehner	1995	702(6)(14)	[12]
Apel, Yndurain	1995	710(11)	[10]
WFSA evaluation	1996	703(8)(14)	

^a Improvement due mostly to fitting function technique. See text.

TABLE II. The Grand Total WFSA $10^{10} a_\mu^{had}$ results

Energy Region [GeV]	WFSA $10^{10} a_\mu^{had} (stat) (sys)$
$\sigma(e^+e^- \rightarrow hadrons) E_\theta - 1.4$	611.332 (7.399) (13.045)
$\sigma(e^+e^- \rightarrow hadrons) 1.4 - 2.0$	32.466 (0.756) (2.379)
R(s) 2.0 - 3.1	28.374 (2.288) (4.400) ^a
J/Ψ (6 states)	9.047 (-) (0.969)
Υ (6 states)	0.109 (-) (0.013)
QCD 3.1 - ∞	21.301 (-) (0.371) ^b
Sub Total < 3.1 + $J/\Psi, \Upsilon$	681.328 (7.782) (14.005)
Sub Total < 1.4	611.332 (7.399) (13.045)
Sub Total > 1.4	91.297 (2.410) (5.108)
Total Figure 1	702.629 (7.782) (14.009)

E_θ refers to the particular thresholds of the exclusive hadronic modes.

^a Represents systematic errors mainly from $\gamma\gamma 2$ detector added linearly for energy regions 1.4 - 2.0 and 2.0 - 3.1 GeV (1.7 + 2.7).

^b Errors determined by $a_\mu^{QCD} (\Lambda_{\overline{MS}} \pm \Delta\Lambda_{\overline{MS}})$

TABLE III. Our WFSA $10^{10} a_\mu^{had}$ results below 2.0 GeV. Upper (lower) contributions listed in the total column are for below (above) 1.4 GeV. Upper (lower) numbers in parentheses are contributions to the statistical (systematic) errors. Systematic (statistical) errors are combined linearly (in quadrature) in each column separately.

Mode	CMD	Olya	ND	DM1A	DM1D	DM2	Other	Total
$\pi^+\pi^-$	- (4.806) (4.789)	- (1.306) (5.807)		- (5.064) (4.021)		- (0.058) (0.110)	- (0.814) (0.732)	502.184 0.781
$\pi^+\pi^-\pi^0$	- (0.509) (0.104)		- (0.856) (1.303)	- (1.209) (0.715)		- (0.041) (0.053)	- (0.049) (0.009)	50.886 0.677
> 2 hadrons	- (0.236) (0.589)	- (0.369) (2.405)	- (0.205) (2.070)		- (0.241) (1.071)	- (0.307) (1.857)	- (0.631) (1.735)	16.782 29.995
K^+K^-	- (0.119) (0.058)	- (0.953) (1.900)				- (0.046) (0.090)	(0.230) ^b	20.623 4.45 ^a 0.759
$K_L K_S$	- (0.173) (0.040)	- (0.156) (0.101)			- (0.033) (0.017)		(0.586) ^b	0.755 14.07 ^a 0.154
$p\bar{p}$					- (0.020) (0.010)			0.100
$\omega \rightarrow \pi^0 \gamma, \eta \gamma$							(0.040) ^b	0.980 ^a
$\phi \rightarrow \pi^0 \gamma, \eta \gamma$							(0.024) ^b	0.602 ^a
Total $E_\theta - 1.4$	- (4.842) (5.580)	- (1.666) (10.213)	- (0.880) (3.373)	(5.206) (4.736)			(0.807) (0.962)	611.33 (7.399) (13.045)
Total 1.4 - 2.0					- (0.244) (1.098)	- (0.318) (2.110)	- (0.642) (1.747) ^c	32.47 (0.756) (2.379)

E_θ refers to the particular thresholds of the exclusive hadronic modes.

DM1A = DM1 at ACO; DM1D = DM1 at DCI, Orsay.

^a Integration of energy dependent width Breit-Wigner in absence of data.

^b Errors determined by m_V , Γ_V , $B_{V \rightarrow ee}$ derivatives of the Breit-Wigner.

^c The $\gamma\gamma 2$ systematic errors from energy regions 1.4 - 2.0 GeV (1.7) and 2.0 - 3.1 GeV (2.7) will be added linearly and presented in Table II.

TABLE IV. Our WFSa $10^{10}a_{\mu}^{had}$ results for > 2 hadrons exclusive modes below 2.0 GeV. Upper (lower) contributions listed in the total column are for below (above) 1.4 GeV. Upper (lower) numbers in parentheses are contributions to the statistical (systematic) errors. Systematic (statistical) errors are combined linearly (in quadrature) in each column separately.

Mode	CMD	Olya	ND	DM1D	DM2	Other	Total	S Factor
$\pi^+\pi^-\pi^0\pi^0$		(0.325) (1.491)	(0.193) (1.514)		(0.256) (0.868)	(0.335) (0.841)	10.783 8.543	1.614 1.692
$\pi^+\pi^-\pi^+\pi^-$	(0.225) (0.527)	(0.174) (0.914)	(0.066) (0.526)	(0.104) (0.623)	(0.070) (0.628)		5.161 10.215	0.486 1.221
$\pi^+\pi^-\pi^+\pi^-\pi^0\pi^0$						(0.494) (0.763)	5.089	0.0
$A_1(\pi^+\pi^-\pi^+\pi^-\pi^0)$	(0.072) (0.062)			(0.124) (0.192)		(0.187) (0.131)	0.305 2.533	0.0 1.010
$K_S K^\pm \pi^\mp$				(0.158) (0.0)	(0.104) (0.119)		0.951	3.199
$K^+ K^- \pi^+ \pi^-$				(0.072) (0.135)	(0.091) (0.123)		0.815	2.709
$K^* K^\pm \pi^\mp$					(0.057) (0.069)		0.692	0.0
$B_1(\omega\pi^0)$			(0.017) (0.030)			(0.079) (0.021)	0.533 0.210	1.132 0.0
$B_2(\eta\pi^+\pi^-)$					(0.039) (0.044)		0.444	0.0
$\pi^+\pi^-\pi^+\pi^-\pi^+\pi^-$				(0.040) (0.117)			0.419	0.0
$B_3(\omega\pi^+\pi^-)$				(0.005) (0.003)	(0.003) (0.004)		0.084	0.982
Total	- (0.236) (0.589)	- (0.369) (2.405)	- (0.205) (2.070)	- (0.241) (1.071)	- (0.307) (1.857)	- (0.631) (1.735)	16.782 29.995 -	

TABLE V. Our WFSa $10^{10}a_{\mu}^{had}$ results for $R(s)$ from 2.0 - 3.1 GeV. Error weighted averages with S -factor in “Total” column. Numbers in upper (lower) parenthesis are WFSa contributions to the statistical (systematic) errors which are combined in quadrature (linearly) in each column separately. Lower right corner errors are quadrature sums of errors in the “Total” row.

Energy Range [GV]	BCF	$\gamma\gamma 2$	Mark I	Total	S Factor
2.0 - 2.6	18.778 (2.149) (0.207)	22.441 (0.527) (2.093)		20.480	0.637
2.6 - 2.87	3.718 (0.469) (0.032)	6.535 (0.130) (0.451)	4.781 (0.236) (0.406)	4.898	1.210
2.87 - 3.0	1.721 (0.202) (0.019)		1.777 (0.061) (0.193)	1.742	0.096
3.0 - 3.1			1.254 (0.062) (0.313)	1.253	0.0
Total	- (2.208) (0.258)	- (0.543) (2.544)	- (0.251) (0.913)	28.374 (2.288) (2.700)	

TABLE VI. WFSA $10^{10}a_{\mu}^{had}$ detector total statistical (upper) and systematic (lower) errors below 1.4 GeV: Row 1) with no new data, Row 2) one new experiment with 3 % stat., 0.5 % syst. errors, Row 3) two new experiments with 3 % stat. and 0.5 % syst. errors.

WFSA $10^{10}a_{\mu}^{had}$ Errors Below 1.4 GeV							
New 2	New 1	CMD	Olya	ND	DM1A	Other	Total
		(4.842)	(1.666)	(0.880)	(5.206)	(0.807)	(7.399)
		(5.580)	(10.213)	(3.373)	(4.736)	(0.962)	(13.045)
	(2.426)	(1.266)	(1.148)	(0.136)	(1.239)	(0.451)	(3.250)
	(2.224)	(1.292)	(3.608)	(0.405)	(1.241)	(0.760)	(4.681)
(1.183)	(1.183)	(0.789)	(0.528)	(0.078)	(0.766)	(0.361)	(2.103)
(1.256)	(1.256)	(0.773)	(1.921)	(0.244)	(0.769)	(0.718)	(2.934)

TABLE VII. WFSA $10^{10}a_{\mu}^{had}$ detector total statistical (upper) and systematic (lower) errors in region 1.4 - 2.0 GeV. Row 1) with no new data, Row 2) new hypothetical data (possibly from τ decay spectral functions) on modes $\pi^+\pi^-\pi^0\pi^0$, $\pi^+\pi^-\pi^+\pi^-\pi^0$ and $\pi^+\pi^-\pi^+\pi^-\pi^0\pi^0$ with 3 % stat., 1.0 % syst. errors, Row 3) including as well the mode $\pi^+\pi^-\pi^+\pi^-$.

WFSA $10^{10}a_{\mu}^{had}$ Errors in 1.4 - 2.0 GeV Region				
New 1	DM1D	DM2	$\gamma\gamma 2$	Total
	(0.244)	(0.318)	(0.628)	(0.745)
	(1.098)	(2.110)	(1.744)	(2.950)
(0.068)	(0.210)	(0.192)	(0.067)	(0.300)
(0.115)	(0.922)	(1.320)	(0.141)	(1.620)
(0.082)	(0.179)	(0.157)	(0.067)	(0.261)
(0.175)	(0.346)	(0.514)	(0.141)	(0.659)

TABLE VIII. WFSA $10^{10}a_{\mu}^{had}$ detector total statistical (upper) and systematic (lower) errors in region 2.0 - 3.1 GeV. Row 1) with no new data, Row 2) one new experiment above 2.2 GeV with 3 % stat., 0.5 % syst. errors

WFSA $10^{10}a_{\mu}^{had}$ Errors in 2.0 - 3.1 GeV Region				
New 1	BCF	$\gamma\gamma 2$	Mark I	Total
	(2.208)	(0.543)	(0.251)	(2.288)
	(0.258)	(2.544)	(0.913)	(2.715)
(0.109)	(0.485)	(0.140)	(0.025)	(0.517)
(0.058)	(0.109)	(0.601)	(0.084)	(0.619)

TABLE IX. Systematic Errors used in our evaluation of the $\pi^+\pi^-$ contribution to a_μ^{had} .

Detector	Systematic error
NA7	.01,.02,.025,.05
CMD	.02
TOF	.035 ^a
DM1	.022
Oly	.04 - .15
M2N	.0 ^b
BCF	.1 ^c
$\mu\pi$	.1 ^c
MEA	.1 ^c
DM2	.12

^a evaluated from errors on radiative correction/efficiency factors.

^b systematic error is included with statistical in quoted errors.

^c Arbitrary value - inconsequential since statistical errors are large.

TABLE X. Systematic Errors used in our evaluation of the $\pi^+\pi^-\pi^0$ contribution to a_μ^{had} .

Detector	Energy Range [GeV]	Systematic error
CMD	.76 - .81	.0 ^a
CMD	.84 - 1.013	.07
ND	.414 - .765	.1
ND	.805 - 1.003	.1
ND	1.036 - 1.379	.2
DM1	0.414 - 1.098	.032
DM2	1.34 - 2.0	.0866
$\gamma\gamma$	1.437 - 2.0	.15

^a systematic errors are less than quoted statistical errors.

TABLE XI. Systematic errors used in our evaluation of higher multiplicity exclusive hadronic mode a_μ^{had} contribution.

Mode	Oly	CMD	ND	ARGUS	DM1	DM2	$\gamma\gamma 2$
$K^+ K^-$	0.1 ^a	0.038	0.1			0.1 ^a	
$K_L K_S$	0.3	0.1	0.1		0.1 ^a		
$\pi^+ \pi^- \pi^+ \pi^-$	0.15	0.1	0.1		0.1	0.1 ^a	
$\pi^+ \pi^- \pi^0 \pi^0$	0.15		0.15			0.1	0.15
$\pi^+ \pi^- \pi^+ \pi^- \pi^0$		0.20			0.12		0.15
$K^+ K^- \pi^+ \pi^-$					0.13	0.1 ^a	
$K_S K^\pm \pi^\mp$					0.0	0.1 ^a	
$K^* K^\pm \pi^\mp$						0.1 ^a	
$\pi^+ \pi^- \pi^+ \pi^- \pi^+ \pi^-$					0.28		
$\pi^+ \pi^- \pi^+ \pi^- \pi^0 \pi^0$							0.15
$p\bar{p}$					0.1 ^a		
$\omega\pi$			0.1	0.1			
$\omega\pi\pi$					0.12	0.082	
$\eta\pi\pi$						0.1 ^a	

^a systematic error not discussed in reference

TABLE XII. Systematic errors used in evaluation of the energy region 2.0 - 3.1 GeV a_μ^{had} contribution.

Energy Region [GeV]	BCF	$\gamma\gamma 2$	Mark I
2.0 - 3.1	0.02	0.21	0.25

TABLE XIII. Data used in the WFSA evaluation

Mode	OLYA	CMD	ND	DM1	DM2	Others
$\pi^+\pi^-$	[31]	[8]		[32]	[33]	M2N BCF $\mu\pi$ MEA NA7 TOF [34] [35] [36] [37] [38] [39]
$\pi^+\pi^-\pi^0$		[40] [41]	[29]	[42]	[43]	GG281 [19]
K^+K^-	[44]	[45]			[46]	MEA [37]
$K_L K_S$	[44]	[45]		[47]		
$\pi^+\pi^-\pi^+\pi^-$	[48]	[49]	[29]	[50]	[51]	
$\pi^+\pi^-\pi^0\pi^0$	[52]		[29]		[46]	GG281 [19]
$\pi^+\pi^-\pi^+\pi^-\pi^0$		[45]		[53]		GG281 [19]
$\pi^+\pi^-\pi^+\pi^-\pi^+\pi^-$				[54]		
$\pi^+\pi^-\pi^+\pi^-\pi^0\pi^0$						GG281 [19]
$K_S K^\pm \pi^\mp$				[55]	[51]	
$K^+ K^- \pi^+ \pi^-$				[56]	[51]	
$K^* K^\pm \pi^\mp$					[51]	
$\omega\pi^0$			[29]			ARGUS [28]
$\eta\pi^+\pi^-$					[57]	
$\omega\pi\pi$				[58]	[59]	
$p\bar{p}$				[60]	[61]	
$R(s)$ 2.0 - 3.1 GeV						GG279 GG281 BCF74 MarkI [18] [19] [21] [20]




FIG. 1. Lowest order hadronic contribution to a_μ^{had} .




FIG. 2. Higher order hadronic contributions to a_μ^{had}

$$\begin{array}{c}
 \rho, \omega, \phi, \rho \dots \quad \omega\pi \quad \omega\pi\pi \quad \eta\pi\pi \\
 \text{wavy} \text{---} \text{cross-hatched circle} \text{---} \text{wavy} = \text{wavy} \text{---} \text{double line} \text{---} \text{wavy} + \text{wavy} \text{---} \text{dome} \text{---} \text{wavy} + \text{wavy} \text{---} \text{two circles} \text{---} \text{wavy} + \text{wavy} \text{---} \text{circle with cross} \text{---} \text{wavy} + \dots
 \end{array}$$

FIG. 3. Vector Meson Dominance representation of hadronic vacuum polarization.