

DFTT 33/96
INFNCA-TH9615
hep-ph/9607334

INCLUSIVE PRODUCTION OF HADRONS IN $\ell^\uparrow p^\uparrow \rightarrow h^\uparrow X$ AND SPIN MEASUREMENTS*

M. ANSELMINO, M. BOGLIONE

*Dipartimento di Fisica Teorica, Università di Torino and
INFN, Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy*

J. HANSSON

Department of Physics, Luleå University of Technology, S-97187 Luleå, Sweden

F. MURGIA

INFN, Sezione di Cagliari, Via A. Negri 18, I-09127 Cagliari, Italy

We discuss the production of polarized hadrons in polarized lepton nucleon interactions and show that the helicity density matrix of the hadron, when measurable, can give information on the spin structure of the nucleon and the spin dependence of the quark fragmentation process. Single spin asymmetries in the $\ell N^\uparrow \rightarrow hX$ process are also briefly discussed.

According to the QCD hard scattering scheme and the factorization theorem^{1-3,4,5} the helicity density matrix of the hadron h inclusively produced in the DIS process $\ell^\uparrow N^\uparrow \rightarrow h^\uparrow X$ is given by

$$\rho_{\lambda_h, \lambda'_h}^{(s, S)}(h) \frac{E_h d^3 \sigma^{\ell, s+N, S \rightarrow h+X}}{d^3 \mathbf{p}_h} = \sum_{q; \lambda_\ell, \lambda_q, \lambda'_q} \int \frac{dx}{\pi z} \frac{1}{16\pi x^2 s^2} \quad (1)$$

$$\rho_{\lambda_\ell, \lambda_\ell}^{\ell, s} \rho_{\lambda_q, \lambda'_q}^{q/N, S} f_{q/N}(x) \hat{M}_{\lambda_\ell, \lambda_q; \lambda_\ell, \lambda_q}^q \hat{M}_{\lambda_\ell, \lambda'_q; \lambda_\ell, \lambda'_q}^{q*} D_{\lambda_h, \lambda'_h}^{\lambda_q, \lambda'_q}(z)$$

where $\rho^{\ell, s}$ is the helicity density matrix of the initial lepton with spin s , $f_{q/N}(x)$ is the number density of unpolarized quarks q with momentum fraction x inside an unpolarized nucleon and $\rho^{q/N, S}$ is the helicity density matrix of quark q inside the polarized nucleon N with spin S . The $\hat{M}_{\lambda_\ell, \lambda_q; \lambda_\ell, \lambda_q}^q$'s are the helicity amplitudes for the elementary process $\ell q \rightarrow \ell q$. The final lepton spin is not observed and helicity conservation of perturbative QCD and QED has already been taken into account in the above equation: as a consequence only the

* Talk delivered by M. Anselmino at DIS96, Rome, April 15-19, 1996.

diagonal elements of $\rho^{\ell,s}$ contribute to $\rho(h)$ and non diagonal elements, present in case of transversely polarized leptons, do not contribute. $D_{\lambda_h, \lambda'_h}^{\lambda_q, \lambda'_q}(z)$ is the product of *fragmentation amplitudes*

$$D_{\lambda_h, \lambda'_h}^{\lambda_q, \lambda'_q}(z) = \oint_{X, \lambda_X} \mathcal{D}_{\lambda_X, \lambda_h; \lambda_q} \mathcal{D}_{\lambda_X, \lambda'_h; \lambda'_q}^* \quad (2)$$

where the \oint_{X, λ_X} stands for a spin sum and phase space integration of the undetected particles, considered as a system X . The usual unpolarized fragmentation function $D_{h/q}(z)$, *i.e.* the density number of hadrons h resulting from the fragmentation of an unpolarized quark q and carrying a fraction z of its momentum, is given by

$$D_{h/q}(z) = \frac{1}{2} \sum_{\lambda_q, \lambda_h} D_{\lambda_h, \lambda_h}^{\lambda_q, \lambda_q}(z) = \frac{1}{2} \sum_{\lambda_q, \lambda_h} D_{h_{\lambda_h}/q_{\lambda_q}}(z), \quad (3)$$

where $D_{\lambda_h, \lambda_h}^{\lambda_q, \lambda_q}(z) \equiv D_{h_{\lambda_h}/q_{\lambda_q}}$ is a polarized fragmentation function, *i.e.* the density number of hadrons h with helicity λ_h resulting from the fragmentation of a quark q with helicity λ_q . Notice that by definition and parity invariance the generalized fragmentation functions (2) obey the relationships

$$D_{\lambda_h, \lambda'_h}^{\lambda_q, \lambda'_q} = \left(D_{\lambda'_h, \lambda_h}^{\lambda'_q, \lambda_q} \right)^* \quad (4)$$

$$D_{-\lambda_h, -\lambda'_h}^{-\lambda_q, -\lambda'_q} = -(-1)^{2S_h} (-1)^{\lambda_q + \lambda'_q + \lambda_h + \lambda'_h} D_{\lambda_h, \lambda'_h}^{\lambda_q, \lambda'_q}, \quad (5)$$

where S_h is the hadron spin; notice also that collinear configuration (intrinsic $\mathbf{k}_\perp = 0$) together with angular momentum conservation in the forward fragmentation process imply

$$D_{\lambda_h, \lambda'_h}^{\lambda_q, \lambda'_q} = 0 \quad \text{when} \quad \lambda_q - \lambda'_q \neq \lambda_h - \lambda'_h. \quad (6)$$

Eq. (1) holds at leading twist, leading order in the coupling constants and large Q^2 values; the intrinsic \mathbf{k}_\perp of the partons have been integrated over and collinear configurations dominate both the distribution functions and the fragmentation processes. For simplicity of notations we have not indicated the Q^2 scale dependences in f and D ; the variable z is related to x by the usual imposition of energy momentum conservation in the elementary $2 \rightarrow 2$ process; more technical details can be found in Ref. ⁵.

The quark helicity density matrix $\rho^{q/N,S}$ can be decomposed as

$$\rho^{q/N,S} = P_P^{q/N,S} \rho^{N,S} + P_A^{q/N,S} \rho^{N,-S} \quad (7)$$

where $P_{P(A)}^{q/N,S}$ (which, in general, depends on x) is the probability that the spin of the quark inside the polarized nucleon N is parallel (antiparallel) to the nucleon spin S and $\rho^{N,S(-S)}$ is the helicity density matrix of the nucleon with spin $S(-S)$. Notice that

$$P^{q/N,S} = P_P^{q/N,S} - P_A^{q/N,S} \quad (8)$$

is the component of the quark polarization vector along the parent nucleon spin direction. In more familiar notations one has, for longitudinally polarized nucleons

$$f_{q/N}(x) P_{P(A)}^{q/N,S_L}(x) = f_{q_{+(-)}/N_+}(x) = f_{q_{-(+)}/N_-}(x) \quad (9)$$

where $f_{q_{+(-)}/N_+}$ is the polarized distribution function, that is the density number of quarks with helicity $+(-)$ inside a nucleon with helicity $+$ and the last equality holds due to parity invariance. This implies

$$f_{q/N} P^{q/N,S_L} = f_{q_{+}/N_+} - f_{q_{-}/N_+} \equiv \Delta q \quad (10)$$

and similarly for transverse polarization T ,

$$f_{q/N} P^{q/N,S_T} = f_{q,S_T/N,S_T} - f_{q,-S_T/N,S_T} \equiv \Delta_T q. \quad (11)$$

where $P^{q/N,S_T}$ is the quark transverse polarization in a transversely polarized nucleon.

We shall now consider several particular cases of Eq. (1) and discuss what can be learned or expected from a measurement of $\rho(h)$. We consider both the case of spin 1 and spin 1/2 hadrons h , at leading twist only; higher twist contribution to single spin asymmetries in $\ell N^\uparrow \rightarrow hX$ processes will be shortly discussed at the end. Somewhat similar analyses can be found in Ref. ⁶ and ⁷.

We choose xz as the hadron production plane with the lepton moving along the z -axis and the nucleon in the opposite direction in the lepton-nucleon centre of mass frame; as usual we indicate by an index L the (longitudinal) nucleon spin orientation along the z -axis, by an index S the (sideway) orientation along the x -axis and by an index N the (normal) orientation along the y -axis.

Spin 1 final hadron; unpolarized leptons and longitudinally polarized nucleons

In this case Eqs. (1) and (6) yield (using obvious shorter notations)

$$\begin{aligned}\rho_{1,1}^{(S_L)}(V) d^3\sigma &= \sum_q \int \frac{dx}{\pi z} f_{q/N} d\hat{\sigma}^q \left[P_A^{q/N, S_L} D_{V_1/q+} + P_P^{q/N, S_L} D_{V_1/q-} \right] \\ \rho_{0,0}^{(S_L)}(V) d^3\sigma &= \sum_q \int \frac{dx}{\pi z} f_{q/N} d\hat{\sigma}^q D_{V_0/q+} \\ \rho_{-1,-1}^{(S_L)}(V) d^3\sigma &= \sum_q \int \frac{dx}{\pi z} f_{q/N} d\hat{\sigma}^q \left[P_A^{q/N, S_L} D_{V_1/q-} + P_P^{q/N, S_L} D_{V_1/q+} \right]\end{aligned}\quad (12)$$

where the apex (S_L) reminds of the nucleon spin configuration.

Spin 1 final hadron; unpolarized leptons and transversely polarized nucleons, $T=S, N$

In this case we have both diagonal

$$\rho_{1,1}^{(S_T)}(V) d^3\sigma = \sum_q \int \frac{dx}{\pi z} f_{q/N} d\hat{\sigma}^q \frac{1}{2} [D_{V_1/q+} + D_{V_1/q-}] \quad (13)$$

$$\rho_{0,0}^{(S_T)}(V) d^3\sigma = \sum_q \int \frac{dx}{\pi z} f_{q/N} d\hat{\sigma}^q D_{V_0/q+} \quad (14)$$

$$\rho_{-1,-1}^{(S_T)}(V) = \rho_{1,1}^{(S_T)}(V) = \frac{1 - \rho_{0,0}^{(S_T)}(V)}{2} \quad (15)$$

and non diagonal matrix elements

$$\rho_{1,0}^{(S_S)}(V) d^3\sigma = \sum_q \int \frac{dx}{\pi z} f_{q/N} \frac{P^{q/N, S_S}}{2} [\text{Re} \hat{M}_+^q \hat{M}_-^{q*}] D_{1,0}^{+,-} \quad (16)$$

$$\rho_{-1,0}^{(S_S)}(V) = \rho_{1,0}^{(S_S)}(V) \quad (17)$$

$$\rho_{1,0}^{(S_N)}(V) = -\rho_{-1,0}^{(S_N)}(V) = i\rho_{1,0}^{(S_S)}(V). \quad (18)$$

which involve the non diagonal fragmentation functions (2); $\hat{M}_{+(-)}$ is a short notation for $\hat{M}_{+,+,+,+(+,-,+,-)}/4\sqrt{s}$.

Spin 1/2 final hadron; unpolarized leptons and polarized nucleons

In case of final spin 1/2 hadrons ($h = B$), with unpolarized leptons and spin S nucleons, we have the non zero results

$$P_x^{(S_S)} d^3\sigma = \sum_q \int \frac{dx}{\pi z} f_{q/N} P^{q/N, S_S} [\text{Re} \hat{M}_+^q \hat{M}_-^{q*}] D_{+,-}^{+,-}$$

$$= \sum_q \int \frac{dx}{\pi z} \Delta_T q \Delta_N \hat{\sigma}^q \Delta_T D_{B/q} \quad (19)$$

$$P_y^{(S_N)} = -P_x^{(S_S)} \quad (20)$$

$$\begin{aligned} P_z^{(S_L)} d^3\sigma &= \sum_q \int \frac{dx}{\pi z} f_{q/N} P^{q/N, S_L} d\hat{\sigma}^q [D_{B_+/q_-} - D_{B_+/q_+}] \\ &- \sum_q \int \frac{dx}{\pi z} \Delta q d\hat{\sigma}^q \Delta D_{B/q}. \end{aligned} \quad (21)$$

where $P_i = \text{Tr}(\sigma^i \rho)$ are the components of the polarization vector in the helicity rest frame of hadron B , and where the apices S_L, S_N and S_S refer to the nucleon spin orientations in the reference frame where we compute the scattering. In the above equations

$$- [\text{Re} \hat{M}_+^q \hat{M}_-^{q*}] = \frac{d\hat{\sigma}^{\ell+q, S_N \rightarrow \ell+q, S_N}}{d\hat{t}} - \frac{d\hat{\sigma}^{\ell+q, S_N \rightarrow \ell+q, -S_N}}{d\hat{t}} \equiv \Delta_N \hat{\sigma}^q, \quad (22)$$

$$D_{+, -}^{+, -} = D_{B, S_N/q, S_N} - D_{B, -S_N/q, S_N} \equiv \Delta_T D_{B/q}, \quad (23)$$

which is a difference of transverse fragmentation functions and

$$D_{B, S_L/q, S_L} - D_{B, -S_L/q, S_L} = D_{B_+/q_+} - D_{B_-/q_+} \equiv \Delta D_{B/q}. \quad (24)$$

Longitudinally polarized leptons and polarized nucleons

We discuss now the case of polarized leptons; as we noticed after Eq. (1) only the diagonal elements of the lepton helicity density matrix $\rho^{\ell, s}$ contribute to $\rho(h)$, so that only longitudinal polarizations could affect the results. We consider then longitudinally polarized leptons, $s = s_L$.

One obtains the same results as in the unpolarized lepton case for the non diagonal matrix elements and slightly different ones for the diagonal elements, for example

$$\rho_{0,0}^{(s_L, S_L)}(V) d^3\sigma_L = \sum_q \int \frac{dx}{\pi z} [f_{q-/N_+} |\hat{M}_+^q|^2 + f_{q+/N_+} |\hat{M}_-^q|^2] D_{V_0/q_+} \quad (25)$$

$$P_z^{(s_L, S_T)}(B) d^3\sigma = \sum_q \int \frac{dx}{\pi z} f_{q/N} \frac{1}{2} [|\hat{M}_+^q|^2 - |\hat{M}_-^q|^2] \Delta D_{B/q}. \quad (26)$$

Experimental measurements

Some elements of the helicity density matrix of the produced hadrons can be measured via the angular distribution of the final hadron h decay; typical examples are the $\rho \rightarrow \pi\pi$ and $\Lambda \rightarrow p\pi$ decays.

For spin 1 hadrons one can measure $\rho_{0,0}$ and $\rho_{1,0}$ [Eqs. (12) and (16)], whereas for weakly decaying spin 1/2 hadrons one measures $P_y^{S_N}$ and $P_z^{S_L}$ [Eqs. (20) and (21)]. Such measurements supply information on the polarized quark fragmentation process and the polarized distribution functions. Further discussion and an estimate of $\rho_{1,0}$ can be found in Ref. ⁵. We only remind here that according to $SU(6)$ wave function the entire Λ polarization is due to the strange quark, so that the difference of polarized fragmentation functions in Eq. (21) is different from zero only for s quarks, $\Delta D_{\Lambda/s} = D_{\Lambda/s}$. Then Eq. (21) reads

$$P_z^{(S_L)} = - \frac{\int dx (\pi z)^{-1} \Delta s d\hat{\sigma}^s D_{\Lambda/s}}{\sum_q \int dx (\pi z)^{-1} f_{q/N} d\hat{\sigma}^q D_{\Lambda/q}} \quad (27)$$

Such a quantity is expected to be rather small; however, any non zero value would offer valuable information on the much debated issue of longitudinal strange quark polarization, Δs , inside a longitudinally polarized nucleon. A similar information on the transverse polarization can be obtained from a measurement of $P_y^{(S_N)}$ and Eq. (20).

Single spin asymmetries and \mathbf{k}_\perp effects in $\ell N^\uparrow \rightarrow hX$

We conclude by mentioning that a fundamental property of quark fragmentation, the quark analysing power

$$A_{h/q} = \frac{\tilde{D}_{h/q^\uparrow}(z, \mathbf{k}_\perp) - \tilde{D}_{h/q^\downarrow}(z, \mathbf{k}_\perp)}{\tilde{D}_{h/q^\uparrow}(z, \mathbf{k}_\perp) + \tilde{D}_{h/q^\downarrow}(z, \mathbf{k}_\perp)} = \frac{\tilde{D}_{h/q^\uparrow}(z, \mathbf{k}_\perp) - \tilde{D}_{h/q^\uparrow}(z, -\mathbf{k}_\perp)}{\tilde{D}_{h/q^\uparrow}(z, \mathbf{k}_\perp) + \tilde{D}_{h/q^\uparrow}(z, -\mathbf{k}_\perp)} \quad (28)$$

has been recently proposed ^{2,3} and suggested to be sizeable. $\tilde{D}_{h/q^\uparrow}(z, \mathbf{k}_\perp)$ denotes the \mathbf{k}_\perp dependent fragmentation function of the polarized quark q .

Such an effect, a leading twist one, might be measured by looking at the production of, say, pions with opposite intrinsic \mathbf{k}_\perp inside the current jet in the scattering of unpolarized leptons off transversely polarized protons; or, equivalently, to the production of pions with a certain \mathbf{k}_\perp resulting from up or down transversely polarized protons.

This same effects might also be responsible, as suggested for the process $p^\uparrow p \rightarrow \pi X$ ⁸⁻¹⁰, for the higher twist single spin asymmetry

$$A_N = \frac{d\sigma^{\ell N^\uparrow \rightarrow hX} - d\sigma^{\ell N^\downarrow \rightarrow hX}}{d\sigma^{\ell N^\uparrow \rightarrow hX} + d\sigma^{\ell N^\downarrow \rightarrow hX}} \quad (29)$$

where integration is performed over the hadron h intrinsic \mathbf{k}_\perp . A measurement of such an asymmetry would provide valuable information.

References

1. J.C. Collins, *Nucl. Phys. B* **394**, 169 (1993).
2. J.C. Collins, *Nucl. Phys. B* **396**, 161 (1993).
3. J.C. Collins, S.H. Heppelmann and G.A. Ladinsky, *Nucl. Phys. B* **420**, 565 (1994).
4. For earlier work on spin asymmetries and helicity density matrices in hard scatterings see also J. Babcock, E. Monsay and D. Sivers, *Phys. Rev. D* **19**, 1483 (1978); M. Anselmino and P. Kroll, *Phys. Rev. D* **30**, 36 (1984); N.S. Craigie, K. Hidaka, M. Jacob and F.M. Renard, *Phys. Rep.* **99**, 69 (1983).
5. M. Anselmino, M. Boglione, J. Hansson and F. Murgia, *preprint* DFTT 61/95 and INFNCA-TH9526, e-Print Archive: hep-ph 9512379, to appear in *Phys. Rev. D*.
6. P.J. Mulders and R.D. Tangerman, *preprint* NIKHEF 95-053, e-Print Archive: hep-ph 9510301.
7. R.L. Jaffe and A. Manohar, *Nucl. Phys. B* **321**, 343 (1989).
8. X. Artru, J. Czyzewski and H. Yabuki, *preprint* TPJU-13-95, e-Print Archive: hep-ph/9508239.
9. M. Anselmino, M. Boglione and F. Murgia, *Phys. Lett. B* **362**, 164 (1995).
10. M. Anselmino, M. Boglione and F. Murgia, in Proceedings of Adriatico Research Conference on *Trends in Collider Spin Physics*, Trieste, Italy, 4-8 Dec 1995, e-Print Archive: hep-ph/9604397.