

WM-96-107  
July 1996

## The Coleman-Weinberg Phase Transition in Extended Higgs Models

Marc Sher

*Physics Department, College of William and Mary, Williamsburg, VA 23187, USA*

In Coleman-Weinberg symmetry breaking, all dimensionful parameters vanish and the symmetry is broken by loop corrections. Before Coleman-Weinberg symmetry breaking in the Standard Model was experimentally ruled out, it had already been excluded on cosmological grounds. In this Brief Report, the cosmological analysis is carried out for Coleman-Weinberg models with extended Higgs sectors, which are not experimentally ruled out, and general constraints on such models are given.

In the Standard Model, supplemented by General Relativity, there are only two dimensionful parameters: the squares of the Higgs mass term,  $\mu$ , and the Planck mass. The ratio of these two numbers must be smaller than  $10^{-34}$ . In a fundamental theory, this ratio must be explained. A significant fraction of research in theoretical physics in the past fifteen years has focused on models which can explain such a small number.

It might eventually be easier to explain this small ratio if it vanishes completely. This is somewhat similar to the assumption made in cosmology that the cosmological constant vanishes; since it must be less (in Planck units) than  $10^{-122}$ , it is generally assumed that zero will be an easier number to explain. Likewise, the  $\mu^2 = 0$  limit could also prove easier to explain than  $\mu^2 \sim 10^{-34}$  in Planck units. The  $\mu^2 = 0$  case was originally considered by Coleman and E. Weinberg(CW)[1], who showed that loop corrections will still break the electroweak symmetry.

Since a parameter has been set to zero, a prediction can be made, and the Higgs mass can be determined in terms of the gauge boson and fermion masses. The result gives  $m_h < 10$  GeV, and requires  $m_t < m_Z$ , which are both in contradiction with experiment. As a result, Coleman-Weinberg symmetry breaking in the Standard Model is ruled out. It is not ruled out, however, in models with more complicated Higgs sectors[2].

Models with extended Higgs sectors have become more popular recently with the realization that sphaleron effects could erase any previously generated baryon asymmetry[3], and that CP violation in the standard model is too small for electroweak baryogenesis[4]. It has been shown[5] that models with extended Higgs sectors can avoid washing out any previous baryon number, as well as generate a sufficient baryon asymmetry. In fact, generation of a baryon asymmetry also requires a first order electroweak phase transition, and CW models always have a first order transition. This leads to greater incentive for considering CW symmetry breaking in extended Higgs models.

Recently, several papers[6, 7] have discussed CW symmetry breaking in extended

Higgs models. In particular, a simple model of Hempfling[7] showed that addition of a singlet field yields a satisfactory mass spectrum in the CW case. The details of the cosmological phase transition in these models, however, were not considered, and this poses a potential difficulty for these models.

Before experiments ruled out CW symmetry breaking in the Standard Model, it had been shown[2, 8] that it was already ruled out on cosmological grounds; either the phase transition never finished or it finished with far too much entropy generation to be acceptable. In this Brief Report, I will examine this issue for the general case, and will present a bound which can be applied to any specific model of electroweak CW symmetry breaking.

First, let us review the cosmology of the CW model in the Standard Model. Minimizing the potential[2], one finds the zero-temperature potential to be

$$V = B\phi^4 \left( \ln(\phi^2/\langle\phi\rangle^2) - \frac{1}{2} \right) \quad (1)$$

where

$$B \equiv \frac{1}{64\pi^2\langle\phi\rangle^4} (6M_W^4 + 3M_Z^4 - 12M_t^4) \quad (2)$$

and we must have  $B > 0$  to have spontaneous symmetry breaking. One immediately sees why CW symmetry breaking is ruled out for the known top quark mass, since that gives  $B < 0$ . However, suppose the top mass were smaller, say 40 GeV; then we would have  $B > 0$ . The Higgs mass would then be given by  $m_H^2 = 8B\langle\phi\rangle^2$ .

At finite temperature, a temperature dependent term must be added. For values of  $T \gg \phi$ , this gives a term

$$V_T = -\frac{1}{90}\eta\pi^2T^4 + \frac{1}{24\langle\phi\rangle^2}\phi^2T^2(6M_W^2 + 3M_Z^2 + 6M_t^2) \quad (3)$$

which causes a positive  $\phi^2$  term to appear, restoring the symmetry at high temperatures, and causing a minimum to occur at the origin at any temperature. As a result, the

Universe will start in the symmetric vacuum, and will have to tunnel through a potential barrier. Although the above expression is only valid at high temperature, the results of tunnelling out of the vacuum at the origin at high temperature are not appreciably affected by using the exact expression (note that this is not the case in determining, for example, the critical temperature at which the symmetric and asymmetric vacua are degenerate[9]).

The tunnelling rate for this model was calculated[10] long ago, and it was found that the transition will not occur until the temperature is below  $10^{-7}$  times the electroweak scale. This will result in an entropy increase during the transition by a factor of more than  $10^{21}$ , washing out any baryon asymmetry, and thus the model was ruled out. However, Witten[11] pointed out that the Universe can not cool below the QCD scale without chiral symmetry breaking giving  $\bar{\psi}\psi$  a vacuum expectation value, which breaks the electroweak gauge symmetry (the Yukawa term,  $\bar{\psi}\psi\Phi$  turns into a linear term when chiral symmetry breaks, thus destabilizing the symmetric vacuum. The entropy generated in this transition is roughly the cube of the ratio of the electroweak to QCD scales, or  $\sim 10^6$ , which is marginally acceptable. Finally, Flores and Sher[8] noted that, in addition to the QCD coupling growing as the Universe cools, the Yukawa coupling also grows. This will cause the coefficient  $B$  in the above to change sign at low temperatures, causing a barrier to appear which persists to zero temperature. Thus, the Universe will get stuck in a metastable vacuum, and if the transition occurs, far too much entropy would be generated. Thus, they concluded that CW symmetry breaking in the Standard Model is ruled out.

What happens in extended Higgs models? The question of CW symmetry breaking in extended Higgs models was first discussed by Gildener and Weinberg[12] and reviewed in Ref. 2. Consider the single Higgs case. There, one can choose the renormalization scale such that the tree-level potential vanishes, i.e. one can choose  $M_R$  such that  $\lambda(M_R) = 0$ .

Calculating the one loop potential, and eliminating  $M_R$  by minimizing and denoting the minimum  $\langle\phi\rangle$  gives the expression in Eq. 1. Now consider the multi-scalar case. The tree-level potential is given by  $V_o = f_{ijkl}\phi_i\phi_j\phi_k\phi_l$  where the  $\phi_i$  label all of the scalars in the model. As Gildener and Weinberg show, one cannot choose the renormalization scale such that  $V_o$  vanishes everywhere, since the scale at which one self-coupling vanishes is not the same as the scale at which another will vanish. However, suppose one defines  $\phi_i = N_i r$ , where  $N_i$  is a unit vector and  $r$  is the distance from the origin of field space, then  $V_o = f_{ijkl}N_iN_jN_kN_lr^4$ . Let the minimum value of  $V_o$  on the unit sphere occur for  $N_i = n_i$ . Then, Gildener and Weinberg show that the renormalization scale can be chosen so that  $V_o$  vanishes along the direction  $N_i = n_i$ .

The important point is that the tree level potential vanishes along a direction in field space, and in any other direction will be positive (if one has positivity of the potential at very large scales). Note, the condition for this to occur is a *single* condition on the  $f_{ijkl}$ ; i.e. one can choose the renormalization scale so that a combination of the  $f_{ijkl}$  vanishes. Now, if the tree-level potential vanishes along the ray  $\phi_i = n_i r$ , then one can calculate the potential along that ray. The result is (minimizing along that ray)

$$V = Br^4 \left( \ln(r^2/\langle r \rangle^2) - \frac{1}{2} \right) \quad (4)$$

where

$$B \equiv \frac{1}{64\pi^2\langle r \rangle^4} (3M_V^4 + M_S^4 - 4M_F^4) \quad (5)$$

and where  $M_V^4$ , for example, refers to the sum of the fourth powers of all vector boson masses. In any other direction, the tree-level potential is positive, the  $f_{ijkl}$  are not small, and radiative corrections are unimportant. The scalar masses-squared are all positive, with the scalar whose mass vanishes at tree-level (corresponding to the flat direction) getting a mass-squared of  $8B\langle r \rangle^2$ , just as in the Standard Model.

Let us give a couple of examples. In the two-Higgs model, considering the neutral

directions only, we can write the doublets as

$$\Phi_1 = \frac{r}{\sqrt{2}} \begin{pmatrix} 0 \\ N_1 \end{pmatrix}, \quad \Phi_2 = \frac{r}{\sqrt{2}} \begin{pmatrix} 0 \\ N_2 \end{pmatrix} \quad (6)$$

Expressing the potential (in the massless theory—see Ref. 2 for the full potential) in terms of these coordinates gives

$$V = \frac{1}{4}r^4[\lambda_1 N_1^4 + \lambda_2 N_2^4 + (\lambda_3 + \lambda_4 + \lambda_5)N_1^2 N_2^2] \quad (7)$$

Suppose the potential has a minimum value (which Gildener and Weinberg show can be chosen to be zero by judicious choice of renormalization scale) on the unit circle along the direction  $N_i = n_i$ . This direction can be found to be

$$n_1 = \frac{\sqrt{\lambda_2}}{\sqrt{\lambda_1} + \sqrt{\lambda_2}}, \quad n_2 = \frac{\sqrt{\lambda_1}}{\sqrt{\lambda_1} + \sqrt{\lambda_2}}, \quad 2\sqrt{\lambda_1 \lambda_2} + \lambda_3 + \lambda_4 + \lambda_5 = 0. \quad (8)$$

The first two of these specify the direction from the origin to the minimum and the third comes from the requirement that the potential vanish along this direction (this is no more “unnatural” than the assumption that a renormalization scale can be found in the Standard Model such that  $\lambda$  is small). The one-loop potential along that direction is then  $V = Br^4[\ln(r^2/\langle r \rangle^2) - \frac{1}{2}]$  where  $\langle r \rangle^2 = \langle \phi_1 \rangle^2 + \langle \phi_2 \rangle^2 = (246 \text{ GeV})^2$  and  $B$ , given in Eq. 5, includes contributions from the  $W, Z, t$  and the five scalars in the model. The lightest Higgs mass-squared is  $8B\langle r \rangle^2$ . This mass can exceed the experimental limit only if some of the scalar masses are quite large (greater than several hundred GeV).

Another example is the model of Hempfling[7], with a singlet  $S$  added. The tree-level potential is  $V_o = \frac{1}{2}\lambda_\phi(\phi^\dagger\phi)^2 + \frac{1}{2}\lambda_S(S^\dagger S)^2 - \lambda_X(\phi^\dagger\phi)(S^\dagger S)$ . Hempfling defines  $\phi \equiv r \sin \beta$  and  $S \equiv r \cos \beta$ ; one sees that  $r$  has the same definition as the above and  $\tan \beta = n_2/n_1$  above. The flat direction is given by  $\tan \beta = \lambda_X/\lambda_\phi$  and the condition that the tree-level potential vanish at the minimum is given by  $\lambda_\phi\lambda_S = \lambda_X^2$ . The one-loop potential is then given by  $V = Br^4[\ln(r^2/\langle r \rangle^2) - \frac{1}{2}]$  where  $B$  is also given in Eq.5. Note that a major

difference is that  $\langle r \rangle$  is not, as in the previous example, constrained to be 246 GeV, but could be much larger, since the singlet vacuum expectation value is unconstrained. This means, for example, that the contributions to  $B$  from the  $W$ ,  $Z$  and top quark are smaller than in the standard model by a factor of  $\left(\frac{246 \text{ GeV}}{r}\right)^4$  (this can be seen in the  $\sin^4 \beta$  factor in Eq. 4 of Hempfling[7]). As a result,  $B$  can be positive without requiring enormous scalar masses. Again, the lightest Higgs mass-squared is  $8B\langle r \rangle^2$ , in agreement with Hempfling. Note that since  $\langle r \rangle$  can be quite large, this mass can easily satisfy the experimental bounds.

We now include the effects of high temperature. Note that the potential is flat along one direction, and large and positive in other directions. We can thus effectively consider the problem to be a one-dimensional problem along the flat direction. In the high temperature limit, the temperature-dependent term can be written as

$$V_T = \frac{1}{24} r^2 T^2 M_2 \quad (9)$$

where

$$M_2 = \frac{3M_V^2 + M_S^2 + 2M_F^2}{\langle r \rangle^2} \quad (10)$$

and  $M_V^2$  ( $M_S^2$ ,  $M_F^2$ ) stands for the sum of the squares of the masses of all of the vector bosons (scalars, fermions) in the model. Note that  $M_2$  is always positive, thus CW models always have symmetry restoration at high temperatures (unlike non-CW models, which, in some cases, can have symmetry anti-restoration[2]). As discussed above, we have verified in several sample cases that the high-temperature limit is satisfactory for this calculation, to within the accuracy stated.

The full potential is thus relatively simple. It has two parameters,  $B$  and  $M_2$ , and is a function only of the radial co-ordinate,  $r$ . We can now examine the phase transition as a function of these two parameters[13]. The procedure is standard and straightforward—the reader is referred to Ref. 2 for a review. The transition temperature is found by

comparing the bubble nucleation rate with the expansion rate (the  $O(3)$  symmetric action is always the lowest here), and the reheating temperature is found by equating the energy densities just prior to and just after the transition. The cube of the ratio (times a factor of  $O(1)$  involving the number of states) gives the increase in entropy generated by the transition.

The ratio of the final to initial entropy is plotted in Fig. 1 for various values of  $B$  and  $M_2$ . Recall that the observed baryon number to entropy ratio today is approximately  $10^{-9}$ , and in many models this requires very efficient baryogenesis. Should the entropy increase in the transition be a factor of 10 or 100, then the transition will not play a significant role (it is unlikely that a specific model will be able to predict the baryon number to entropy ratio more accurately than an order of magnitude). Should the entropy increase be a factor of  $10^4 - 10^6$ , then an extremely efficient mechanism for baryogenesis must be found. The entropy increase cannot be much more than  $10^6$ , since the transition temperature will be near or below the QCD scale, and the transition will be driven by QCD condensate formation[14].

For regions of parameter-space in which the transition is driven by QCD, the entropy increase of at least  $10^6$  requires an initial baryon-number to entropy ratio close to  $10^{-3}$ . It is difficult, but perhaps not impossible, to construct a baryogenesis mechanism which can generate such a large ratio. However, such regions of parameter-space have another, more serious, problem.

As the Universe cools in the symmetric phase, all of the quarks, vector bosons and Higgs bosons are massless. At what temperature will  $\bar{\psi}\psi$  condense? The QCD beta function is flatter than in the standard model, since all six quarks contribute to very low scales—this will lower the chiral symmetry breaking transition temperature by roughly (very roughly) a factor of 3. However another factor will increase the transition temperature. The top quarks will not only attract each other via QCD, but also via Higgs



exchange (recall that the Higgs boson is massless). As the Universe cools, the top quark Yukawa coupling will rapidly reach its fixed point, which (at one-loop) is  $\frac{4}{3}$  times the QCD coupling. This additional attraction will break the  $SU(6)$  symmetry and cause top quark condensation to occur at a higher temperature than that of the other quarks. The transition temperature will thus be somewhat similar to the chiral symmetry breaking temperature (roughly 200 MeV). Note that the idea of top quark condensation driving the electroweak symmetry breaking has been of great interest[15], but here the condensation drives electroweak symmetry breaking below the GeV scale!

The serious problem occurs because of the growth of the top quark Yukawa coupling as the Universe cools in the symmetric phase. How does this affect the potential? As discussed in detail in Ref. 2, the potential we have considered is *not* “renormalization-group improved”, and neglected terms are approximately of  $O(\frac{g^2}{4\pi} \ln(\frac{Q_1}{Q_2}))$  times those included, where  $g$  is a coupling and  $Q_1$  and  $Q_2$  are the largest and smallest scales in the region of interest. Since we are only considering values of  $r$  between the QCD and electroweak scales, the logarithm is not too large, and the effects are only relevant for the top quark Yukawa coupling. In this case, it is sufficient to “run” the top quark Yukawa coupling (and QCD coupling, which enters in the beta function for the Yukawa coupling) in the expression for  $B$  (this has been verified using the full expression). Thus, as the top quark Yukawa coupling grows,  $B$  will eventually change sign. The potential near the origin, instead of being negative at zero temperature, will be positive, leading to a barrier at zero temperature. Even when chiral symmetry breaks, this barrier is likely to persist, and the Universe will get stuck in a metastable state; significantly increasing the entropy generation further. Although perturbative techniques are questionable at these low scales, it does appear that such models, already endangered by the entropy production required, are in further jeopardy, and likely ruled out.

The results of this Brief Report can be summarized as follows. For any CW model,

one finds the sum of the squares of all scalar vacuum expectation values, calling it  $\langle r \rangle^2$ . Plugging that and the masses of all of the particles in the model into Eqs. 5 and 10, one finds  $B$  and  $M_2$ . Looking at Figure 1, the entropy generation can be determined. If it is less than  $10^2$ , the effects of the transition are small; if it is greater than  $10^6$ , in the QCD region, the transition is apparently fatal. In between, very efficient baryogenesis is required. Let us illustrate this procedure with two simple examples. In the Standard Model with a 40 GeV top quark, we have  $\langle r \rangle = 246$  GeV and thus  $M_2 = 1.22$  and  $64\pi^2 B = 0.12$ , and from Figure 1 we see that the model is excluded, in agreement with previous results. In the Hempfling model, in the limit in which  $\langle r \rangle \gg 246$  GeV, the only relevant scalar mass is given by  $M_S^2 = 6\lambda_S \langle r \rangle^2$  and there is a gauge boson of mass  $g_X \langle r \rangle$ . This gives a value of  $M_2 = 6\lambda_S + 3g_X^2$  and  $64\pi^2 B = 36\lambda_S^2 + 3g_X^4$ . For  $\alpha_X = .01$ , we find that the entropy generation is less than  $10^2$  for  $\lambda_S > .15$ , is between  $10^2$  and  $10^6$  for  $.10 < \lambda_S < .15$  and is greater than  $10^6$  (i.e. the model is excluded) for  $\lambda_S < .10$ . A similar procedure can be applied to any Coleman-Weinberg electroweak model.

I thank Carl Carlson for useful discussions. This work was supported by the National Science Foundation.

## References

- [1] S. Coleman and E. Weinberg, Phys. Rev. **D7**, 1888 (1973).
- [2] M. Sher, Phys. Reports **179**, 273 (1989).
- [3] V.A. Kuzmin, V.A. Rubakov and M.E. Shaposhnikov, Phys. Lett. **B155**, 36 (1985);  
M.E. Shaposhnikov Nucl. Phys. **B287**, 757 (1987); See the Proceedings of the NATO  
Advanced Research Workshop: Electroweak physics and the early universe, Sintra,  
1994 (Plenum Press), and references therein.

- [4] M.B. Gavela, P. Hernandez, J. Orloff and O. Pene, Mod. Phys. Lett. **A9**, 795 (1994);  
P. Huet and E. Sather, Phys. Rev. **D51**, 379 (1995).
- [5] A.I. Bochkarev, S.V. Kuzmin and M.E. Shaposhnikov, Phys. Lett. **B244**, 275 (1990), Phys. Rev. **D43**, 369 (1991); A.G. Cohen, D.B. Kaplan and A.E. Nelson, Phys. Lett. **B245**, 561 (1990), Nucl. Phys. **B349**, 727 (1991). B. Kastening, R.D. Peccei, X. Zhang, Phys. Lett. **B266**, 413 (1991); N. Turok and J. Zadrozny, Nucl. Phys. **B369**, 729 (1991); S. Myint, Phys. Lett. **B287**, 325 (1992); J.R. Espinosa, M. Quiros and F. Zwirner, Phys. Lett. **B307**, 106 (1993).
- [6] S. Bornholdt, N. Tetradis and C. Wetterich, Phys. Lett. **B348**, 89 (1995); K. Takemura, Prog. Theor. Phys. **95**, 609, 1996.
- [7] R. Hempfling, Phys. Lett. **B379**, 153 (1996).
- [8] R. A. Flores and M. Sher, Nucl. Phys. **B238**, 702 (1984).
- [9] R. Easther and W. Moreau, J. Phys. G **18**, 1869 (1992).
- [10] A. Guth and E.J. Weinberg, Phys. Rev. Lett. **45**, 1131 (1980).
- [11] E. Witten, Nucl. Phys. **B177**, 477 (1981).
- [12] E. Gildener, Phys. Rev. **D13**, 1025 (1976); E. Gildener and S. Weinberg, Phys. Rev. **D13**, 3333 (1976).
- [13] Our approach is similar to that of O. Bertolami, Phys. Lett. **B176**, 416 (1986).  
In that work, however, the expression for the high-temperature limit is incorrect, the thin wall approximation (which may not be valid) is used, and the author only considers the condition in which the transition is driven by QCD.

- [14] In making this statement, we assumed that  $r$  was 246 GeV at the minimum. In models with singlets, it could be closer to the TeV scale, in which case the entropy generation which occurs if the transition is driven by QCD condensates is more like  $10^8$  than  $10^6$ . As one can see from Fig. 1, this makes very little difference in the result.
- [15] W.A. Bardeen, C.T. Hill and M. Lindner, Phys. Rev. **D41**, 1647 (1990).

Figure 1: For any electroweak CW model, the values of  $M_2$  and  $B$ , as defined in the text, can be determined. Here is plotted the entropy generated in the phase transition as a function of those parameters. In the region below the bottom line, the transition is broken by  $\bar{t}t$  condensation; since the entropy generated is at least  $10^6$ , and a metastable vacuum is likely to form, this region is apparently excluded. For a given  $M_2$ , the uncertainty in  $64\pi^2 B$  due to the high-temperature approximation and due to uncertainties in the expansion rate is approximately 0.2. For  $M_2$  below 0.5, the uncertainties are larger, however the value of  $M_2$  exceeds 0.5 in all models that have appeared to date.