

# Dispersion Analysis of the Strange Vector Form Factors of the Nucleon

Hilmar Forkel

*European Centre for Theoretical Studies in Nuclear Physics and Related Areas,*

*Villa Tambosi, Strada delle Tabarelle 286, I-38050 Villazzano, Italy*

(September 1995)

## Abstract

We analyze the nucleon matrix element of the strange quark vector current in a nucleon-model independent dispersive approach with input from the current world data set for the isoscalar electromagnetic form factors. The update of Jaffe's minimal 3-pole ansatz for the spectral functions yields a 40% larger (Sachs) strangeness radius,  $(r_s^2)_{Sachs} = 0.20 \text{ fm}^2$ , and a by 20% reduced magnitude of the strangeness magnetic moment,  $\mu_s = -0.26$ . In the pole approximation these values are shown to be upper bounds. After extending the ansatz in order to implement the asymptotic QCD momentum dependence (which the 3-pole form factors cannot reproduce), we find the magnitude of the 3-pole results reduced by up to a factor of 2.5. The signs of the leading moments originate primarily from the large  $\phi$ -meson couplings and are generic in the pole approximation.

PACS numbers: 14.20.Dh, 12.40Vv, 12.15Mm, 12.40.-y, 12.39.Ki

## I. INTRODUCTION

Nonvalence quark distributions in hadrons arise from subtle and little understood quantum effects in the hadronic wave functions, which provide a unique key to hadron structure beyond the naive quark model [1]. The sea quark distributions of the nucleon in the strangeness and charm sectors are particularly interesting and much studied examples. While charm admixtures are mainly probed in hard scattering processes [2], strangeness fluctuations in the nucleon can produce (due to the smaller strange quark mass) much larger effects which are in some Lorentz channels directly measurable at low energies. Mounting experimental evidence indeed indicates significant values for the nucleon matrix elements of the strange scalar [3–5] and axial vector [6–8] currents.

In order to further advance the understanding of the nucleon’s strangeness content, both experimental and theoretical studies beyond these two channels are crucial. For once, the channel dependence provides insight into the dynamical origin of the strange-quark distribution. Flavor mixing instanton-induced interactions, for example, reveal themselves in a pronounced and characteristic channel dependence pattern [9]. Furthermore, some controversial assumptions in the analysis of the existing data (for example in the extrapolation and small- $Q^2$  evolution and in the treatment of  $SU(3)$  violations) can be avoided in other channels.

The present paper deals with the still unmeasured vector channel, i.e. with the nucleon matrix element of the strange vector current. This matrix element is experimentally accessible at low energies and has some useful theoretical properties. In close analogy with and as a part of the electromagnetic distributions it describes the nucleon’s strangeness charge and current distributions by Dirac and Pauli form factors. Furthermore, strangeness conservation renders these form factors scale independent (up to weak corrections), which avoids complications due to nonperturbative evolution from the outset.

As already mentioned, essentially no experimental information on the vector form factors exists up to now (apart from a reanalysis of older neutrino scattering data [10] with too poor

statistics to be conclusive). With the present experimental techniques, however, they can be directly measured by parity-violating lepton scattering off different hadronic targets [11–13]. Four experiments of this type are in preparation at CEBAF [13–16] and MAMI [13,17], while SAMPLE [18] at Bates already started to take data. These experiments will measure for the first time sea quark effects in hadrons at low energies.

In anticipation of the forthcoming data several theoretical estimates of the strange form factors, primarily on the basis of nucleon models, have appeared in the literature. Since sea quark effects arise from a delicate interplay of quantum effects in QCD, their reproduction in hadron models is much more challenging than the calculation of the standard static observables. Reflecting these difficulties, present nucleon model estimates [19–26] contain large and often uncontrolled theoretical uncertainties. For the Dirac form factor, in particular, the predictions vary by over an order of magnitude and in their sign. A comparison of some of these estimates can be found in Ref. [24]. Lattice simulations of the strange form factors have not yet been carried out since the computational demands increase substantially when quark-line disconnected contributions have to be taken into account (see however [27]).

In the present paper we bypass the problems associated with these dynamical calculations by pursuing a dispersive, nucleon-model independent approach. It was initiated by Jaffe [28] and becomes practicable since the isoscalar electromagnetic current carries the same quantum numbers as the strange current and thus couples to the nucleon through the same intermediate states. The available experimental data on the electromagnetic nucleon form factors can therefore be used as input for the strange form factor analysis.

After updating the minimal dispersive analysis of Jaffe with input from the current world data set for the electromagnetic form factors, we will focus on extensions of the spectral functions which implement information from QCD at high momentum transfers. We will be particularly interested in the implications of the QCD asymptotics for the low-momentum behavior of the form factors and their first non-vanishing moments, i.e. the strangeness radius and magnetic moment. A better understanding of the low-momentum regime is also needed for the experimental determination of the moments: since one has to extrapolate

the data to zero momentum transfer, the resulting values will be sensitive to the assumed low-momentum dependence of the form factors [24].

The paper is organized as follows. In the next section we outline the general ideas behind the dispersive treatment of the strange form factors and describe their implementation in some detail. We then update Jaffe's minimal 3-pole estimate in Section III with input from new fits to the electromagnetic form factors. In Section IV we discuss several extensions of the approach in pole approximation which are designed to adapt the large (spacelike) momentum behavior to predictions from QCD counting rules. Some generic features of the dispersive analysis and their impact on the sign of the moments are pointed out in section V. Section VI, finally, summarizes the main results and contains our conclusions. A partial summary of these results was reported previously in [29].

## II. STRANGE VECTOR FORM FACTORS AND 3-POLE ESTIMATE

In the absence of time reversal violations the nucleon matrix element of the strange quark vector current<sup>1</sup>

$$J_\mu^s = \bar{q}\gamma_\mu(B - Y)q = \bar{s}\gamma_\mu s, \quad \text{with} \quad Y = B - S = \frac{\lambda^8}{\sqrt{3}} \quad (2.1)$$

is completely determined by two invariant amplitudes, the strange Dirac and Pauli form factors  $F_1^{(s)}$  and  $F_2^{(s)}$ :

$$\langle p' | \bar{s}\gamma_\mu s | p \rangle = \bar{N}(p') \left( \gamma_\mu F_1^{(s)}(q^2) + i \frac{\sigma_{\mu\nu} q^\nu}{2M_N} F_2^{(s)}(q^2) \right) N(p). \quad (2.2)$$

(Here  $q = p' - p$  is the momentum transfer of the current and  $N(p)$  denotes the nucleon spinor.) The above decomposition is analogous to that of the electromagnetic current matrix element. Strangeness conservation and the nucleon's zero overall strangeness charge imply, however, a different normalization,  $F_1^{(s)}(0) = 0$ , of the Dirac form factor.

---

<sup>1</sup>Note the nonstandard sign convention for the strangeness charge [28], which carries over to the sign of its hypercharge contribution.

Alternatively, the matrix element (2.2) can be described in terms of the electric and magnetic Sachs form factors:

$$\begin{aligned} G_E^{(s)}(q^2) &= F_1^{(s)}(q^2) + \frac{q^2}{4M_N^2} F_2^{(s)}(q^2), \\ G_M^{(s)}(q^2) &= F_1^{(s)}(q^2) + F_2^{(s)}(q^2). \end{aligned} \quad (2.3)$$

Due to their association with the strangeness charge and current distributions in the Breit frame the Sachs form factors have a somewhat more direct physical interpretation.

The strangeness radius  $r_s^2$  and the strangeness magnetic moment  $\mu_s$  are defined as the first nonvanishing moments of either the Dirac/Pauli or the Sachs form factors,

$$r_s^2 = 6 \frac{d}{dq^2} F_1^{(s)}(q^2)|_{q^2=0}, \quad (r_s^2)_{Sachs} = 6 \frac{d}{dq^2} G_E^{(s)}(q^2)|_{q^2=0}, \quad \mu_s = F_2^{(s)}(0) = G_M^{(s)}(0). \quad (2.4)$$

Both definitions are currently in use.

Our analysis of the strange form factors starts from the dispersion relations

$$F_i^{(s)}(q^2) = \frac{1}{\pi} \int_{s_0}^{\infty} ds \frac{Im\{F_i^{(s)}(s)\}}{s - q^2}. \quad (2.5)$$

Subtraction terms are omitted since they play no role in the following discussion. The singularities of the form factors are located above the three-pion threshold (in the limit of good G-parity), i.e. at real, time-like  $q^2 \geq s_0 = (3m_\pi)^2$ . The spectral functions  $\pi^{-1} Im\{F_i^{(s)}\}$  receive contributions from all on-shell intermediate states with  $I^G J^{PC} = 0^- 1^{--}$  through which the strangeness current couples to the nucleon.

Our aim in the remainder of this paper will be to construct N-pole approximations

$$\frac{1}{\pi} Im\{F_i^{(s)}(s)\} = \sum_{v=1}^N B_i^{(v)} m_v^2 \delta(s - m_v^2), \quad i \in \{1, 2\}, \quad (2.6)$$

for the spectral densities, i.e. to represent the intermediate states by sharp vector meson poles. This ansatz provides an excellent description of the two lowest-lying resonances in the isoscalar channel, the  $\omega$  and  $\phi$  mesons, with their respective widths of 8 and 4 MeV. The additional poles summarize strength from higher-lying resonances and from the multi-particle continuum.

The  $N$ -pole ansatz (2.6) contains  $3N$  a priori undetermined mass and coupling parameters. It is crucial for the reliability of the dispersion analysis that these parameters, and in particular those of the low-lying poles, are determined as accurately as possible. Fortunately, the couplings and masses of the  $\omega$  and  $\phi$  as well as the mass of the third pole can be estimated model-independently from experimental input, as noted by Jaffe [28]. This estimate relies on the fact that the isoscalar electromagnetic current  $J_\mu^{(I=0)} = J_\mu^Y/2$  shares the quantum numbers of the strange current and thus couples through the same intermediate states to the nucleon.

Since the isoscalar form factors  $F_i^{(I=0)}$  have been measured in a large momentum range ( $0 \leq Q^2 = -q^2 \leq 30 \text{ GeV}$ ) and are well fitted [30–32] by a dispersive 3-pole ansatz analogous to (2.6), the masses and couplings of these three pole states in the strange form factors can be estimated from the fit parameters: The three masses  $m_1 - m_3$  in (2.6) are identified with the pole positions found in the fits (which implies in particular  $m_1 = m_\omega, m_2 = m_\phi$ ) and the strange  $\omega$  and  $\phi$  couplings<sup>2</sup>  $B_{1,2}^{(\omega,\phi)}$  are related to the fitted values of the corresponding isoscalar couplings  $A_{1,2}^{(\omega,\phi)}$  by exploiting the known flavor structure

$$\begin{aligned} |\omega\rangle &= \cos \epsilon \frac{1}{\sqrt{2}} \left( |\bar{u}\gamma_\mu u\rangle + |\bar{d}\gamma_\mu d\rangle \right) - \sin \epsilon |\bar{s}\gamma_\mu s\rangle, \\ |\phi\rangle &= \sin \epsilon \frac{1}{\sqrt{2}} \left( |\bar{u}\gamma_\mu u\rangle + |\bar{d}\gamma_\mu d\rangle \right) + \cos \epsilon |\bar{s}\gamma_\mu s\rangle, \end{aligned} \quad (2.7)$$

of the  $\omega$  and  $\phi$  poles. (The physical flavor wave functions deviate only slightly from the ideally mixed states, i.e. the mixing angle  $\epsilon = 0.053$  [33] is small.) The couplings  $g(V, J)$  of the vector mesons  $V = \omega, \phi$  to the neutral currents  $J = J^{(I=0)}, J^{(s)}$  (defined via  $\langle 0 | J_\mu | V \rangle = g(V, J) m_V^2 \varepsilon_\mu$ , where  $\varepsilon_\mu$  is the polarization vector of the  $V$ ) are determined under the assumption that the quark current of flavor  $i$  couples with universal strength  $\kappa$  exclusively to the component of the vector-meson wave function  $(\bar{q}_j q_j)_V$  with the same

---

<sup>2</sup>Depending on the context we will sometimes substitute the symbols  $\omega$  and  $\phi$  for the values  $v = 1, 2$  of the pole index.

flavor, i.e.

$$\langle 0 | \bar{q}_i \gamma_\mu q_i | (\bar{q}_j q_j)_V \rangle = \kappa m_V^2 \delta_{ij} \varepsilon_\mu. \quad (2.8)$$

This formula reproduces the empirical values of the electromagnetic coupling ratios to within a few percent. For the couplings to the neutral currents it leads to the expressions

$$\begin{aligned} g(\omega, J^{I=0}) &= \frac{\kappa}{\sqrt{6}} \sin(\theta_0 + \epsilon), & g(\omega, J^s) &= -\kappa \sin \epsilon, \\ g(\phi, J^{I=0}) &= -\frac{\kappa}{\sqrt{6}} \cos(\theta_0 + \epsilon), & g(\phi, J^s) &= \kappa \cos \epsilon, \end{aligned} \quad (2.9)$$

where  $\theta_0$  is the “magic angle” with  $\sin^2 \theta_0 = 1/3$ .

After parametrizing the vector-meson nucleon couplings as  $g_i(\omega_0, N) = g_i \cos \eta_i$ ,  $g_i(\phi_0, N) = g_i \sin \eta_i$  (the index  $i = 1, 2$  refers to the  $\gamma_\mu$  and  $\sigma_{\mu\nu} q^\nu$  couplings and  $\omega_0, \phi_0$  denote the ideally mixed states), the four couplings  $B_{1,2}^{(\omega, \phi)}$  can be related to the corresponding (fitted) isoscalar couplings  $A_{1,2}^{(\omega, \phi)}$  (which determine phenomenological values for  $\eta_i$  and  $\kappa g_i$ ) through the relations

$$A_i^{(\omega)} = \frac{\kappa g_i}{\sqrt{6}} \sin(\theta_0 + \epsilon) \cos(\eta_i + \epsilon), \quad B_i^{(\omega)} = -\kappa g_i \sin \epsilon \cos(\eta_i + \epsilon), \quad (2.10)$$

$$A_i^{(\phi)} = -\frac{\kappa g_i}{\sqrt{6}} \cos(\theta_0 + \epsilon) \sin(\eta_i + \epsilon), \quad B_i^{(\phi)} = \kappa g_i \cos \epsilon \sin(\eta_i + \epsilon). \quad (2.11)$$

The numerical values of the couplings  $B_{1,2}^{(\omega, \phi)}$  for the different fits to the electromagnetic form factors (cf. Table I) are listed in Table II.

The normalization of the Dirac form factor requires (in the given framework) at least one more pole besides the  $\omega$  and the  $\phi$ . In order to complete the construction of the spectral densities (2.6) for the minimal 3-pole ansatz we therefore have to determine two more couplings,  $B_1^{(3)}$  and  $B_2^{(3)}$ . For this purpose flavor symmetry arguments offer no help since the flavor content of the strength associated with the third pole is unknown. Instead, we fix these couplings by imposing moderate constraints (i.e. superconvergence relations) on the asymptotic behavior of the form factors,

$$\lim_{q^2 \rightarrow \infty} F_1^{(s)}(q^2) \rightarrow 0 \quad \Rightarrow \quad \sum_v B_1^{(v)} = 0, \quad (2.12)$$

$$\lim_{q^2 \rightarrow \infty} q^2 F_2^{(s)}(q^2) \rightarrow 0 \quad \Rightarrow \quad \sum_v B_2^{(v)} m_v^2 = 0. \quad (2.13)$$

The first of these relations is needed in any case since it also normalizes the Dirac form factor. Table II contains the numerical values of the third-pole couplings which follow from solving these constraints. Since the normalization condition can be used to write  $F_1^{(s)}$  with one subtraction, the final expressions for the 3-pole form factors become

$$F_1^{(s)}(q^2) = q^2 \sum_{v=1}^3 B_1^{(v)} \frac{1}{m_v^2 - q^2}, \quad (2.14)$$

$$F_2^{(s)}(q^2) = \sum_{v=1}^3 B_2^{(v)} \frac{m_v^2}{m_v^2 - q^2}. \quad (2.15)$$

In this form they were used by Jaffe [28], with parameters extracted from three of Höhler *et al.*'s fits [30] (cf. Tables I and II). The resulting form factors are plotted in Figs. 1a and 1b. Note that the overall scale of the form factors depends rather sensitively on which parameter set is used. For an estimate of the strangeness radius and magnetic moment Jaffe averaged over all three fits and obtained  $r_s^2 = (0.16 \pm 0.06) \text{ fm}^2$ ,  $(r_s^2)_{Sachs} = (0.14 \pm 0.07) \text{ fm}^2$  and  $\mu_s = -(0.31 \pm 0.09)$  (in units of the nuclear magneton). The indicated errors originate solely from differences between the fits and provide therefore at best a rough lower bound on the complete error.

The 3-pole results for the leading moments are surprisingly large, of the order of the neutron charge radius  $r_n^2 = -0.11 \text{ fm}^2$  and the nucleon's isoscalar magnetic moment  $\mu^{(I=0)} = 0.44$ , respectively. The main origin of these values can be traced to the large couplings of the  $\phi$ . As will be shown below, the dominant role of the  $\phi$  pole in the low- $s$  region of the spectral functions determines together with the asymptotic constraints most of the momentum dependence of the form factors<sup>3</sup>.

We conclude this section by noting that the simplicity of the 3-pole ansatz implies both advantages and limitations. On the one hand it requires a minimal number of parameters to be fixed and avoids the increasingly less reliable description of higher-lying strength in

---

<sup>3</sup>The results of the dispersive analysis rely therefore strongly on the identification of the second pole in the isoscalar form factor fits with the physical  $\phi$  meson.



terms of additional poles. On the other hand it leads to a dipole behavior of the form factors (see the following section) and cannot accomodate the faster decays which QCD counting rules predict at large space-like  $q^2$ . This issue will be addressed in Section IV.

### III. UPDATE OF THE 3-POLE ESTIMATE

The 3-pole estimate of the last section was based on mass and coupling parameters derived from the twenty year old Höhler fits. A recently performed new fit to the current world data set for the electromagnetic form factors by Mergell, Meissner and Drechsel (MMD) [31] permits an update of this analysis, which will be the main subject of the present section. We will also discuss some characteristic features of the resulting spectral densities and their impact on the momentum dependence of the form factors.

The MMD fits were designed to reproduce both the asymptotic power behavior<sup>4</sup> [34] of the electromagnetic form factors,  $F_i \rightarrow q^{-2(i+1)}$ , and the logarithmic QCD corrections [35]. Since the asymptotic behavior arises at least partially from continuum contributions, a more complete description should reduce the continuum contamination of the extracted pole couplings. One would expect this effect to be strongest for the third, effective pole, and indeed the MMD couplings  $A_i^{(3)}$  are substantially smaller than the average couplings  $(\bar{A}_1^{(3)})_H$  of Höhler et al.:

$$\frac{(A_1^{(3)})_{MMD}}{(\bar{A}_1^{(3)})_H} = 0.19, \quad \frac{(A_2^{(3)})_{MMD}}{(\bar{A}_2^{(3)})_H} = 0.67. \quad (3.1)$$

The accuracy of the fitted  $\omega$  and  $\phi$  couplings (which change considerably less, cf. Table I, but are of pivotal importance in the determination of the corresponding strange current couplings) should also benefit from the improved continuum description and enhance the reliability of the strange form factor analysis.

---

<sup>4</sup>One of the necessary superconvergence relations for this behavior was not imposed by Höhler *et al.*

As perhaps another consequence of the improved continuum description, the MMD fit finds the third pole at the mass of a well-established resonance, the  $\omega(1600)$ . This strengthens the rationale for adopting the same  $m_3$  in the strange form factors: if the third pole mostly summarized continuum contributions (as assumed in [30]), the response to the strange and hypercharge currents could in principle be centered at different invariant masses in the spectral functions. Still, the association with the  $\omega(1600)$  should not be taken too seriously since the quality of the data, the ill-posed fitting problem [36] and limitations of the pole ansatz do not allow a very accurate determination of the pole positions<sup>5</sup>. In the spectral functions of the strange form factors, furthermore, strength around 1600 MeV may originate from the nearby  $\phi(1680)$  resonance with its larger strangeness content. Some support for this possibility will emerge in section IV.

Concluding this brief discussion of improvements in the MMD fit which are beneficial for the strange form factor analysis, we stress that its probably most important new feature is the considerably expanded experimental data base. It consists of the current world data set for the electromagnetic form factors, which has grown, in particular for the neutron electric form factor, substantially over the last two decades since Höhler *et al.*s fits appeared.

The parameter update for the strange 3-pole form factors follows essentially the procedure of the preceding section. The three pole masses are identified with those of the MMD fits, i.e.  $m_\omega = 781$  MeV,  $m_\phi = 1019$  MeV, and  $m_3 = 1600$  MeV. An additional step is required to extract the pole couplings, since the MMD couplings

$$A_i^{(v)}(q^2) = A_i^{(v)} \frac{L(q^2)}{L(m_v^2)}, \quad L(q^2) = \left[ \ln \left( \frac{9.733 - q^2}{0.350} \right) \right]^{-2.148} \quad (3.2)$$

(in the “multiplicative” parametrization of Ref. [31]) have an effective momentum dependence devised to reproduce the logarithmic QCD corrections in the asymptotic region. However, since the  $A_i^{(v)}(q^2)$  vary by less than 10% in the low-momentum region

---

<sup>5</sup>Höhler *et al.* [30], for example, could fit the older data with comparable accuracy for values of  $m_3$  between 1400 and 1800 MeV.

$0 \leq Q^2 = -q^2 \leq 2 \text{ GeV}^2$  and hardly affect the momentum dependence of the form factors, we will use the on-shell values, i.e. the pole residua  $A_i^{(v)} \equiv A_i^{(v)}(m_v^2)$ , which are listed in Table I. (The singularity of  $L(q^2)$  at timelike  $q^2$  is an artefact of the parametrization without physical significance [31].)

The  $\omega$  and  $\phi$  couplings  $B_i^{(\omega,\phi)}$  can now be calculated from the  $A_i^{(\omega,\phi)}$  via Eqs. (2.10) and (2.11). The new values of the coupling and mixing parameters  $\kappa g_i$  and  $\eta_i$ ,

$$\tan(\eta_1 + \epsilon) = 0.4583, \quad \tan(\eta_2 + \epsilon) = 0.6137, \quad \kappa g_1 = 5.3218, \quad \kappa g_2 = -0.9294, \quad (3.3)$$

establish the relation between the isoscalar and strange current couplings. The results for the latter are given in the last rows of Table II. Together with the pole masses  $m_\omega, m_\phi$  and  $m_3$  they form the common parameter set on which all the following descriptions of the spectral functions will be based.

The two remaining couplings  $B_{1,2}^{(3)}$  of the 3-pole ansatz are again fixed by imposing the asymptotic conditions (2.12) and (2.13), and their values are also listed in Table II. Note, incidentally, that logarithmic corrections to the  $B_i^{(v)}$  (analogous to those for the isoscalar couplings (3.2)) would not improve the description of the strange form factors since the 3-pole ansatz does not even reproduce the asymptotic power behavior predicted by QCD (see below).

We are now ready to discuss the updated results. As the new values for the strangeness radius and magnetic moment we find

$$r_s^2 = 0.22 \text{ fm}^2, \quad (r_s^2)_{Sachs} = 0.20 \text{ fm}^2, \quad \mu_s = -0.26. \quad (3.4)$$

Comparison with the values of Section II shows that the square radius increases by 40 % whereas the absolute value of the magnetic moment is reduced by about 20 %. The bulk of these changes can be traced to differences in the values of Höhler's and MMD's  $\phi$ -nucleon couplings. While both fits find surprisingly large values for these couplings<sup>6</sup>, MMD's  $A_1^{(\phi)}$

---

<sup>6</sup>The large  $g_i(\phi NN)$  were interpreted as one of the first indications for OZI-rule violations in the

still exceeds those of all Höhler fits and is almost 30 % larger than their average  $(\bar{A}_1^{(\phi)})_H$ . The Pauli coupling, on the other hand, becomes smaller in the new fits,

$$\frac{(A_1^{(\phi)})_{MMD}}{(\bar{A}_1^{(\phi)})_H} = 1.28, \quad \frac{(A_2^{(\phi)})_{MMD}}{(\bar{A}_2^{(\phi)})_H} = 0.84. \quad (3.5)$$

The strange current couplings  $B_i^{(\phi)}$  move in the same directions and bring about most of the mentioned changes in the values of the moments.

It would clearly be useful to have a reliable error estimate for the updated moments. Unfortunately, however, it is practically impossible to assess the systematic errors associated with the input data, the fit procedure, the limitations of the 3-pole ansatz etc. Any error analysis would therefore necessarily be incomplete and potentially misleading. Jaffe's partial error estimate from the variance between different isoscalar form factor fits, incidentally, cannot be applied to the results (3.4) since MMD provide only one set of fit parameters.

Figs. 1a and 1b show the momentum dependence of the updated 3-pole form factors at spacelike momenta up to  $Q^2 = 1\text{GeV}^2$ , which includes the range to be probed in the Bates, CEBAF and MAMI experiments. For comparison these figures also contain the 3-pole form factors of the last section with input from Höhler *et al.*'s fits 7.1, 8.1 and 8.2.

The new Dirac form factor follows rather closely the one derived from Höhler *et al.*'s fit 8.2 (which has by far the largest  $\phi$  coupling under Höhler's fits), while it is up to twice as large as those from the fits 7.1 and 8.1. The updated Pauli form factor, on the other hand, is similar to those from both fits 8.1 and 8.2 (although the  $m_3$  of these fits differ by 400 MeV), whereas it has almost twice the size of that from fit 7.1. The average deviation of the updated form factors from the ones based on Höhler's fits is not insignificant and leads to the changed predictions (3.4).

It is instructive to compare the variations under the fits to the isoscalar *electromagnetic* form factors with the induced variations between the strange form factors. In the considered momentum range the former vary by at most a few percent while the corresponding strange

---

nucleon [37], although they might at least partially originate from  $K\bar{K}$  continuum strength [38].

form factors show much larger differences. The main source of these differences lies in the increased sensitivity of the strange form factor analysis to the  $\phi$  couplings.

Also the detailed momentum dependence of the form factors (beyond their overall scale) depends sensitively on the relative sizes of the pole couplings. The leading  $1/q^2$  dependence of the  $\phi$  pole, for example, cannot be cancelled by the  $\omega$  pole alone since the  $|B_i^{(\omega)}|$  are about an order of magnitude smaller than the  $|B_i^{(\phi)}|$ . In order to satisfy the asymptotic constraints (2.12) and (2.13), the third-pole couplings must therefore have a comparable magnitude, but the opposite sign of the  $\phi$  couplings. A glance at Table II shows that this is indeed the case:  $B_1^{(3)} \simeq -B_1^{(\phi)}$  and  $B_2^{(3)} \simeq -(1/3)B_2^{(\phi)}$ . One consequently expects a dipole behavior for  $F_2^{(s)}$  with a mass scale between  $m_2$  and  $m_3$ , and an almost perfect fit for all space-like momenta is obtained in the form

$$F_1^{(s)}(q^2) = \frac{1}{6} \frac{r_s^2 q^2}{(1 - \frac{q^2}{M_1^2})^2}, \quad (3.6)$$

$$F_2^{(s)}(q^2) = \frac{\mu_s}{(1 - \frac{q^2}{M_2^2})^2}, \quad (3.7)$$

with  $M_1 = 1.31 \text{ GeV} \simeq M_2 = 1.26 \text{ GeV}$  (for the MMD parameters, i.e. with  $r_s^2 = 5.680 \text{ GeV}^{-2}$  and  $\mu_s = -0.257$ ). The above parametrization also realizes the asymptotic behavior (2.12), (2.13) explicitly.

It is well known that an analogous “bump-dip” structure, generated by the  $\omega$  and  $\phi$  poles, leads to the approximate dipole behavior of the electromagnetic form factors. As a consequence, the dipole mass parameter lies between the  $\omega$  and  $\phi$  masses and the couplings to the third pole are about an order of magnitude smaller (cf. Table I). The strange form factors are thus much harder, and this probably rather generic feature should be kept in mind if one chooses to parametrize them in dipole form<sup>7</sup>.

We will argue in the next section that the 3-pole results probably overestimate the

---

<sup>7</sup>Sometimes Galster dipole parametrizations [39] with the same mass as in the electromagnetic form factors have been used in the literature, e.g. in [10].

strange form factors, because constraints from QCD demand a faster asymptotic decay than (2.12), (2.13). Since the slow falloff also casts doubt on the results at small momenta (see below) it would be desirable to realize at least the maximal decay power of the 3-pole ansatz by imposing the additional superconvergence conditions  $\sum B_1^{(v)} m_v^2 = 0$  and  $\sum B_2^{(v)} m_v^4 = 0$ . However, no free parameters are left to accomodate further constraints, and Table I shows that the given parameters do not conspire to satisfy them automatically:  $\sum B_1^{(v)} m_v^2 = -2.87 \text{ GeV}^2$  and  $\sum B_2^{(v)} m_v^4 = 0.72 \text{ GeV}^4$ .

#### IV. BEYOND THE 3-POLE ANSATZ

In the remainder of this paper we will discuss minimal extensions of the 3-pole ansatz which correct the asymptotic  $q^2$  dependence of the form factors in order to match the power behavior predicted by QCD counting rules [35].

Let us start with a general comment. Even if the 3-pole description of the spectral functions is incomplete since it cannot reproduce the QCD asymptotics, this need not necessarily imply an inadequate description of the form factors in the *low-momentum region* ( $Q^2 \leq 1 \text{ GeV}^2$ ) probed at Bates, MAMI and CEBAF. Indeed, the impact of the correct asymptotics on the behavior at low momenta is not *a priori* clear: it depends both on how much of this behavior is determined by just the low- $s$  strength in the spectral functions and on how exclusively the asymptotic decay originates from higher-lying strength. These two questions are of course directly related to the (at present unknown) momentum scale at which the asymptotic behavior sets in.

Moreover, even the question of *which* asymptotic behavior to implement for an optimal low-momentum description of the form factors depends on this scale. QCD dimensional counting rules [34,35] predict that elastic interactions of the strangeness current at large spacelike  $q^2$  are (up to logarithms) suppressed as  $(1/q^2)^n$ , where  $n$  corresponds to the number of hard gluon propagators needed to distribute the momentum transfer of the probe in the target nucleon. At very large  $Q^2 = -q^2$  the leading asymptotic power behavior arises from

extrinsic<sup>8</sup> radiative corrections, which renormalize the strangeness current. Thus they are suppressed by additional factors of the strong coupling  $\alpha_s$  and decay with the same powers of  $(1/q^2)$  as the electromagnetic form factors, i.e.

$$F_1^{(s)}(q^2) \rightarrow \left(\frac{-1}{q^2}\right)^2, \quad F_2^{(s)}(q^2) \rightarrow \left(\frac{-1}{q^2}\right)^3. \quad (4.1)$$

However, enforcing this behavior might not be the best choice for an optimal description of the strange form factors at *small and intermediate* momenta in the pole approximation. Alternatively, one could implement the large- $Q^2$  behavior of the intrinsic contributions, which originate from  $s\bar{s}$  admixtures to the nucleon wave function. Although the intrinsic contributions are asymptotically subleading,

$$F_1^{(s)}(q^2) \rightarrow \left(\frac{-1}{q^2}\right)^4, \quad F_2^{(s)}(q^2) \rightarrow \left(\frac{-1}{q^2}\right)^5, \quad (4.2)$$

they lack the additional  $\alpha_s$ -suppression of their extrinsic counterparts and might thus determine the momentum dependence of the form factors at intermediate spacelike  $q^2$ , i.e. before the extrinsic behavior ultimately begins to dominate. In this situation, the form factors would up to the onset of the extrinsic behavior be better described by imposing the intrinsic asymptotics (4.2). The mismatch with the extrinsic behavior at very large momenta, furthermore, would then have very little impact in the low-momentum region of interest.

An early transition to the asymptotic regime improves both the accuracy of the pole approximation and the chances for the appearance of an intrinsic region at intermediate momenta. At present, however, it is unknown at which momenta the asymptotic behavior sets in and whether intrinsic or extrinsic contributions dominate initially. In the following sections we will therefore consider both possibilities and analyze the minimal extensions of the 3-pole ansatz which implement either of the asymptotics (4.1) or (4.2).

In the pole approximation (2.6) these extensions correspond to the implementation of additional poles, which represent both higher-lying resonances and strength from multi-particle

---

<sup>8</sup>For an early discussion of intrinsic and extrinsic contributions in the context of charm quark admixtures see Ref. [2]. I am indebted to Stan Brodsky for very helpful discussions on this subject.

intermediate states like  $(2n + 1)\pi, K\bar{K}, N\bar{N}$  and  $\Lambda\bar{\Lambda}$ . An explicit inclusion of continuum cuts (in particular from the  $K\bar{K}$  states) is beyond the scope of the present paper and will be reserved for a separate investigation [40].

### A. The 4-pole ansatz with extrinsic asymptotics

The phenomenological values of the three lowest-lying masses and of the  $\phi$  and  $\omega$  couplings (as found in Section III) imply that at least four poles are required to realize the asymptotic behavior (4.1). The minimal ansatz for the form factors with extrinsic asymptotics is therefore

$$F_i^{(s)}(q^2) = \sum_{v=1}^4 B_i^{(v)} \frac{m_v^2}{m_v^2 - q^2}, \quad i \in \{1, 2\}, \quad (4.3)$$

together with the two superconvergence relations

$$\sum_{v=1}^4 B_2^{(v)} m_v^{2n} = 0, \quad n = \{1, 2\} \quad (4.4)$$

for the (unsubtracted) Pauli form factor and the normalization and superconvergence conditions

$$\sum_{v=1}^4 B_1^{(v)} m_v^{2n} = 0, \quad n = \{0, 1\} \quad (4.5)$$

for the Dirac form factor.

Since the couplings  $B_i^{(\omega, \phi)}$  and the first three masses are already fixed, there are five parameters (four couplings and one mass) left to be determined. Writing the above constraints in the form

$$\begin{pmatrix} m_3^2 & m_4^2 \\ m_3^4 & m_4^4 \end{pmatrix} \begin{pmatrix} m_3^{-2} & 0 \\ 0 & m_4^{-2} \end{pmatrix} \begin{pmatrix} B_1^{(3)} \\ B_1^{(4)} \end{pmatrix} = - \begin{pmatrix} C_1^{(3)} \\ C_1^{(4)} \end{pmatrix} \quad (4.6)$$

and

$$\begin{pmatrix} m_3^2 & m_4^2 \\ m_3^4 & m_4^4 \end{pmatrix} \begin{pmatrix} B_2^{(3)} \\ B_2^{(4)} \end{pmatrix} = - \begin{pmatrix} C_2^{(3)} \\ C_2^{(4)} \end{pmatrix} \quad (4.7)$$



$(C_1^{(i)} \equiv \sum_{j=1}^2 B_1^{(j)} m_j^{2(i-3)}, C_2^{(i)} \equiv \sum_{j=1}^2 B_2^{(j)} m_j^{2(i-2)})$  shows immediately that these equations have a unique solution for the couplings  $B_i^{(3,4)}$  as a function of the mass  $m_4$ . (The Vandermonde mass matrices are regular since the 4-pole ansatz implies  $m_3 \neq m_4$ .) This solution is

$$B_1^{(v)} = -B_1^{(1)} \prod_{i \neq v} \frac{m_1^2 - m_i^2}{m_v^2 - m_i^2} - B_1^{(2)} \prod_{i \neq v} \frac{m_2^2 - m_i^2}{m_v^2 - m_i^2}, \quad (4.8)$$

$$B_2^{(v)} = -B_2^{(1)} \frac{m_1^2}{m_v^2} \prod_{i \neq v} \frac{m_1^2 - m_i^2}{m_v^2 - m_i^2} - B_2^{(2)} \frac{m_2^2}{m_v^2} \prod_{i \neq v} \frac{m_2^2 - m_i^2}{m_v^2 - m_i^2}. \quad (4.9)$$

$(v, i \in \{3, 4\}.)$

After eliminating the  $B_i^{(3,4)}$  from the constrained 4-pole ansatz with the help of (4.8) and (4.9), one obtains the form factors as functions of the already determined masses and couplings, as well as of  $m_4$ :

$$F_1^s(q^2) = q^2 \left[ \frac{B_1^{(1)}}{m_1^2 - q^2} \prod_{i=3}^4 \frac{m_i^2 - m_1^2}{m_i^2 - q^2} + \frac{B_1^{(2)}}{m_2^2 - q^2} \prod_{i=3}^4 \frac{m_i^2 - m_2^2}{m_i^2 - q^2} \right], \quad (4.10)$$

$$F_2^s(q^2) = \frac{B_1^{(1)} m_1^2}{m_1^2 - q^2} \prod_{i=3}^4 \frac{m_i^2 - m_1^2}{m_i^2 - q^2} + \frac{B_1^{(2)} m_2^2}{m_2^2 - q^2} \prod_{i=3}^4 \frac{m_i^2 - m_2^2}{m_i^2 - q^2}. \quad (4.11)$$

These expressions for the strange form factors with extrinsic asymptotics are useful since they contain only unconstrained parameters.

Before turning to their numerical evaluation we still have to determine the position of the fourth pole. In contrast to the three low-lying poles, theoretical estimates of its mass would have to rely on uncontrolled assumptions since the 3-pole fits to the electromagnetic form factors contain no information about this pole. Relating the couplings of the third and fourth pole in a mixing model (e.g. for the  $\omega(1600)$  and the  $\phi(1680)$ ) in order to determine  $m_4$  from Eqs. (4.8) or (4.9), for example, would be unreliable since these resonances have a large width and since their flavor content can be estimated only roughly.

Alternatively, one could attempt to fix the fourth pole mass at the value of another known  $(0^- 1^{--})$  resonance. Although it becomes increasingly unreliable for higher-lying poles, a similar procedure was followed in Ref. [41] to estimate the electromagnetic form factors.

In the strangeness channel, however, not all of the five established resonances seem to be good candidates for additional poles. The  $\omega(1420)$  did not require a pole in the MMD fit<sup>9</sup> and thus appears to couple only weakly to the nucleon. Due to its dominant flavor content (as estimated from the observed partial decay widths), moreover, it will probably couple even weaker to the strangeness current. The remaining two established  $0^-1^{--}$  resonances, the  $\omega(1600)$  and the  $\phi(1680)$ , cannot be resolved in the pole approximation since their widths of 280 and 150 MeV are much larger than their mass difference. Thus only three strong and well separated centers of strength are known in the isoscalar channel. Since those are already represented by the first three poles, they offer no guidance in finding the value of  $m_4$ .

For the following numerical estimates we will therefore restrict the value of  $m_4$  only by the requirement that it should exceed the third pole mass ( $m_3 = 1.6 \text{ GeV}$ ) by at least a typical width<sup>10</sup> of  $\sim 300 \text{ MeV}$ . Under this condition neighboring strength positions do not coalesce and have to be described by separate poles. For masses  $1.9 \text{ GeV} \leq m_4 \leq \infty$  we find the strangeness radius and magnetic moment to interpolate monotonically in the ranges

$$\begin{aligned} 0.15 \text{ fm}^2 &\leq r_s^2 \leq 0.22 \text{ fm}^2, \\ 0.14 \text{ fm}^2 &\leq (r_s^2)_{Sachs} \leq 0.20 \text{ fm}^2, \\ -0.18 &\geq \mu_s \geq -0.26. \end{aligned} \tag{4.12}$$

If  $m_4$  becomes large, the fourth pole does not affect the momentum dependence of the form factors at  $Q^2 \ll m_4^2$ . For  $m_4 \rightarrow \infty$  the 4-pole form factors become identical to those from the 3-pole ansatz of Section III, which thus provide the upper bounds on  $r_s^2$  and  $|\mu_s|$  in (4.12). (To the quoted accuracy these values are reached at  $m_4 \sim 12 \text{ GeV}$ .) A fourth pole in

---

<sup>9</sup>Only the third pole of Höhler *et al.*s Fit 8.1 at 1.4 GeV gives some indication for a (small) contribution from the  $\omega(1420)$ .

<sup>10</sup>Continuum strength around these invariant masses would probably have an even broader distribution.

the region around 2 GeV, on the other hand, reduces the 3-pole predictions for  $r_s^2$  and  $|\mu_s|$  by about one third<sup>11</sup> while the signs of the form factors and moments remain unchanged.

Due to the asymptotic constraints on the 4-pole spectral densities, the size of the third-pole couplings remains comparable to that of the  $\phi$  couplings for all admissible values of  $m_4$  (cf. Eqs. (4.8), (4.9)). In view of MMD's identification of this pole with the  $\omega(1600)$  it is surprising that these couplings are much larger than those of the  $\omega(782)$  (even if growing continuum contributions to higher-lying poles should be expected). This suggests that in the strangeness channel the third pole receives significant strength from the  $\phi(1680)$  resonance, which presumably has much stronger couplings to the strange current.

If  $m_4$  lies in the region around 2 GeV even the fourth-pole couplings reach a size comparable to those of the second and third pole. In this case one expects the form factors to show a quadrupole momentum dependence. For the conservative choice  $m_4 = 1.9$  GeV (i.e. for the (in absolute magnitude) smallest 4-pole form factors with extrinsic asymptotics) we indeed find to a very good approximation

$$F_1^{(s)}(q^2) = \frac{1}{6} \frac{r_s^2 q^2}{(1 - \frac{q^2}{M_1^2})^3}, \quad (4.13)$$

$$F_2^{(s)}(q^2) = \frac{\mu_s}{(1 - \frac{q^2}{M_s^2})^3}, \quad (4.14)$$

with  $r_s^2 = 0.1482 \text{ fm}^2$ ,  $\mu_s = -0.1789$ ,  $M_1 = 1.47 \text{ GeV}$  and  $M_2 = 1.43 \text{ GeV}$ . The values of the mass parameters correspond rather closely to the weighted mean of the three dominant pole positions.

The 4-pole form factors with  $m_4 = 1.9 \text{ GeV}$  are shown in Figs. 2a and 2b. For comparison we also show the 3-pole form factors, i.e. the  $m_4 \rightarrow \infty$  limit of the 4-pole ansatz. All 4-pole form factors with  $m_4 \geq 1.9 \text{ GeV}$  lie in the band between these two bounds and are monotonically reached by increasing  $m_4$ . At  $Q^2 \sim 1 \text{ GeV}^2$  the 4-pole Dirac form factor is

---

<sup>11</sup>Even if one chooses  $m_4$  overly close to  $m_3$  in order to minimize  $r_s^2$  and  $|\mu_s|$ , their values cannot be reduced much further. For  $m_4 = 1.65 \text{ GeV}$ , e.g., one finds  $r_s^2 = 0.12 \text{ fm}^2$  and  $\mu_s = -0.15$ .

about two times smaller than that of the 3-pole ansatz; the strangeness radii differ by a factor of 0.7.

From the figures it is apparent that the weaker (and incorrect) asymptotic decay of the 3-pole bounds affects the momentum dependence of the form factors already at small and intermediate momenta. Thus the 3-pole predictions probably overestimate the strange form factors and the magnitude of their moments.

### B. The 6-pole ansatz with intrinsic asymptotics

Realizing the intrinsic decay (4.2) by an ansatz of the form (2.6) requires minimally five poles. The 5-pole ansatz, however, (which has the attractive feature that the number of its free parameters is matched by an equal number of constraints) turns out to be overconstrained: no exact solutions of the constraints exist and the approximate ones have unphysical features. (In particular, they require a pronounced additional pole in close vicinity to the  $\phi$ , but with couplings of opposite sign. A detailed discussion of the 5-pole ansatz is relegated to appendix B.)

Therefore, the minimal description of the intrinsic asymptotics in the pole approximation is given by the 6-pole ansatz

$$F_i^{(s)}(q^2) = \sum_{v=1}^6 B_i^{(v)} \frac{m_v^2}{m_v^2 - q^2}, \quad i \in \{1, 2\} \quad (4.15)$$

with the constraints

$$\sum_{v=1}^6 B_1^{(v)} m_v^{2k} = 0, \quad k = \{0, 1, 2, 3\}, \quad (4.16)$$

$$\sum_{v=1}^6 B_2^{(v)} m_v^{2n} = 0, \quad n = \{1, 2, 3, 4\}. \quad (4.17)$$

We solve these constraints by a straightforward extension of the approach described in Section IV A. The resulting expressions for the couplings and form factors are generalizations of (4.8), (4.9) and (4.10), (4.11), with the product indices now ranging up to  $N = 6$ . The

couplings are again uniquely determined as functions of the masses, but two more pole positions,  $m_5$  and  $m_6$ , are left a priori unrestricted.

As in the last section, we base the further analysis of the 6-pole form factors on the premise that higher-lying poles should have a minimal mass difference of  $\Delta m = 300$  MeV. Under this assumption the most conservative form factor estimate (i.e. the one smallest in magnitude) corresponds to the mass values  $\{m_4, m_5, m_6\} = \{1.9, 2.2, 2.5\}$  GeV. Increasing  $m_6, m_5$  and  $m_4$  up to infinity leads to monotonically increasing magnitudes of the form factors (and thus of  $r_s^2$  and  $|\mu_s|$ ) until finally, via the 5-pole and 4-pole results, the 3-pole form factors are reached as upper bounds. In the considered domain of pole positions we find the leading moments in the ranges

$$\begin{aligned} 0.089 \text{ fm}^2 &\leq r_s^2 \leq 0.22 \text{ fm}^2, \\ 0.081 \text{ fm}^2 &\leq (r_s^2)_{Sachs} \leq 0.20 \text{ fm}^2, \\ -0.114 &\geq \mu_s \geq -0.26. \end{aligned} \tag{4.18}$$

Relative to the 3-pole estimates the intrinsic asymptotics can thus reduce the size of the strangeness radius and magnetic moment by up to a factor of 2.5, i.e. considerably more than the extrinsic asymptotics.

As in the case of the 3-pole and 4-pole ansätze before, we find the 6-pole form factors well fitted by the simplest multipole formulae which match their asymptotic behavior:

$$F_1^{(s)}(q^2) = \frac{1}{6} \frac{r_s^2 q^2}{(1 - \frac{q^2}{M_1^2})^5}, \tag{4.19}$$

$$F_2^{(s)}(q^2) = \frac{\mu_s}{(1 - \frac{q^2}{M_s^2})^5}, \tag{4.20}$$

with  $r_s^2 = 0.08879 \text{ fm}^2$  and  $\mu_s = -0.1136$ . For  $\{m_4, m_5, m_6\} = \{1.9, 2.2, 2.5\}$  GeV, in particular, we find  $M_1 = 1.72$  GeV (for  $Q^2 \leq 10$  GeV a somewhat better fit is obtained with  $M_1 = 1.79$  GeV) and  $M_2 = 1.66$  GeV. Similarly to the previously discussed cases, the momentum dependence of Eqs. (4.19) and (4.20) reflects the fact that all five couplings  $B_i^{(2)} - B_i^{(6)}$  are of comparable size and have alternating signs.

Figures 2a and 2b contain, in addition to the 3- and 4-pole form factors, the exact 6-pole form factors with the above set of higher-lying strength positions,  $\{m_4, m_5, m_6\} = \{1.9, 2.2, 2.5\}$  GeV. The graphs demonstrate the discussed hierarchy of magnitudes. The bands between the lower bounds from the 6-pole form factors and the upper bounds from the 3-pole form factors can be continuously covered by increasing the three highest-lying masses of the 6-pole ansatz from their minimal values to infinity.

## V. GENERIC ASPECTS OF THE LEADING MOMENTS

The leading nonvanishing moments characterize the low-momentum behavior of the strange vector form factors and set their scale. They will also be the first to be measured in the forthcoming experiments, and for both reasons they are the focus of most current theoretical work in the strangeness vector channel.

Despite these efforts, however, no consensus on the size or even on the signs of these moments has been reached. Nucleon model predictions, in particular, involve large theoretical uncertainties and led to both positive and negative signs for the strangeness radius<sup>12</sup>. Model-independent information on the sign of the moments would thus be very useful. Moreover, it could give valuable hints towards the dynamical origin of the nucleon's strangeness distribution.

A kaon cloud (e.g. a  $K - \Lambda$  component in the nucleon wave function [21,22,24]), for example, generates a negative contribution to the radius. Neglecting recoil effects, this can be qualitatively understood from the fact that the kaon is less than half as heavy as the  $\Lambda$  (or any other hyperon). Its strangeness distribution thus reaches out farther from the nucleon's center of mass, and its (in our convention) negative strangeness charge determines the sign

---

<sup>12</sup>The mentioned model calculations find, with the exception of Ref. [23], a negative sign for the strange magnetic moment. Hadron models do agree better on the value of  $\mu_s$  than on that of  $r_s^2$ , probably because the moment is less sensitive to details of the strangeness distribution.

of  $r_s^2$ . An analogous argument (with the even lighter pions in the cloud) has been used to explain the negative charge radius of the neutron. Garvey *et al.*'s reanalysis of the BNL neutrino scattering data [10], incidentally, seems to favor a negative sign of the strangeness radius, although the statistics of the data is too poor to reach a reliable conclusion. The majority of the present nucleon model calculations also find a negative  $r_s^2$ .

All our dispersive estimates, on the other hand, led (independently of the number of poles involved and of the required asymptotic behavior) to the opposite, positive sign of the strangeness radius. This (as well as the negative sign of  $\mu_s$ ) is a rather generic property of the pole approximation, which can be read off from the general N-pole expressions for the lowest moments,

$$r_s^2 = \sum_v^N \frac{B_1^{(v)}}{m_v^2}, \quad \mu_s = \sum_v B_2^{(v)}. \quad (5.1)$$

These expressions imply that the signs of the characteristically large  $\phi$  couplings (which are positive (negative) in the Dirac (Pauli) form factor) determine the signs of the leading moments. The impact of the higher pole couplings is attenuated by their alternating sign (as required by the superconvergence relations) and by the mass factor in the expression for  $r_s^2$ . This explains also why we find the signs of the moments to be independent of the number of higher-lying poles.

The opposite signs of the pole and kaon-cloud predictions for  $r_s^2$  might point towards missing physics in either framework. A positive sign from experiment could indicate, for example, that the  $K - \Lambda$  intermediate states – although they are the lightest accessible ones – do not provide the main contribution to the nucleon's strangeness distribution. Indeed, their impact could be overcompensated by the more numerous intermediate states with higher mass in which the negative strangeness is carried by the heavier particle. This issue is currently under investigation [42].

A negative sign of the strangeness radius (due to the kaon cloud contribution or any other mechanism), on the other hand, would expose a serious shortcoming of the pole approximation, probably due to the neglect of continuum contributions. Indications in this direction

come from the generalized vector meson dominance framework of Ref. [24], which emphasizes the importance of the  $K\bar{K}$  continuum in addition to the  $\omega$  and  $\phi$  poles. In this approach the kaonic intermediate states are consistently incorporated through extended vector–meson nucleon vertices which describe the intrinsic strangeness distribution of the nucleon. Despite the large and positive Dirac coupling of the  $\phi$  these intrinsic contributions turn the sign of  $r_s^2$  negative. This finding suggests that the role of  $K\bar{K}$  continuum states in the dispersive analysis should be examined explicitly [40].

## VI. SUMMARY AND CONCLUSIONS

We have analyzed the nucleon’s strange vector form factors in a dispersive approach which circumvents dynamical model calculations and relies instead on experimental input from isoscalar electromagnetic form factor data. We emphasize in particular the impact of QCD-imposed constraints on the asymptotic behavior of the form factors.

All intermediate states through which the strangeness current couples to the nucleon, including higher-lying resonance and continuum contributions, are described in the pole approximation, i.e. by isoscalar vector meson states of zero width. This amounts to a generalization of the vector meson dominance principle which successfully accounts for electromagnetic interactions of hadrons. Because of its largely generic character we expect this approach to be a useful starting point also for estimates of the strange form factors.

After updating the results of Jaffe’s minimal 3–pole ansatz with input from a new fit to the world data set on the electromagnetic form factors, we extend the pole approximation in order to implement the asymptotic momentum dependence which QCD counting rules predict. In the following we collect and discuss our main results and comment on some directions for their refinement:

- 1) The update of Jaffe’s 3-pole analysis on the basis of the current world data set for the electromagnetic form factors leads to a by 40 % increased strangeness radius and a by 20 % reduced magnitude of the strange magnetic moment:



$$r_s^2 = 0.22 \text{ fm}^2, \quad (r_s^2)_{Sachs} = 0.20 \text{ fm}^2, \quad \mu_s = -0.26.$$

The considerably extended data base of the new fits and their more complete description of the isoscalar form factor asymptotics should improve the accuracy of these results.

Since both the  $\phi$  pole and the third pole contribute with comparable weight and with opposite sign to the spectral functions, the 3-pole form factors have, to a very good approximation, dipole form. The dipole mass parameter lies between the masses of the two dominant poles at about 1.3 GeV, which makes the strange form factors considerably harder than the electromagnetic ones (with a cutoff mass of 0.84 GeV). This presumably rather generic property of the strange form factors should be kept in mind if one chooses to parametrize their momentum dependence in dipole form.

- 2) The main advantage of the 3-pole ansatz lies in its simplicity. On the other hand, it cannot describe the fast decay of the strange form factors which QCD counting rules predict at asymptotically large momenta. In order to correct this shortcoming one has to implement either the asymptotic behavior of the ultimately dominating extrinsic contributions or the faster decay of the intrinsic ones, which may determine the momentum dependence at intermediate momenta. We consider both alternatives since it is at present unclear which strategy will lead to a better description at the low and intermediate momenta of interest.

The asymptotics of the extrinsic contributions requires minimally four poles, and with the conservative choice  $m_4 = 1.9 \text{ GeV}$  for the fourth pole position one finds

$$r_s^2 = 0.15 \text{ fm}^2, \quad (r_s^2)_{Sachs} = 0.14 \text{ fm}^2, \quad \mu_s = -0.18,$$

which amounts to a reduction of the 3-pole results by about a third. Increasing the value of  $m_4$  leads to monotonically increasing values for  $r_s^2$  and  $|\mu_s|$  and, in the limit  $m_4 \rightarrow \infty$ , back to the 3-pole results.

Minimally six poles are needed in order to realize the intrinsic asymptotics, and for a similarly conservative choice  $\{m_4, m_5, m_6\} = \{1.9, 2.2, 2.5\}$  GeV of the higher pole positions one obtains

$$r_s^2 = 0.089 \text{ fm}^2, \quad (r_s^2)_{Sachs} = 0.081 \text{ fm}^2, \quad \mu_s = -0.114.$$

Compared to the 3-pole results the magnitude of the moments decreases by about a factor 2.5, i.e. considerably more than in the case of the extrinsic asymptotics. Again the 3-pole results can be recovered by shifting the higher-lying pole masses to infinity.

The above results mark out the range of values for the leading moments which are accessible in the pole approximation. The predictions for  $r_s^2$  are more sensitive to the momentum dependence of the form factor and thus probably less reliable than those for  $\mu_s$ . The momentum dependence of the 3-, 4- and 6-pole form factors is well fitted by the simplest multipole parametrizations which match their asymptotic behavior.

- 3) For several reasons it does not seem useful to extend the pole approximation beyond the minimal 4(6)-pole ansätze considered above. While each new pole introduces three additional and a priori undetermined parameters, there are no further asymptotic constraints which could be used to fix them. Moreover, the zero-width approximation becomes less reliable for higher-lying strength and the impact of additional poles on the low-momentum behavior of the form factors would decrease since no further superconvergence conditions keep their couplings large.
- 4) The 3-pole ansatz yields the (in absolute magnitude) largest possible strange form factors in the pole approximation. Due to the unrealistic asymptotics, however, it probably overestimates the size of their leading moments by up to a factor 2.5. It is quite remarkable that the QCD-prescribed modifications of the asymptotic behavior can have such a strong impact on the low-momentum predictions.
- 5) In all the considered pole-ansatz variants the couplings to the third pole turn out to be an order of magnitude larger than the corresponding couplings in the isoscalar

electromagnetic form factors. The reason is that the third pole has to balance the strong  $\phi(1019)$  pole in order to achieve even the modest asymptotic decay of the 3-pole form factors (where no superconvergence constraint is imposed on the normalized Dirac form factor). The third-pole couplings are also four to eight times larger than the strange couplings to the  $\omega(781)$ . It seems therefore unlikely that the third pole can be identified (as in the electromagnetic case) with the  $\omega(1600)$ , which has a mostly nonstrange flavor content. An alternative and more convincing provenance for at least part of this strength could be the neighboring  $\phi(1680)$ . (Of course, both resonances cannot be distinguished in a pole ansatz since their widths are much larger than their mass difference.) If correct, this hypothesis would imply that the two established  $\phi$  resonances alone determine to a large part the low-momentum behavior of the strange vector form factors.

- 6) The signs of the leading form-factor moments turn out to be independent of the number of poles and of the implemented asymptotics: in the pole approximation they follow those of the large  $\phi$  couplings and yield a *positive* square strangeness radius and a negative strange magnetic moment. These results underline once more the crucial importance of the  $\phi(1019)$  resonance for the behavior of the vector form factors.

Most model calculations agree with the negative sign of the strange magnetic moment (see Table III), but the majority of them predicts the opposite, negative sign for the strangeness radius. This might point towards the relevance of the so far neglected  $K\bar{K}$  intermediate states in the dispersive analysis. On the basis of its large strangeness content and comparatively small invariant mass, the  $K\bar{K}$  continuum can in any case be expected to generate a significant low-energy cut. Its inclusion into the dispersion analysis is in progress [40]. Other relevant contributions to the spectral densities could come from the  $(2n+1)\pi, N\bar{N}, \Lambda\bar{\Lambda}$  (and possibly higher-lying) continuum cuts. An explicit estimate of these contributions could test if such cuts are sufficiently well accounted for by zero-width poles, as we have assumed.

Our main intent in the present study was to examine the impact of the strange form factor asymptotics on the low-momentum predictions of the dispersive analysis. In particular, we determined the range of values for the leading moments (in pole approximation) which are consistent with our present knowledge of the QCD asymptotics. Even if this range remains rather large, some further and more general conclusions can be drawn from our analysis. The most important ones are probably (i) that the third pole plays (independently of the required asymptotics) a much more important role than in the isoscalar electromagnetic form factors, (ii) that its interplay with the  $\phi(1019)$  makes the strange form factors harder than the electromagnetic ones and (iii) that the signs of the leading moments, which originate mainly from the  $\phi$  pole, are generic in the given framework and might point towards limitations of the pole approximation. Our findings suggest that the study of kaonic continuum states is the most promising direction for further extensions of the dispersive strange form factor analysis.

## ACKNOWLEDGMENTS

I would like to thank Stan Brodsky for helpful correspondence on the asymptotic behavior of the strange form factors and Wolfram Weise for an interesting discussion. Financial support by the European Community through the HCM programme is also gratefully acknowledged.

## APPENDIX A: RATIONALIZED N-POLE FORM FACTORS

In this appendix we collect several useful formulae which are encountered in deriving the results of the preceding sections. We consider a general  $N$ -pole ansatz

$$F_i^{(s)}(q^2) = \sum_{v=1}^N B_i^{(v)} \frac{m_v^2}{m_v^2 - q^2}, \quad (\text{A1})$$

for the form factors and impose the normalization and superconvergence relations (SCRs)

$$\sum_{v=1}^N B_i^{(v)} m_v^{2n} = 0, \quad n = \{0, 1, 2, \dots, k\}. \quad (\text{A2})$$

(Note that the above constraints are homogeneous and do not fix the common scale of the couplings  $B^{(v)}$ .) Of course, the form factors (A1) can satisfy only a limited number of superconvergence relations, which restricts the maximal power of  $1/q^2$  in their asymptotic decay. For the strange Pauli form factor this power is  $N$ , while the normalization condition at  $q^2 = 0$  reduces it to  $N - 1$  for the Dirac form factor.

The leading asymptotic behavior of the Dirac form factor<sup>13</sup> can be seen more explicitly by writing the normalization condition and the  $N - 1$  SCRs as a system of  $N$  linear equations  $M \cdot B = 0$  for the coefficient vector  $B$ . Since  $M$  is a  $N \times N$  Vandermonde matrix with the determinant

$$\det(M) = \prod_{v < w}^{N-1} (m_v^2 - m_w^2) \quad (\text{A3})$$

and since none of the pole masses are equal, the unique solution is  $B = 0$ , i.e. the form factor vanishes identically. A nontrivial  $F_1$  with  $F_1(0) = 0$  allows therefore maximally  $N - 2$  SCRs, and the  $N$ -pole Dirac form factor cannot decay faster than  $F_1 \rightarrow (1/q^2)^{N-1}$ .

In order to exhibit the impact of the normalization and SCR constraints on the form factors and their asymptotic behavior explicitly, one can rewrite the  $N$ -pole ansatz (A1) in rationalized form. We will list below the resulting expressions for the 3-, 4- and 5-pole form factors, which can be easily generalized to larger  $N$ . We have for the 3-pole ansatz

$$\begin{aligned} \sum_{v=1}^3 B_i^{(v)} \frac{m_v^2}{m_v^2 - q^2} = & \left( \prod_{v=1}^3 \frac{1}{m_v^2 - q^2} \right) \left\{ S_i^{(3,0)} \prod_{v=1}^3 m_v^2 + \right. \\ & \left. + (-q^2) \left[ (s^{(3,1)} S_i^{(3,1)} - S_i^{(3,2)}) \right] + (-q^2)^2 S_i^{(3,1)} \right\}, \end{aligned} \quad (\text{A4})$$

for the 4-pole ansatz

$$\sum_{v=1}^4 B_i^{(v)} \frac{m_v^2}{m_v^2 - q^2} = \left( \prod_{v=1}^4 \frac{1}{m_v^2 - q^2} \right) \left\{ S_i^{(4,0)} \prod_{v=1}^4 m_v^2 + \right.$$

---

<sup>13</sup>A similar argument can be given for the Pauli form factor.

$$\begin{aligned}
& + (-q^2) \left[ \frac{1}{2} \left[ (s^{(4,1)})^2 - s^{(4,2)} \right] S_i^{(4,1)} - s^{(4,1)} S_i^{(4,2)} + S_i^{(4,3)} \right] \\
& + (-q^2)^2 \left[ s^{(4,1)} S_i^{(4,1)} - S_i^{(4,2)} \right] + (-q^2)^3 S_i^{(4,1)} \Big\}, \tag{A5}
\end{aligned}$$

and for the 5-pole ansatz

$$\begin{aligned}
\sum_{v=1}^5 B_i^{(v)} \frac{m_v^2}{m_v^2 - q^2} &= \left( \prod_{v=1}^5 \frac{1}{m_v^2 - q^2} \right) \left\{ S_i^{(5,0)} \prod_{v=1}^5 m_v^2 + \right. \\
&+ (-q^2) \left[ \frac{1}{6} S_i^{(5,1)} \left[ (s^{(5,1)})^3 - 3s^{(5,2)} s^{(5,1)} + 2s^{(5,3)} \right] \right. \\
&- \frac{1}{2} S_i^{(5,2)} \left[ (s^{(5,1)})^2 - s^{(5,2)} \right] + S_i^{(5,3)} s^{(5,1)} - S_i^{(5,4)} \Big] \\
&+ (-q^2)^2 \left[ \frac{1}{2} S_i^{(5,1)} \left[ (s^{(5,1)})^2 - s^{(5,2)} \right] - S_i^{(5,2)} s^{(5,1)} + S_i^{(5,3)} \right] \\
&\left. \left. + (-q^2)^3 \left[ S_i^{(5,1)} s^{(5,1)} - S_i^{(5,2)} \right] + (-q^2)^4 S_i^{(5,1)} \right\}, \tag{A6}
\end{aligned}$$

where we have introduced the abbreviations

$$S_i^{(N,n)} = \sum_{v=1}^N B_i^{(v)} m_v^{2n}, \quad s^{(N,n)} = \sum_{v=1}^N m_v^{2n}. \tag{A7}$$

Note that these expressions for the  $S_i^{(N,n)}$  appear in the superconvergence relations, so that their impact on the momentum dependence of the constrained form factors can be directly red off from Eqs. (A4) – (A6). The correctly normalized 4-pole form factors with extrinsic asymptotics, for example, satisfy the constraints (4.5) and (4.4) or in our present notation

$$S_1^{(4,m)} = 0, \quad m = \{0, 1\}; \quad S_2^{(4,n)} = 0, \quad n = \{1, 2\}, \tag{A8}$$

and thus have the rationalized form

$$\begin{aligned}
F_1^{(s)}(q^2) &= \left( \prod_{v=1}^4 \frac{1}{m_v^2 - q^2} \right) \left\{ (-q^2) \left[ -s^{(4,1)} S_1^{(4,2)} + S_1^{(4,3)} \right] - (-q^2)^2 S_1^{(4,2)} \right\}, \\
F_2^{(s)}(q^2) &= \left( \prod_{v=1}^4 \frac{1}{m_v^2 - q^2} \right) \left\{ S_2^{(4,0)} \prod_{v=1}^4 m_v^2 + (-q^2) S_2^{(4,3)} \right\}. \tag{A9}
\end{aligned}$$

In this form the cancellation of the asymptotic behavior from individual poles up to the required order and the large- $q^2$  behavior (4.1) become explicit. The rationalized expressions (A4) – (A6) confirm the above result for the maximal leading decay power of the  $N$ -pole ansatz.

## APPENDIX B: THE 5-POLE ANSATZ WITH INTRINSIC QCD ASYMPTOTICS

At least five poles are needed to realize the intrinsic asymptotic behavior (4.2) in the pole approximation. Besides being in this sense minimal, the 5-pole ansatz has the additional advantage that the number of normalization and superconvergence constraints matches the number of free parameters. A physically acceptable solution of these constraints would therefore determine the form factors without further input.

In order to provide the 5-pole ansatz

$$F_i^s(q^2) = \sum_{v=1}^5 B_i^{(v)} \frac{m_v^2}{m_v^2 - q^2}, \quad i \in \{1, 2\}, \quad (\text{B1})$$

with the correct normalization and the intrinsic asymptotics (i.e.  $\lim_{Q^2 \rightarrow \infty} Q^8 F_2^s = 0$  and  $\lim_{Q^2 \rightarrow \infty} Q^6 F_1^s = 0$ ), the constraints

$$\sum_{v=1}^5 B_2^{(v)} m_v^{2n} = 0, \quad n = \{1 - 4\}; \quad \sum_{v=1}^5 B_1^{(v)} m_v^{2n} = 0, \quad n = \{0 - 3\}. \quad (\text{B2})$$

have to be imposed. Since the 4 couplings  $B_i^{(\omega, \phi)}$  and the 3 masses  $m_\omega, m_\phi, m_3$  are already fixed at the values of Section III, 8 parameters are left to be determined from the 8 constraints (B2). To this end we first derive the couplings  $B_i^{(v)}, v \in \{3, 4, 5\}$  as a function of the masses  $m_3 - m_5$  by using the 6 constraints with the lowest mass weights and find the unique solution

$$B_1^{(v)} = -B_1^{(1)} \prod_{i \neq v} \frac{m_1^2 - m_i^2}{m_v^2 - m_i^2} - B_1^{(2)} \prod_{i \neq v} \frac{m_2^2 - m_i^2}{m_v^2 - m_i^2}, \quad (\text{B3})$$

$$B_2^{(v)} = -B_2^{(1)} \frac{m_1^2}{m_v^2} \prod_{i \neq v} \frac{m_1^2 - m_i^2}{m_v^2 - m_i^2} - B_2^{(2)} \frac{m_2^2}{m_v^2} \prod_{i \neq v} \frac{m_2^2 - m_i^2}{m_v^2 - m_i^2}. \quad (\text{B4})$$

( $v, i \in \{3, 4, 5\}$ .)

The two masses  $m_4$  and  $m_5$  in the above expressions are related by the two remaining constraints, which both have the form of cubic equations for  $m_4^2$ . Out of its 3 real solutions, two are trivial,  $m_4^2 = m_3^2$  and  $m_4^2 = m_5^2$ , and can be discarded. The remaining one gives

$$m_4^2 = m_2^2 \frac{1 + \epsilon_1 \mu^2 M(m_5^2)}{1 + \epsilon_1 M(m_5^2)} \quad \text{from} \quad \sum_{v=1}^5 B_1^{(v)} m_v^6 = 0, \quad (\text{B5})$$

and

$$m_4^2 = m_2^2 \frac{1 + \epsilon_2 \mu^4 M(m_5^2)}{1 + \epsilon_2 \mu^2 M(m_5^2)} \quad \text{from} \quad \sum_{v=1}^5 B_2^{(v)} m_v^8 = 0, \quad (\text{B6})$$

where we defined

$$M(m_5^2) \equiv \frac{(m_1^2 - m_3^2)(m_1^2 - m_5^2)}{(m_2^2 - m_3^2)(m_2^2 - m_5^2)}, \quad \mu = \frac{m_1}{m_2} = 0.767. \quad (\text{B7})$$

Either of the above equations (B5) and (B6) implies characteristic restrictions on the allowed values of  $m_4^2$  and  $m_5^2$ . Their origin can be traced to the dominance of the  $\phi$  pole over the  $\omega$  pole, which is reflected in the small coupling ratios

$$\epsilon_1 \equiv \frac{B_1^{(\omega)}}{B_1^{(\phi)}} = -0.116, \quad \epsilon_2 \equiv \frac{B_2^{(\omega)}}{B_2^{(\phi)}} = -0.086. \quad (\text{B8})$$

After expanding (B5) in  $\epsilon_1 M$ , which is possible as long as  $m_5^2$  is not too close to  $m_2^2$  (so that  $M$  is of order one), we find

$$m_4^2 = m_2^2 \left[ 1 + \epsilon_1 (\mu^2 - 1) M(m_5^2) + O((\epsilon_1 M)^2) \right]. \quad (\text{B9})$$

This equation depends only weakly on  $m_5^2$ . For  $m_5^2 > m_3^2$ , it requires  $m_4^2$  to lie very close to the asymptotic value  $m_4^2 = 1.114 m_2^2$  (where  $m_5^2 \rightarrow \infty$ ) and thus very close to the  $\phi$  pole and quite far from the next known resonance, the  $\omega(1420)$ . For  $m_5^2$  close to the  $\phi$ , on the other hand, the singularity in (B5) becomes dominant and all values of  $m_4^2$  can be found as solutions for  $m_5^2$  values  $1.072 m_2^2$ .

In other words, one of the two largest pole masses remains always close to the  $\phi$  mass, whereas the other one is practically unconstrained. Qualitatively the same conclusion can be drawn from eq. (B6), which is even more restrictive since  $|\epsilon_2| < |\epsilon_1|$  and since  $\mu$  appears in higher powers.

Inspection of the solutions for the couplings

$$B_1^{(v)} = -B_1^{(2)} \prod_{i=1,3,4,5 \neq v} \frac{m_2^2 - m_i^2}{m_v^2 - m_i^2}, \quad (\text{B10})$$

$$B_2^{(v)} = -B_2^{(2)} \frac{m_2^2}{m_v^2} \prod_{i=1,3,4,5 \neq v} \frac{m_2^2 - m_i^2}{m_v^2 - m_i^2}, \quad v \in \{3, 4, 5\}, \quad (\text{B11})$$



which follow from eqs. (B3) and (B4) after the additional constraints are implemented, clarifies the significance of this result. The above expressions show that the coupling of the pole close to the  $\phi$  approximately equals the  $\phi$  coupling with opposite sign, while the remaining two couplings stay small. The large contribution of the  $\phi$  at asymptotic momenta can therefore only be canceled (as required by the superconvergence relations) by a very closely neighboring pole.

It is worth emphasizing that this result rests on only two qualitative features of the spectral density, namely the dominance of the  $\phi$  coupling over the  $\omega$  coupling and the asymptotic decay of the form factors. Taking also  $m_3$  as a free parameter, in particular, would not change the above conclusions, except that in this case one out of *three* pole masses (i.e.  $m_3, m_4$  or  $m_5$ ) would have to stay close to  $m_\phi$ , again with an approximately equal coupling of opposite sign.

With the help of the solutions (B10) and (B11), we can now eliminate the couplings  $B_i^{(3)} - B_i^{(5)}$  from the form factors, and after rationalizing the results we obtain

$$F_1^s(q^2) = q^2 \sum_{v=1}^5 B_1^{(v)} m_v^8 \prod_{i=1}^5 \frac{1}{m_i^2 - q^2} = q^2 B_1^{(2)} \frac{\prod_{v \neq 2}^5 (m_v^2 - m_2^2)}{\prod_{v=1}^5 (m_v^2 - q^2)}, \quad (\text{B12})$$

where  $m_4^2$  and  $m_5^2$  are related by eq. (B5), and

$$F_2^s(q^2) = \sum_{i=1}^5 B_2^{(i)} \prod_{j=1}^5 \frac{m_j^2}{m_j^2 - q^2} = B_2^{(2)} m_2^2 \frac{\prod_{v \neq 2}^5 (m_v^2 - m_2^2)}{\prod_{v=1}^5 (m_v^2 - q^2)}, \quad (\text{B13})$$

where  $m_4^2$  and  $m_5^2$  are related by eq. (B6).

In order to fix the values of  $m_4$  and  $m_5$  we now have to search for simultaneous solutions of Eqs. (B5) and (B6). Unfortunately, it is straightforward to see that only trivial solutions exist:  $m_4^2 = m_1^2, m_5^2 = m_2^2$  and  $m_4^2 = m_2^2, m_5^2 = m_1^2$ . (In this case the fourth and fifth pole just cancel the  $\omega$  and  $\phi$  poles and  $B_i^{(3)} = 0$ , i.e. both form factors vanish identically.) Thus the 5-pole ansatz with intrinsic asymptotics is overconstrained, even if the value of  $m_3$  is taken as a free parameter.

Nontrivial approximate solutions of this ansatz do exist, however. To see this, note that for  $\epsilon_2 = \epsilon_1 \mu^2$  the two equations (B5) and (B6) would coincide and therefore not fix

$m_4$  and  $m_5$  individually. The actual parameter values are not too far from this situation ( $\epsilon_2 - \epsilon_1 \mu^2 = -0.0458$ ) and thus approximate solutions are possible. The one with the smallest error has  $m_5^2$  close to the  $\phi$  mass and a large value of  $m_4^2$ .

Alternatively, one could allow  $m_5$  to take slightly different values in both form factors. This is not ruled out a priori since the fifth pole summarizes the strength of higher lying resonance and continuum contributions with potentially different individual Dirac and Pauli couplings. However, none of these approximate solutions resolves the problem that neighboring strength in the spectral functions would have to balance the  $\phi$  pole. This probably makes them unphysical since the  $\phi$  meson is the only established  $0^{-}1^{-}$  resonance with a mass in the vicinity of 1 GeV and since there is no indication for comparable continuum strength of opposite sign in its neighborhood.

We have also considered a modified 5-pole ansatz with the highest two superconvergence relations (which are most sensitive to the higher lying spectral strength and thus the least reliable<sup>14</sup>) relaxed, i.e. with  $\sum B_1^{(v)} m_v^6 \neq 0$  and  $\sum B_2^{(v)} m_v^8 \neq 0$ . This leaves the two masses  $m_4$  and  $m_5$  undetermined and leads to an asymptotic decay intermediate between that of the intrinsic and extrinsic contributions.

The modified 5-pole ansatz leads to  $r_2^2$  and  $\mu_s$  values in the ranges

$$-0.14 \geq \mu_s \geq -0.26, \quad 0.11 \text{ fm}^2 \leq r_s^2 \leq 0.22 \text{ fm}^2 \quad 0.10 \text{ fm}^2 \leq (r_s^2)_{\text{Sachs}} \leq 0.20 \text{ fm}^2, \quad (\text{B14})$$

for  $m_4 \leq 1.9 \text{ GeV}$ ,  $m_5 \leq 2.2 \text{ GeV}$  to  $m_{4,5} \rightarrow \infty$ . Two further poles in the region around 2 GeV thus reduce the 3-pole results by up to a factor of two. The modified 5-pole form factors with  $m_4 = 1.9 \text{ GeV}$ ,  $m_5 = 2.2 \text{ GeV}$  are well fitted by

$$F_1^s(q^2) = \frac{1}{6} \frac{r_s^2 q^2}{(1 - \frac{q^2}{M_1^2})^4}, \quad (\text{B15})$$

$$F_2^s(q^2) = \frac{\mu_s}{(1 - \frac{q^2}{M_s^2})^4}, \quad (\text{B16})$$

---

<sup>14</sup>Some dispersion theoretical studies [43] have found that SCRs involving higher powers of the masses can overburden the pole approximation.

with  $M_1 = 1.61\text{GeV}$ ,  $M_2 = 1.54\text{GeV}$ . As expected (and confirmed by inspection of the higher-pole couplings) all poles except the first contribute significantly in this case, and the weighted average of their positions corresponds to the values of the mass parameters  $M_1$  and  $M_2$  above.

## REFERENCES

- [1] A. DeRujula, H. Georgi and S.L. Glashow, Phys. Rev. D 12, 147 (1975); N. Isgur and G. Karl, Phys. Rev. D 18, 4187 (1978).
- [2] S.J. Brodsky, C. Peterson and N. Sakay, Phys. Rev. D23, 2745 (1981).
- [3] T.P. Cheng and R. Dashen, Phys. Rev. Lett. 26, 594 (1971); T.P. Cheng, Phys. Rev. D 13, 2161 (1976).
- [4] J.F. Donoghue and C.R. Nappi, Phys. Lett. B168, 105 (1986).
- [5] J. Gasser, H. Leutwyler and M.E. Sainio, Phys. Lett. B253, 252 (1991).
- [6] L.A. Ahrens et al., Phys. Rev. D 35, 785 (1987).
- [7] J. Ashman et al., Phys. Lett. B206, 364 (1988); Nucl. Phys. B328, 1 (1989).
- [8] P. Amaundruz et al., Phys. Rev. Lett. 66, 2712 (1991); CERN-PPE/93-117; B. Adeva et al., Phys. Lett. B 302, 533 (1993); Phys. Lett. B 320, 400 (1994).
- [9] see, for example, B.L. Ioffe and M. Karliner, Phys. Lett. B 247, 387 (1990).
- [10] G.T. Garvey, W.C. Louis and D.H. White, Phys. Rev. C 48, 761 (1993).
- [11] R.D. McKeown, Phys. Lett. B219, 140 (1989).
- [12] D.H. Beck, Phys. Rev. D 39, 3248 (1989).
- [13] M.J. Musolf, T.W. Donnelly, J. Dubach, S.J. Pollock, S. Kowalski and E.J. Beise, Phys. Rep. 239, 1 (1994).
- [14] CEBAF proposal #PR-91-017, D.H. Beck, spokesperson.
- [15] CEBAF proposal #PR-91-004, E.J. Beise, spokesperson.
- [16] CEBAF proposal #PR-91-010, J.M. Finn and P.A. Souder, spokespeople.
- [17] MAMI proposal A4/1-93, D. von Harrach, spokesperson.

- [18] MIT/Bates proposal # 89-06, R. McKeon and D.H. Beck, contact people.
- [19] N.W. Park, J. Schechter and H. Weigel, Phys. Rev. D 43, 869 (1991); N.W. Park and H. Weigel, Nucl. Phys. A541, 453 (1992).
- [20] H. Weigel, R. Alkofer and H. Reinhardt, Nucl. Phys. B387, 638 (1992).
- [21] M.J. Musolf and M. Burkardt, Z. Phys. C61, 433 (1994).
- [22] T.D. Cohen, H. Forkel and M. Nielsen, Phys. Lett. B316, 1 (1993).
- [23] S. Hong and B. Park, Nucl. Phys. A561, 525 (1993).
- [24] H. Forkel, M. Nielsen, X. Jin, and T.D. Cohen, Phys. Rev. C 50, 3108 (1994).
- [25] W. Koepf and E.M. Henley, Phys. Rev. C 49, 2219 (1994).
- [26] H.-C. Kim, T. Watabe and K. Goeke, preprint RUB-TPII-11/95.
- [27] S.J. Dong and K.F. Liu, Phys. Lett. B328, 130 (1994).
- [28] R.L. Jaffe, Phys. Lett. B229, 275 (1989).
- [29] H. Forkel, Plenary talk at the International School for Nuclear Physics, Erice, 17th Course: “Quarks in Hadrons and Nuclei”, Sept. 19-27, 1995, Prog. Part. Nucl. Phys. **36**, 229 (1996).
- [30] G. Höhler et al., Nucl. Phys. B114, 505 (1976).
- [31] P. Mergell, U.-G. Meissner and D. Drechsel, Mainz and Bonn University preprint TK 9515, MKPH-T-95-07, hep-ph/9506375
- [32] G. Höhler,  $\pi - N$  Newsletter 9, 108 (1993).
- [33] P. Jain et al., Phys. Rev. D 37, 3252 (1988).
- [34] S.J. Brodsky and G.R. Farrar, Phys. Rev. D11, 1309 (1975).
- [35] S.J. Brodsky and G.P. Lepage, Phys. Rev. D22, 2157 (1980).

- [36] S. Ciulli, C. Pomponiu and I. Sabba Stefanescu, Phys. Rep. 17, 133 (1975); I. Sabba Stefanescu, J. Math. Phys. 21, 175 (1980).
- [37] H. Genz and G. Höhler, Phys. Lett. B61, 389 (1976).
- [38] M.F. Gari and W. Krümpelmann, Phys. Lett. B274, 159 (1992).
- [39] S. Galster et al., Nucl. Phys. B32, 221 (1971).
- [40] H. Forkel, in preparation
- [41] S. Dubnicka, A.Z. Dubnickova and P. Strizenec, Dubna preprint E2-92-520 (1992); S.I. Bilenkaya et al., Nuovo Cim. A 105 (1992) 1421.
- [42] H. Forkel, M.J. Musolf and M. Nielsen, in preparation
- [43] N.M. Queen and G. Violini, Dispersion Theory in High-Energy Physics, John Wiley, New York, 1974.

TABLE I. Fit parameters from Refs. [30] and [31] with  $F_1^{I=0}(q^2) = \sum_v A_1^{(v)} m_v^2 / (m_v^2 - q^2)$ ,  $F_2^{I=0}(q^2) = \sum_v A_2^{(v)} m_v^2 / (m_v^2 - q^2)$ . For the (weakly momentum-dependent, see section III) couplings of the MMD fits we list the on-shell values.

Fit				
8.1	$m_v^2$ (GeV <sup>2</sup> )	0.61	1.04	1.96
	$A_1^{(v)}$	1.13	-0.52	-0.11
	$A_2^{(v)}$	-0.23	0.19	-0.036
8.2	$m_v^2$ (GeV <sup>2</sup> )	0.61	1.04	3.24
	$A_1^{(v)}$	1.17	-0.62	-0.040
	$A_2^{(v)}$	-0.18	0.13	-0.0062
7.1	$m_v^2$ (GeV <sup>2</sup> )	0.61	1.04	2.79
	$A_1^{(v)}$	1.12	-0.53	-0.086
	$A_2^{(v)}$	-0.26	0.24	-0.029
MMD	$m_v^2$ (GeV <sup>2</sup> )	0.611	1.039	2.560
	$A_1^{(v)}$	1.223	-0.711	-0.0149
	$A_2^{(v)}$	-0.200	0.156	-0.0159

TABLE II. Mass and coupling parameters in the 3-pole strange form factors  $F_1^{(s)}(q^2) = q^2 \sum_v B_1^{(v)}/(m_v^2 - q^2)$ ,  $F_2^{(s)}(q^2) = \sum_v B_2^{(v)}m_v^2/(m_v^2 - q^2)$ , as obtained from Höhler et al. [30] (Fits 8.1, 8.2, 7.1) and from Mergell, Meissner and Drechsel (MMD) [31].

fit number				
8.1	$m_v^2$ (GeV <sup>2</sup> )	0.61	1.04	1.96
	$B_1^{(v)}$	-0.24	1.62	-1.38
	$B_2^{(v)}$	0.048	-0.60	-0.30
8.2	$m_v^2$ (GeV <sup>2</sup> )	0.61	1.04	3.24
	$B_1^{(v)}$	-0.24	1.92	-1.67
	$B_2^{(v)}$	0.038	-0.39	0.12
7.1	$m_v^2$ (GeV <sup>2</sup> )	0.61	1.04	2.79
	$B_1^{(v)}$	-0.23	1.65	-1.41
	$B_2^{(v)}$	0.055	-0.75	0.27
MMD	$m_v^2$ (GeV <sup>2</sup> )	0.611	1.039	2.560
	$B_1^{(v)}$	-0.256	2.214	-1.958
	$B_2^{(v)}$	0.0420	-0.486	0.187



TABLE III. Theoretical estimates of the strangeness radius and magnetic moment

Source	$\mu_s$ ( $\mu_N$ )	$r_s^2$ ( $10^{-2} \text{ fm}^2$ )	$(r_s^2)_{Sachs}$ ( $10^{-2} \text{ fm}^2$ )	Ref.
3 poles	$-0.31 \pm 0.09$	16	$14 \pm 0.07$	[28]
3 poles, updt.	$-0.26$	22	20	this work
4 poles, extr.	$-(0.18 - 0.26)$	$15 - 22$	$14 - 20$	this work
6 poles, intr.	$-(0.11 - 0.26)$	$8.9 - 22$	$8.1 - 20$	this work
Kaon-loops & SU(3)	$-(0.31 - 0.40)$	$-(0.67 - 0.59)$	$-(2.71 - 3.23)$	[21]
Kaon-loops	$-(0.24 - 0.32)$	$-(0.67 - 0.69)$	$-(2.23 - 2.76)$	[24,22]
Gen. VMD	$-(0.24 - 0.32)$	$-(2.43 - 2.45)$	$-(3.99 - 4.51)$	[24,22]
CQ-NJL			1.69	[24]
SU(3) Skyrme	$-0.13$	$-10$	$-11$	[19]
Bos. NJL I	$-(0.05 - 0.25)$	$-(10 - 20)$	$-(10 - 22)$	[20]
Bos. NJL II	$-0.45$	$-17$	$-20$	[26]

FIG. 1a. The strange Dirac form factor from the 3-pole ansatz with parameters (cf. Table II) based on Fit 8.1 (dashed), Fit 7.1 (dot-dashed), and Fit 8.2 (dotted) of Höhler et al. [30], and on the fit of MMD [31] (full line).

FIG 1b. The strange Pauli form factor from the 3-pole ansatz with lines defined as in Fig. 1.

FIG 2a. The strange Dirac form factor from the 3-pole (full line) ansatz, the 4-pole ansatz with extrinsic asymptotics (dotted line) and the 6-pole ansatz with intrinsic asymptotics (dashed line).

FIG 2b. The strange Pauli form factor from the 3-pole, 4-pole and 6-pole ansätze in the same notation as in FIG 2a.







