

IJHET 335

May 1996

CPT, STRINGS, AND NEUTRAL MESONS

V A . KOSTELECKY

Physics Department, Indiana University, Bloomington, IN 47405, U.S.A.

A sketch of our results on CPT and strings is given. A mechanism for spontaneous CPT violation in string theory is briefly reviewed, and recent theoretical progress is summarized. Possible CPT-violating contributions to a four-dimensional low-energy effective theory could produce effects at levels testable in neutral-meson systems. Likely experimental signatures and attainable bounds are outlined in candidate experiments with correlated and uncorrelated neutral mesons.

CPT invariance is a fundamental theoretical property of local relativistic field theories of point particles.¹ Moreover, interferometric experiments with neutral kaons have tested it to a high degree of precision.² These features make CPT violation an ideal candidate signature for new, non-particle physics.³

Currently, string theory is among the most promising frameworks for a unified description of the fundamental forces. Since strings are extended objects, they do not automatically satisfy the axioms leading to CPT invariance. Indeed, an explicit stringy mechanism for CPT breaking is known,³ connected to spontaneous Lorentz breaking.⁴ Briefly, it turns out that couplings in string field theory exist that are impossible in conventional four-dimensional renormalizable gauge theories. When scalars acquire expectation values, these couplings can induce instabilities in effective potentials for Lorentz tensors. Nonzero expectation values for the latter can ensue, leading under suitable circumstances to CPT violation.

Additional support for this scenario is provided by a recent detailed analysis of Lorentz- and CPT-breaking solutions of the equations of motion for the open bosonic string.⁵ Dominant terms of the action in a level-truncation scheme are computed analytically, and the equations of motion are solved. For some solutions, over 20,000 nonvanishing terms in the action are used. The associated nonzero expectation values for various tensor fields are found to exhibit predicted features.

If a string theory indeed describes the Universe, then the above mechanism might induce CPT-violating terms in the effective four-dimensional low-energy theory, i.e., in the standard model. Since no zeroth-order CPT violation is observed, it must be suppressed. The natural suppression factor is the ratio

10^{-17} of the low-energy scale to the Planck scale. It turns out that a suppression of this magnitude can produce effects around or just below those detectable in current experiments with neutral-meson systems.^{3,6}

The issues of modeling and detecting these effects have been examined^{6;7;8} in the K system, the D system, and the two B systems. Candidate experiments include ones with uncorrelated mesons or with correlated mesons (factories). Several interesting features emerge from the analyses.

In the string scenario, the magnitude of possible CPT violation depends on coupling constants that can vary among different quark flavors as, for example, do the usual Yukawa couplings. This means it would be desirable to test for CPT violation in all the neutral meson systems, not only the K system.

The only bounds in the literature to date are those on the components of the K -system CPT-violating parameter ϵ_K . These may be improved soon, e.g., in factories, under favorable circumstances possibly to one part in 10^5 .

Testing CPT in the D system is hard because the mixing is so small. Nonetheless, definite tests can be performed at a c -charm factory. They could include bounding the K -system parameter ϵ_K and the corresponding quantity ϵ_D in the D system.

Testing CPT in the neutral B_d system is particularly interesting because it involves the b quark, and in some string scenarios CPT violation is largest in this system. Although no bound is currently in the literature, sufficient data already exist to bound the CPT-violating parameter ϵ_B at about the 10% level. Prospects at the B factories appear interesting.

My thanks to Don Colladay, Rob Potting, Stuart Samuel, and Rick Van Kooten for enjoyable collaborations leading to the results described here.

1. See, e.g., R.F. Streater and A.S. Wightman, *PCT, Spin and Statistics, and All That*, Benjamin Cummings, Reading, 1964.
2. See, e.g., Review of Particle Properties, *Phys. Rev. D* 50, 1173 (1994).
3. V.A. Kostecky and R. Potting, *Nucl. Phys. B* 359 (1991) 545.
4. V.A. Kostecky and S. Samuel, *Phys. Rev. D* 39 (1989) 683; *ibid.*, 40 (1989) 1886; *Phys. Rev. Lett.* 63 (1989) 224; *ibid.*, 66 (1991) 1811.
5. V.A. Kostecky and R. Potting, *Phys. Lett. B*, in press (hep-th/9605088).
6. V.A. Kostecky and R. Potting, *Phys. Rev. D* 51, 3923 (1995); see also V.A. Kostecky, R. Potting and S. Samuel, in S. Hegarty et al., eds., *Proc. 1991 Joint Intl. Lepton-Photon Symp. and Europhys. Conf. on High Energy Phys.* (World Scientific 1992); V.A. Kostecky and R. Potting, in D.B. Cline, ed., *Gamma Ray (Neutrino Cosmology and Planck Scale Physics)* (World Scientific 1993) (hep-th/9211116).
7. D. Colladay and V.A. Kostecky, *Phys. Lett. B* 344, 259 (1995); *Phys. Rev. D* 52, 6224 (1995).
8. V.A. Kostecky and R. Van Kooten, *Phys. Rev. D*, in press (hep-ph/9607449).