

Renormalization group analysis of the Higgs boson mass in a noncommutative differential geometry

Yoshitaka OKUMURA*

Department of Natural Science, Chubu University, Kasugai, 487, Japan

Abstract

Within one loop approximation, the renormalization group analysis of the Higgs boson mass is performed with an initial condition induced by $m_H = \sqrt{2}m_w$ which is presented in the new scheme of our noncommutative differential geometry for the reconstruction of the standard model. This initial condition holds at energy derived from $g^2 = (5/3)g'^2$ with the running $SU(2)_L$ and $U(1)_Y$ gauge coupling constants. $\sin^2\theta_w = 3/8$ is obtained under same conditions. However, contrary to $SU(5)$ GUT without supersymmetry, the grand unification of coupling constant is not realized in this scheme. The physical mass of the Higgs boson considerably depends on the top quark mass m_{top} since the top quark Yukawa coupling constants gives a large negative contribution to β function of the Higgs quartic coupling constant in wide range. The Higgs boson mass varies from 153.42GeV to 191.94GeV corresponding to $168\text{GeV} \leq m_{top} \leq 192\text{GeV}$. We find $m_H = 164.01\text{GeV}$ for $m_{top} = 175\text{GeV}$ and $m_H = 171.92\text{GeV}$ for $m_{top} = 180\text{GeV}$.

PACS number(s): 11.15.Ex,02.40.-k,12.15.-y,12.10.Kt

Keywords: spontaneously broken gauge theory; Higgs mechanism; noncommutative geometry

1 Introduction

Noncommutative differential geometry (NCG) on the discrete space such as $M_4 \times Z_2$ and, in general, $M_4 \times Z_N$ ($N = 2, 3, \dots$); Minkowski space multiplied by several points space allows one to regard the Higgs field as the gauge field on the same footing as the ordinary gauge field. A number of articles in various versions of NCG [1]-[13] have confirmed this fact by reconstructing the standard model and the grand unified theory (GUT) [3], [7], [8] without any theoretical and experimental contradictions. This is because the NCG approach to understand the Higgs mechanism apparently shows the reason for the existence of the Higgs boson field and, in addition, it does not requires any other extra physical mode except for those in the standard model and GUT.

The bosonic Lagrangian of the spontaneously broken gauge theory derived from the NCG approach has the restrictions on the gauge and the Higgs quartic coupling constants. These restrictions yield the numerical estimate of the Weinberg angle and the mass relation between the Higgs boson and other particle such as gauge boson and top quark, *though these relations hold only in tree level*. Several works have been done [11], [12] to estimate the quantum effects of these relations by assuming them to hold at some renormalization point. Shinohara, Nishida, Tanaka and Sogami [12] have recently calculated in a very clear formulation the renormalization group (RG) evolution of the mass relation $m_H = \sqrt{2}m_{top}$ for the Higgs boson and top quark, which was proposed by Sogami [13] in a variant of NCG with the generalized covariant derivative with gauge and Higgs boson fields operating on the total fermion field.

Since Connes proposed the first idea [1] concerning NCG, many versions of the NCG approach [2]-[13] have appeared to understand the Higgs mechanism, among which the present author [6], [7] has also proposed a characteristic formulation of NCG. Our formulation of a NCG is a generalization of the usual differential geometry on an ordinary manifold to the discrete manifold $M_4 \times Z_N$. The reconstruction of $SO(10)$ GUT [8] and left-right symmetric gauge model [9] had been already performed according to our NCG scheme. In a NCG on $M_4 \times Z_2$, the extra differential one-form χ in Z_2 is introduced in addition

*e-mail address: okum@isc.chubu.ac.jp

to the usual one-form dx^μ in M_4 , and therefore our formulation is very similar to ordinary differential geometry, whereas, in Connes' original one and his follower's NCG, the Dirac matrices γ_μ and γ_5 are used to describe the generalized gauge field. In a NCG, the gauge and the Higgs boson fields are given as coefficients of dx^μ and χ in the generalized connection on $M_4 \times Z_2$, respectively. However, there is no symmetry to mix dx^μ and χ , and, therefore, the ordinary gauge field can not be transformed to the Higgs boson field. In Ref.[10], three generations of fermions including left and right-handed quarks and leptons and the strong interaction are incorporated in a new scheme of our NCG to reconstruct the standard model. By setting coupling constants on the $y = +$ and $y = -$ planes with an argument $y(= \pm)$ in Z_2 to be equal, we could obtain $\sin^2 \theta_W = 3/8$ and $m_H = \sqrt{2}m_W$. Following Shinohara and co-workers [12], we will perform in this article the RG analysis of the mass relation $m_H = \sqrt{2}m_W$ for the gauge and Higgs bosons, which holds at the energy derived from $g^2 = (5/3)g'^2$ as for the running $SU(2)_L$ and $U(1)_Y$ gauge coupling constants. In other version of NCG including Sogami's approach [13], the mass relation between the Higgs boson and top quark is presented, whereas ours is the mass relation between the Higgs and charged gauge bosons. This is because the Yukawa coupling constants written in matrix form in the generation space are not contained in the generalized gauge field in our formulation.

This article consists of five sections. The next section presents the brief review of our new scheme of a NCG in Ref.[10] which incorporates the strong interaction. In the third section, the reconstruction of the standard model based on the new scheme of our NCG presented in the second section is reviewed. A characteristic point is to take the fermion field to be a vector in 24-dimensional space containing color and three generations indices. The fourth section, the main section in this paper, includes the numerical estimation of the RG evolution of the mass relation $m_H = \sqrt{2}m_W$ on the assumption that it holds at a energy of $g^2 = (5/3)g'^2$ with $SU(2)_L$ and $U(1)$ gauge coupling constants. The last section is devoted to concluding remarks.

2 Our NCG on $M_4 \times Z_2$

We review our previous formulation [10] proposed to reconstruct the standard model in which three generations of left and right-handed quarks and leptons are taken into account and also the strong interaction is incorporated. A characteristic point of this formulation is to take the left and right-handed fermions $\psi(x, y)$ with arguments x and $y(= \pm)$ in M_4 and Z_2 , respectively as

$$\psi(x, +) = \begin{pmatrix} u_L^r \\ u_L^g \\ u_L^b \\ \nu_L \\ d_L^r \\ d_L^g \\ d_L^b \\ e_L \end{pmatrix}, \quad \psi(x, -) = \begin{pmatrix} u_R^r \\ u_R^g \\ u_R^b \\ 0 \\ d_R^r \\ d_R^g \\ d_R^b \\ e_R \end{pmatrix}, \quad (2.1)$$

where subscripts L and R denote the left-handed and right-handed fermions, respectively and superscripts r , g and b represent the color indices. It should be noticed that $\psi(x, y)$ has the index for the three generation and so do the explicit expressions for fermions on the right hand sides of Eq.(2.1). In the strict expressions, u , d , ν and e in Eq.(2.1) should be replaced by

$$u \rightarrow \begin{pmatrix} u \\ c \\ t \end{pmatrix}, \quad d \rightarrow \begin{pmatrix} d \\ s \\ b \end{pmatrix}, \quad \nu \rightarrow \begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix}, \quad e \rightarrow \begin{pmatrix} e \\ \mu \\ \tau \end{pmatrix}, \quad (2.2)$$

respectively. Thus, $\psi(x, \pm)$ is a vector in the 24-dimensional space. In order to construct the Dirac Lagrangian of standard model corresponding to $\psi(x, \pm)$ in Eqs.(2.1), we must need 24-dimensional generalized covariant derivative composed of gauge and Higgs fields on $M_4 \times Z_2$. This is achieved by developing a NCG on the discrete space in the follows.

Let us start with the equation of the generalized gauge field $\mathcal{A}(x, y)$ written in one-form on the discrete space $M_4 \times Z_2$.

$$\mathcal{A}(x, y) = \sum_i a_i^\dagger(x, y) \mathbf{d}a_i(x, y) + \sum_j b_j^\dagger(x, y) \mathbf{d}b_j(x, y), \quad (2.3)$$

where $a_i(x, y)$ and $b_j(x, y)$ are the square-matrix-valued functions and are taken so as to commute with each other, because $\sum_i a_i^\dagger(x, y) \mathbf{d}a_i(x, y)$ is the flavor sector including the flavor gauge and the Higgs fields and $\sum_j b_j^\dagger(x, y) \mathbf{d}b_j(x, y)$ correspond with color gauge field. i and j are variables of the extra internal space which we can not now identify. \mathbf{d} in Eq. (2.3) is the generalized exterior derivative defined as follows.

$$\mathbf{d} = d + d_\chi, \quad (2.4)$$

$$da_i(x, y) = \partial_\mu a_i(x, y) dx^\mu, \quad (2.5)$$

$$d_\chi a_i(x, y) = [-a_i(x, y)M(y) + M(y)a_i(x, -y)]\chi, \quad (2.6)$$

$$db_j(x, y) = \partial_\mu b_j(x, y) dx^\mu \quad (2.7)$$

$$d_\chi b_j(x, y) = 0. \quad (2.8)$$

Here dx^μ is ordinary one-form basis, taken to be dimensionless, in Minkowski space M_4 , and χ is the one-form basis, assumed to be also dimensionless, in the discrete space Z_2 . If the operator ∂_y is defined as

$$\partial_y a_i(x, y) = [-a_i(x, y)M(y) + M(y)a_i(x, -y)], \quad (2.9)$$

Eq.(2.6) is rewritten as

$$d_\chi a_i(x, y) = \partial_y a_i(x, y)\chi, \quad (2.10)$$

just in the same form as in Eq.(2.5). The operator ∂_y is a difference operator on the discrete space with the modification by $M(y)$. $M(y)$ in Eq.(2.9) is required because $a_i(x, +)$ and $a_i(x, -)$ are in general square matrices with different degree and, therefore, $M(y)$ ensures the consistent calculations of matrices. ∂_y in Z_2 is an alternative of ∂_μ in M_4 . We have introduced x -independent matrix $M(y)$ whose hermitian conjugation is given by $M(y)^\dagger = M(-y)$. The matrix $M(y)$ determines the scale and pattern of the spontaneous breakdown of the gauge symmetry. Thus, Eq.(2.8) means that the color symmetry of the strong interaction does not break spontaneously.

In order to find the explicit forms of gauge and Higgs fields according to Eq.(2.3), we need the following important algebraic rule of non-commutative geometry.

$$f(x, y)\chi = \chi f(x, -y), \quad (2.11)$$

where $f(x, y)$ is a field defined on the discrete space such as $a_i(x, y)$, gauge field, Higgs field or fermion fields. It should be noticed that Eq.(2.11) does not express the relation between the matrix elements of $f(x, +)$ and $f(x, -)$ but insures the product between the fields expressed in differential form on the discrete space. Using Eq.(2.11) and some other algebraic rules in Eqs.(2.4)-(2.8), $\mathcal{A}(x, y)$ is rewritten as

$$\mathcal{A}(x, y) = A_\mu(x, y)dx^\mu + \Phi(x, y)\chi + G_\mu(x)dx^\mu, \quad (2.12)$$

where

$$A_\mu(x, y) = \sum_i a_i^\dagger(x, y)\partial_\mu a_i(x, y), \quad (2.13)$$

$$\begin{aligned} \Phi(x, y) &= \sum_i a_i^\dagger(x, y) (-a_i(x, y)M(y) + M(y)a_i(x, -y)) \\ &= a_i^\dagger(x, y) \partial_y a_i(x, y), \end{aligned} \quad (2.14)$$

$$G_\mu(x) = \sum_j b_j^\dagger(x)\partial_\mu b_j(x). \quad (2.15)$$

$A_\mu(x, y)$, $\Phi(x, y)$ and $G_\mu(x)$ are identified with the gauge field in the flavor symmetry, Higgs fields, and the color gauge field responsible for the strong interaction, respectively.

In order to identify $A_\mu(x, y)$ and $G_\mu(x)$ with true gauge fields, the following conditions have to be imposed.

$$\sum_i a_i^\dagger(x, y)a_i(x, y) = 1, \quad (2.16)$$

$$\sum_j b_j^\dagger(x)b_j(x) = \frac{1}{g_s}, \quad (2.17)$$

where g_s is a constant related to the coupling constant of the strong interaction.

Before constructing the gauge covariant field strength, we address the gauge transformation of $a_i(x, y)$ and $b_j(x)$ which is defined as

$$\begin{aligned} a_i^g(x, y) &= a_i(x, y)g_f(x, y), \\ b_j^g(x) &= b_j(x)g_c(x), \end{aligned} \quad (2.18)$$

where $g_f(x, y)$ and $g_c(x)$ are the gauge functions with respect to the flavor unitary group and the color $SU(3)_c$ group, respectively. It should be noticed that $g_c(x)$ can be taken to commute with $a_i(x, y)$ and $M(y)$ and at the same time $g_f(x, y)$ is taken to commute with $b_j(x)$. $g_f(x, y)$ and $g_c(x)$ commute with each other. Then, we obtain the gauge transformation of $\mathcal{A}(x, y)$ to be

$$\mathcal{A}^g(x, y) = g_f^{-1}(x, y)g_c^{-1}(x)\mathcal{A}(x, y)g_f(x, y)g_c(x) + g_f^{-1}(x, y)\mathbf{d}g_f(x, y) + \frac{1}{g_s}g_c^{-1}(x)dg_c(x), \quad (2.19)$$

where use has been made of Eq.(2.3) and Eq. (2.18), and as in Eqs.(2.5) and (2.6),

$$\mathbf{d}g_f(x, y) = \partial_\mu g_f(x, y)dx^\mu + \partial_y g_f(x, y)\chi \quad (2.20)$$

with the operator ∂_y defined in Eq.(2.9). Using Eqs.(2.19) and (2.20), we can find the gauge transformations of gauge and Higgs fields as

$$A_\mu^g(x, y) = g_f^{-1}(x, y)A_\mu(x, y)g_f(x, y) + g_f^{-1}(x, y)\partial_\mu g_f(x, y), \quad (2.21)$$

$$\Phi^g(x, y) = g_f^{-1}(x, y)\Phi(x, y)g_f(x, -y) + g_f^{-1}(x, y)\partial_y g_f(x, y), \quad (2.22)$$

$$G_\mu^g(x) = g_c^{-1}(x)G_\mu(x)g_c(x) + \frac{1}{g_s}g_c^{-1}(x)\partial_\mu g_c(x). \quad (2.23)$$

Equation(2.22) is very similar to two other equations and so it strongly indicates that the Higgs field is a kind of gauge field on the discrete space. From Eqs.(2.20) and (2.22) it is rewritten as

$$\Phi^g(x, y) + M(y) = g_f^{-1}(x, y)(\Phi(x, y) + M(y))g_f(x, -y), \quad (2.24)$$

which makes it obvious that

$$H(x, y) = \Phi(x, y) + M(y) \quad (2.25)$$

is un-shifted Higgs field whereas $\Phi(x, y)$ denotes shifted one with the vanishing vacuum expectation value.

In addition to the algebraic rules in Eqs.(2.4)-(2.8) we add one more important rule that

$$d_\chi M(y) = M(y)M(-y)\chi \quad (2.26)$$

which yields together with Eqs.(2.4)-(2.8) the nilpotency of the generalized exterior derivative \mathbf{d} ;

$$\mathbf{d}^2 f(x, y) = (d^2 + dd_\chi + d_\chi d + d_\chi^2)f(x, y) = 0 \quad (2.27)$$

with the condition $dx^\mu \wedge \chi = -\chi \wedge dx^\mu$. In proving the nilpotency, the following rule must be taken into account that whenever the d_χ operation jumps over $M(y)$, minus sign is attached, for example

$$\begin{aligned} d_\chi \{a(x, y)M(y)b(x, -y)\} &= (d_\chi a(x, y))M(y)b(x, -y) \\ &\quad + a(x, y)(d_\chi M(y))b(x, -y) - a(x, y)M(y)(d_\chi b(x, -y)). \end{aligned} \quad (2.28)$$

With these considerations we can construct the gauge covariant field strength.

$$\mathcal{F}(x, y) = F(x, y) + \mathcal{G}(x), \quad (2.29)$$

where $F(x, y)$ and $\mathcal{G}(x)$ are the field strengths of flavor and color gauge fields, respectively and given as

$$\begin{aligned} F(x, y) &= \mathbf{d}A(x, y) + A(x, y) \wedge A(x, y), \\ \mathcal{G}(x) &= dG(x) + g_s G(x) \wedge G(x), \end{aligned} \quad (2.30)$$

where it should be noted that $\mathbf{d}A(x, y) = \sum_i \mathbf{d}a_i^\dagger(x, y) \wedge \mathbf{d}a_i(x, y)$ and $dG(x) = \sum_j db_j^\dagger(x) \wedge db_j(x)$ are followed because of the nilpotency of \mathbf{d} and d . We can easily find the gauge transformation of $\mathcal{F}(x, y)$ as

$$\mathcal{F}^g(x, y) = g^{-1}(x, y)\mathcal{F}(x, y)g(x, y), \quad (2.31)$$

where $g(x, y) = g_f(x, y)g_c(x)$. The algebraic rules defined in Eqs.(2.4)-(2.11) and (2.26) yield

$$F(x, y) = \frac{1}{2}F_{\mu\nu}(x, y)dx^\mu \wedge dx^\nu + D_\mu\Phi(x, y)dx^\mu \wedge \chi + V(x, y)\chi \wedge \chi, \quad (2.32)$$

where

$$F_{\mu\nu}(x, y) = \partial_\mu A_\nu(x, y) - \partial_\nu A_\mu(x, y) + [A_\mu(x, y), A_\nu(x, y)], \quad (2.33)$$

$$D_\mu\Phi(x, y) = \partial_\mu\Phi(x, y) + A_\mu(x, y)(M(y) + \Phi(x, y)) - (\Phi(x, y) + M(y))A_\mu(x, -y), \quad (2.34)$$

$$V(x, y) = (\Phi(x, y) + M(y))(\Phi(x, -y) + M(-y)) - Y(x, y). \quad (2.35)$$

$Y(x, y)$ in Eq.(2.35) is auxiliary field and expressed as

$$Y(x, y) = \sum_i a_i^\dagger(x, y)M(y)M(-y)a_i(x, y), \quad (2.36)$$

which may be independent or dependent of $\Phi(x, y)$ and/or may be a constant field. In contrast to $F(x, y)$, $\mathcal{G}(x)$ is simply denoted as

$$\begin{aligned} \mathcal{G}(x) &= \frac{1}{2}G_{\mu\nu}(x)dx^\mu \wedge dx^\nu \\ &= \frac{1}{2}\{\partial_\mu G_\nu(x) - \partial_\nu G_\mu(x) + g_s[G_\mu(x), G_\nu(x)]\}dx^\mu \wedge dx^\nu. \end{aligned} \quad (2.37)$$

With the same metric structure on the discrete space $M_4 \times Z_2$ as in Ref.[6] that

$$\begin{aligned} \langle dx^\mu, dx^\nu \rangle &= g^{\mu\nu}, \quad g^{\mu\nu} = \text{diag}(1, -1, -1, -1), \\ \langle \chi, dx^\mu \rangle &= 0, \\ \langle \chi, \chi \rangle &= -1 \end{aligned} \quad (2.38)$$

we can obtain the gauge invariant Yang-Mills-Higgs lagrangian(YMH)

$$\begin{aligned} \mathcal{L}_{YMH}(x) &= -\text{Tr} \sum_{y=\pm} \frac{1}{g_y^2} \langle \mathcal{F}(x, y), \mathcal{F}(x, y) \rangle \\ &= -\text{Tr} \sum_{y=\pm} \frac{1}{2g_y^2} F_{\mu\nu}^\dagger(x, y)F^{\mu\nu}(x, y) \\ &\quad + \text{Tr} \sum_{y=\pm} \frac{1}{g_y^2} (D_\mu\Phi(x, y))^\dagger D^\mu\Phi(x, y) \\ &\quad - \text{Tr} \sum_{y=\pm} \frac{1}{g_y^2} V^\dagger(x, y)V(x, y) \\ &\quad - \text{Tr} \sum_{y=\pm} \frac{1}{2g_y^2} G_{\mu\nu}^\dagger(x)G^{\mu\nu}(x), \end{aligned} \quad (2.39)$$

where g_y is a constant relating to the coupling constant of the flavor gauge field and Tr denotes the trace over internal symmetry matrices including the color, flavor symmetries and generation space. The third term on the right hand side is the potential term of Higgs particle.

The fermion sector of the standard model was also reconstructed in Ref.[10]. However, it is not presented in this article because it is not necessary to perform the RG analysis. With these preparations, we can apply the formulation in this section to reconstruction of the Yang-Mills-Higgs Lagrangian in the standard model.

3 Reconstruction of standard model

According to a NCG in the previous section, the standard model is reconstructed. This section is also the review of Ref.[10]. The gauge fields $A_\mu(x, y)$ and $G_\mu(x)$ in the covariant derivative must be the differential representation of the fermion fields in Eqs.(2.1) and (2.2) and, therefore, they are expressed in

24 \times 24 matrices. The Higgs field $\Phi(x, y)$ is also taken to give the correct Yukawa interaction in the Dirac Lagrangian and expressed in the 24 \times 24 matrix. We specify $A_\mu(x, y)$, $\Phi(x, y)$ and $G_\mu(x)$ in Eq.(2.12) as follows:

$$A_\mu(x, +) = -\frac{i}{2} \left\{ \sum_{k=1}^3 \tau^k \otimes 1^4 A_{L\mu}^k + a B_\mu \right\} \otimes 1^3, \quad (3.1)$$

where $A_{L\mu}^k$ and B_μ are $SU(2)_L$ and $U(1)$ gauge fields, respectively and so τ^k is the Pauli matrices. 1^3 represents the unit matrix in the generation space and a is the $U(1)$ hypercharge matrix corresponding to $\psi(x, +)$ in Eq.(2.1) and expressed as

$$a = \text{diag} \left(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, -1, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, -1 \right). \quad (3.2)$$

$$A_\mu(x, -) = -\frac{i}{2} b B_\mu \otimes 1^3, \quad (3.3)$$

where b is the $U(1)$ hypercharge matrix corresponding to $\psi(x, -)$ in Eq.(2.1) and so it is 8 \times 8 diagonal matrix expressed in

$$b = \text{diag} \left(\frac{4}{3}, \frac{4}{3}, \frac{4}{3}, 0, -\frac{2}{3}, -\frac{2}{3}, -\frac{2}{3}, -2 \right). \quad (3.4)$$

$G_\mu(x)$ is denoted by

$$G_\mu(x) = -\frac{i}{2} \sum_{a=1}^8 \tau^0 \otimes \lambda'^a G_\mu^a \otimes 1^3, \quad (3.5)$$

where λ'^a is 4 \times 4 matrix made of the Gell-Mann matrix λ^a by adding 0 components to fourth line and column.

$$\lambda'^a = \begin{pmatrix} & & & 0 \\ & \lambda^a & & 0 \\ & & & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (3.6)$$

Higgs field $\Phi(x, y)$ is represented in 24 \times 24 matrix by

$$\begin{aligned} \Phi(+) &= \begin{pmatrix} \phi_0^* & \phi^+ \\ -\phi^- & \phi_0 \end{pmatrix} \otimes 1^4 \otimes 1^3, \\ \Phi(-) &= \begin{pmatrix} \phi_0 & -\phi^+ \\ \phi^- & \phi_0^* \end{pmatrix} \otimes 1^4 \otimes 1^3. \end{aligned} \quad (3.7)$$

Corresponding to Eq.(3.7), symmetry breaking function $M(y)$ is given by

$$M(+) = \begin{pmatrix} \mu & 0 \\ 0 & \mu \end{pmatrix} \otimes 1^4 \otimes 1^3, \quad M(-) = M(+)^{\dagger}. \quad (3.8)$$

It should be noted that G_μ is taken so as to commute with $A_\mu(y)$ and $\Phi(y)$. With these specifications, all quantities needed to express the explicit expression of $\mathcal{F}(x, y)$ in Eq.(2.29) can be explicitly written down as follows.

$$F_{\mu\nu}(x, +) = -\frac{i}{2} \left\{ \sum_{k=1}^3 \tau^k \otimes 1^4 F_{L\mu\nu}^k + \tau^0 \otimes a B_{\mu\nu} \right\} \otimes 1^3, \quad (3.9)$$

$$F_{\mu\nu}(x, -) = -\frac{i}{2} b B_{\mu\nu} \otimes 1^3, \quad (3.10)$$

$$G_{\mu\nu}(x) = -\frac{i}{2} \sum_{a=1}^8 \tau^0 \otimes \lambda'^a G_{\mu\nu}^a \otimes 1^3, \quad (3.11)$$

where

$$F_{L\mu\nu}^k = \partial_\mu A_{L\nu}^k - \partial_\nu A_{L\mu}^k + \epsilon^{klm} A_{L\mu}^l A_{L\nu}^m, \quad (3.12)$$

$$B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu, \quad (3.13)$$

$$G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g_s f^{abc} G_\mu^b G_\nu^c. \quad (3.14)$$

$D_\mu \Phi(x, y)$ in Eq.(2.34) is represented in

$$D_\mu \Phi(+) = (D_\mu \Phi(-))^\dagger = \left\{ \partial_\mu h' - \frac{i}{2} \left(\sum_{k=1}^3 \tau^k A_{L\mu}^k h' + h' c B_\mu \right) \right\} \otimes 1^4 \otimes 1^3, \quad (3.15)$$

where h' and c are given as

$$h' = \begin{pmatrix} \phi_0^* + \mu & \phi^+ \\ -\phi^- & \phi_0 + \mu \end{pmatrix}, \quad c = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (3.16)$$

The matrix c comes from

$$\tau^0 \otimes a - b = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \otimes 1^4 = c \otimes 1^4 \quad (3.17)$$

to insure that Higgs doublet $h = (\phi^+, \phi_0 + \mu)^t$ has plus one hypercharge and $\tilde{h} = i\tau^2 h^*$ minus one. $V(x, y)$ in Eq.(2.35) is expressed in

$$\begin{aligned} V(x, +) &= (h' h'^\dagger - \mu^2) \otimes 1^4 \otimes 1^3, \\ V(x, -) &= (h'^\dagger h' - \mu^2) \otimes 1^4 \otimes 1^3. \end{aligned} \quad (3.18)$$

It should be noticed that $Y(x, y)$ in Eq.(2.36) can be estimated by use of Eq.(3.8) to be

$$\begin{aligned} Y(x, \pm) &= \sum_i a_i^\dagger(x, \pm) M(\pm) M(\mp) a_i(x, \pm) \\ &= \mu^2 \sum_i a_i^\dagger(x, \pm) a_i(x, \pm) = \mu^2 1^{24}, \end{aligned} \quad (3.19)$$

where use has benedequationn made of Eq.(2.16).

Putting above equations into Eq.(2.39) and rescaling gauge and Higgs fields we can obtain Yang-Mills-Higgs lagrangian for the standard model as follows:

$$\begin{aligned} \mathcal{L}_{YMH} &= -\frac{1}{4} \sum_{k=1}^3 (F_{\mu\nu}^k)^2 - \frac{1}{4} B_{\mu\nu}^2 \\ &\quad + |D_\mu h|^2 - \lambda (h^\dagger h - \mu^2)^2 \\ &\quad - \frac{1}{4} \sum_{a=1}^8 G_{\mu\nu}^a \dagger G^{a\mu\nu}, \end{aligned} \quad (3.20)$$

where

$$F_{\mu\nu}^k = \partial_\mu A_\nu^k - \partial_\nu A_\mu^k + g\epsilon^{klm} A_\mu^l A_\nu^m, \quad (3.21)$$

$$B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu, \quad (3.22)$$

$$D^\mu h = \left[\partial_\mu - \frac{i}{2} \left(\sum_k \tau^k g A_{L\mu}^k + \tau^0 g' B_\mu \right) \right] h, \quad h = \begin{pmatrix} \phi^+ \\ \phi_0 + \mu \end{pmatrix}, \quad (3.23)$$

$$G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g_c f^{abc} G_\mu^b G_\nu^c, \quad (3.24)$$

with

$$g^2 = \frac{g_+^2}{12}, \quad (3.25)$$

$$g^2 = \frac{2g_+^2 g_-^2}{3g_-^2 \text{Tra}^2 + 3g_+^2 \text{Trb}^2} = \frac{g_+^2 g_-^2}{16g_+^2 + 4g_-^2}, \quad (3.26)$$

$$\lambda = \frac{g_+^2 g_-^2}{24(g_+^2 + g_-^2)}, \quad (3.27)$$

$$g_c^2 = g_s^2 \frac{g_+^2 g_-^2}{6(g_+^2 + g_-^2)}. \quad (3.28)$$

Equation(3.26) yields the Weinberg angle with the parameter $\delta = g_+/g_-$ to be

$$\sin^2 \theta_W = \frac{3}{4(\delta^2 + 1)}. \quad (3.29)$$

The gauge transformation affords the Higgs doublet h to take the form that

$$h = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ \eta + v \end{pmatrix} \quad (3.30)$$

which makes possible along with Eqs.(3.20)~(3.23) to expect the gauge boson and Higgs particle masses.

$$m_W^2 = \frac{1}{48}g_+^2v^2, \quad (3.31)$$

$$m_Z^2 = \frac{1 + \delta^2}{12(4\delta^2 + 1)}g_+^2v^2, \quad (3.32)$$

$$m_H^2 = \frac{1}{12(\delta^2 + 1)}g_+^2v^2. \quad (3.33)$$

In the limit of equal coupling constant $g_+ = g_-$, $g^2 = (5/3)g'^2$ is followed and, therefore, $\sin^2 \theta_W = 3/8$ which is the same value expected by the $SU(5)$ and $SO(10)$ GUTs. This fact makes possible to conclude that the limit of $g_+ = g_-$ yields the relations that hold at the energy derived from $g^2 = (5/3)g'^2$ with the running $SU(2)_L$ and $U(1)_Y$ gauge coupling constants. However, it should be noted that the color gauge coupling constant g_c includes an additional parameter g_s as in Eq.(3.28), and so g_c does not necessarily coincide with the $SU(2)_L$ gauge coupling constant g . Thus, contrary to $SU(5)$ GUT without supersymmetry, the grand unification of coupling constants is not realized in the present scheme. In the next section, in order to predict the physical Higgs mass, we will perform the RG analysis with the initial condition induced by $m_H = \sqrt{2}m_W$ at the energy derived from $g^2(t) = (5/3)g'^2(t)$ with the running coupling constants.

4 Numerical estimation of the Higgs mass

In order to perform the RG analysis, the renormalizable Lagrangian is necessary. We have the Yang-Mills-Higgs Lagrangian of the standard model expressed in Eq.(3.20). However, it has the special restriction on the gauge couplings and the Higgs quartic coupling which yield the mass relation for the Higgs and gauge bosons. In a sense, the restriction on coupling constants is an obstacle for renormalizability. Thus, we consider that the NCG approach provides the Lagrangian at some renormalization point. The more general Lagrangian is obtained by adding the term for the Higgs potential considered by Sitarz. He defined the new metric $g_{\alpha\beta}$ with α and β running over 0, 1, 2, 3, 4 by $g^{\alpha\beta} = \text{diag}(+, -, -, -, -)$. The fifth index represents the discrete space Z_2 . Then, $dx^\alpha = (dx^0, dx^1, dx^2, dx^3, \chi)$ is followed. The generalized field strength $F(x, y)$ in Eq.(2.22) is written by $F(x, y) = F_{\alpha\beta}(x, y)dx^\alpha \wedge dx^\beta$ where $F_{\alpha\beta}(x, y)$ is denoted in Eq.(2.23). Then, it is easily derived that $\text{Tr}\{g^{\alpha\beta}F_{\alpha\beta}(x, y)\}$ is gauge invariant. Thus, the term

$$|\text{Tr}\{g^{\alpha\beta}F_{\alpha\beta}(x, y)\}|^2 = \{\text{Tr}V(x, y)\}^\dagger \{\text{Tr}V(x, y)\} \quad (4.1)$$

can be added to Eq.(3.20). This procedure gives rise to the Lagrangian without any restriction on Higgs quartic coupling constants. In this article, we assume that this Sitarz term vanishes at a point that Eqs.(3.26)-(3.28) hold and perform the renormalization group analysis by adopting the mass relation $m_H = \sqrt{2}m_W$ as an initial condition which holds at a renormalization point giving $g_+ = g_-$ in Eq.(3.26). In this context, the way to obtain the BRST invariant Lagrangian of spontaneously broken gauge theory in NCG has already presented by Ref.[14] and [15], which includes the Fadeef-Popov ghosts and Nakanishi-Lautrup field for the gauge fixing. Thus, we can obtain the β functions for various coupling constants in the standard model just as already done.[16]

Shinohara, Nishida, Tanaka and Sogami [12] have estimated the Higgs boson mass by changing the renormalization point μ_0 at which the mass relation $m_H = \sqrt{2}m_{top}$ holds and obtained the result that m_H approaches to m_{top} from the upper side for large values of μ_0 . Following Shinohara and co-workers, the RG equations for coupling constants and the vacuum expectation value of the Higgs field are used to estimate the physical Higgs boson mass. In our case, the renormalization point at which the mass

relation $m_H = \sqrt{2}m_W$ holds is determined by looking for the point giving the relation $g^2 = (5/3)g'^2$ which results in $\sin^2\theta_W = 3/8$.

Let us start by writing the RG equations for coupling constants in the standard model. With notations

$$g_3 = g_c, \quad g_2 = g, \quad g_1 = \sqrt{\frac{5}{3}}g' \quad (4.2)$$

for $SU(3)_c$, $SU(2)_L$ and $U(1)_Y$ gauge coupling constants, respectively, the RG equations are written as

$$\mu \frac{\partial \alpha_i}{\partial \mu} = \beta_i, \quad \alpha_i = \frac{g_i^2}{4\pi} \quad (i = 1, 2, 3), \quad (4.3)$$

where [16]

$$\beta_1 = \frac{1}{2\pi} \left(\frac{4}{3}n_f + \frac{1}{10} \right) \alpha_1^2, \quad (4.4)$$

$$\beta_2 = -\frac{1}{2\pi} \left(\frac{22}{3} - \frac{3}{4}n_f - \frac{1}{6} \right) \alpha_2^2, \quad (4.5)$$

$$\beta_3 = -\frac{1}{2\pi} \left(\frac{33}{3} - \frac{4}{3}n_f \right) \alpha_3^2 \quad (4.6)$$

with the generation number n_f . The RG equation of the quartic Higgs coupling constant λ is expressed as

$$\mu \frac{\partial \alpha_H}{\partial \mu} = \beta_H, \quad \alpha_H = \frac{\lambda}{4\pi}, \quad (4.7)$$

where [16]

$$\begin{aligned} \beta_H = & \frac{6}{\pi} \alpha_H^2 - \frac{9}{20\pi} \alpha_H \alpha_1 - \frac{9}{4\pi} \alpha_H \alpha_2 + \frac{27}{800\pi} \alpha_1^2 + \frac{9}{80\pi} \alpha_1 \alpha_2 + \frac{9}{32\pi} \alpha_2^2 \\ & + \frac{1}{4\pi^2} \text{Tr} \left\{ (A_e^\dagger A_e) + 3(A_u^\dagger A_u) + 3(A_d^\dagger A_d) \right\} \alpha_H - \frac{1}{32\pi^3} \left\{ (A_e^\dagger A_e)^2 + 3(A_u^\dagger A_u)^2 + 3(A_d^\dagger A_d)^2 \right\} \end{aligned} \quad (4.8)$$

A_e , A_u and A_d in Eq.(4.8) are the Yukawa coupling constants written in 3×3 matrices for the lepton, up quark and down quark sectors, respectively. In view of the large top quark mass in the low energy region, we now assume the top quark dominance in the evaluation of traces with respect to the Yukawa coupling constants in Eq.(4.8). In this approximation, β_H is rewritten [16] as

$$\beta_H = \frac{6}{\pi} \alpha_H^2 - \frac{9}{20\pi} \alpha_H \alpha_1 - \frac{9}{4\pi} \alpha_H \alpha_2 + \frac{27}{800\pi} \alpha_1^2 + \frac{9}{80\pi} \alpha_1 \alpha_2 + \frac{9}{32\pi} \alpha_2^2 + 3 \frac{1}{\pi} \alpha_Y \alpha_H - \frac{3}{2\pi} \alpha_Y^2 \quad (4.9)$$

with the definition that

$$\alpha_Y = \frac{|(A_u)_{33}|^2}{4\pi}. \quad (4.10)$$

α_Y is subject to the following RG equation.

$$\mu \frac{\partial \alpha_Y}{\partial \mu} = \beta_Y, \quad (4.11)$$

where the β -function is given as

$$\beta_Y = \frac{\alpha_Y}{4\pi} \left(9\alpha_Y - 16\alpha_3 - \frac{9}{2}\alpha_2 - \frac{17}{10}\alpha_1 \right). \quad (4.12)$$

The RG equations for α_i ($i = 1, 2, 3$) in Eq.(4.3) are analytically solved for $n_f = 3$ and the solutions are given as

$$\alpha_1(t) = \frac{\alpha_1(0)}{1 - \alpha_1(0) \frac{41}{20\pi} t}, \quad (4.13)$$

$$\alpha_2(t) = \frac{\alpha_2(0)}{1 + \alpha_2(0) \frac{19}{12\pi} t}, \quad (4.14)$$

$$\alpha_3(t) = \frac{\alpha_3(0)}{1 + \alpha_3(0) \frac{7}{2\pi} t}, \quad (4.15)$$

where $t = \log(\mu/\mu^0)$. By use of these solutions, the RG equation for α_Y can be solved to be

$$\alpha_Y(t) = F(t) \left[\frac{\alpha_Y(0)}{1 - \alpha_Y(0) \frac{9}{4\pi} \int_0^t F(t') dt'} \right], \quad (4.16)$$

where the function $F(t)$ is defined [18] as

$$F(t) = \left(\frac{\alpha_1(t)}{\alpha_1(0)} \right)^{-17/82} \left(\frac{\alpha_2(t)}{\alpha_2(0)} \right)^{27/38} \left(\frac{\alpha_3(t)}{\alpha_3(0)} \right)^{8/7} \quad (4.17)$$

with the condition $F(0) = 1$.

Masses of all particles in the standard model are introduced by the vacuum expectation value v of the Higgs field. From Eqs.(3.26)-(3.28) and (3.31)-(3.33),

$$m_W^2 = \frac{\pi}{4} \alpha_2 v^2, \quad m_Z^2 = \frac{\pi}{4} \left(\alpha_2 + \frac{3}{5} \alpha_1 \right) v^2 \quad m_H^2 = 8\pi \alpha_H v^2 \quad (4.18)$$

are given by use of the vacuum expectation value v . Top quark mass is also expressed with v as

$$m_{top} = (A_u)_{33} \frac{v}{\sqrt{2}} = \sqrt{2\pi\alpha_Y} v. \quad (4.19)$$

In order to analyze the RG evolution of these masses, we need the RG evolution of v . It is given [16] in the following equation.

$$\mu \frac{\partial v}{\partial \mu} = -\frac{v}{4\pi} \left(3\alpha_Y - \frac{9}{4} \alpha_2 - \frac{9}{20} \alpha_1 \right) \quad (4.20)$$

with the same approximation as in the equations for coupling constants. It should be noted that all variables in Eqs.(4.18)-(4.20) depend on $t = \log(\mu/\mu^0)$.

Eq.(4.20) is analytically solved to result in

$$v(t) = v(0) \left(\frac{\alpha_Y(t)}{\alpha_Y(0)} \right)^{-1/3} \left(\frac{\alpha_1(t)}{\alpha_1(0)} \right)^{-7/492} \left(\frac{\alpha_2(t)}{\alpha_2(0)} \right)^{-9/76} \left(\frac{\alpha_3(t)}{\alpha_3(0)} \right)^{8/21} \quad (4.21)$$

with $t = \log(\mu/\mu^0)$. In these RG equations, we take the renormalization point μ_0 to be the neutral gauge boson mass $m_Z = 91.187\text{GeV}$ [17]. Thus, we must determine the gauge coupling constants in Eqs.(4.13)-(4.15) and $v(0)$ in Eq.(4.21) at $\mu = m_Z$. They takes the values [17]

$$\alpha_1(0) = 0.017, \quad \alpha_2(0) = 0.034, \quad \alpha_3(0) = 0.12 \quad (4.22)$$

and

$$v(0) = \sqrt{\frac{1}{\sqrt{2}G_F}} = 246\text{GeV}. \quad (4.23)$$

According to Eq.(4.19), the physical top quark mass m_{top} satisfies the equation

$$m_{top} = \sqrt{2\pi\alpha_Y(t_{top})} v(t_{top}), \quad (4.24)$$

where $t_{top} = \log(m_{top}/m_Z)$. This equation along with Eqs.(4.16), (4.17) and (4.22) fixes the numerical value of $\alpha_Y(0)$ which depends on the physical mass m_{top} .

Let us consider to solve the RG equation in Eq.(4.7) by adopting the mass relation $m_H = \sqrt{2}m_W$ as the initial condition. From Eqs.(3.26), (3.31) and (3.33), this mass relation leads to the relation $g_1 = g_2$ in Eq.(4.2). Then, according to Eqs.(4.13) and (4.14) together with Eq.(4.22), we can determine the renormalization point t_0 at which $m_H = \sqrt{2}m_W$ holds. The numerical value of t_0 is

$$t_0 = 25.431. \quad (4.25)$$

From Eq.(4.18), the initial condition of Eq.(4.7) is determined as

$$\alpha_H(t_0) = \frac{1}{16} \alpha_2(t_0). \quad (4.26)$$

With these considerations, we can find the running Higgs boson mass from Eq.(4.18) as

$$m_H(t) = \sqrt{8\pi\alpha_H(t)}v(t). \quad (4.27)$$

The physical Higgs boson mass m_H is given by imposing the condition that

$$m_H = \sqrt{8\pi\alpha_H(t_H)}v(t_H) \quad (4.28)$$

with $t_H = \log(m_H/m_Z)$.

Top quark mass m_{top} reported by particle data group [17] is given as

$$m_{top} = 180 \pm 12\text{GeV}. \quad (4.29)$$

We investigate the Higgs boson mass by varying the top quark mass in the range of Eq.(4.29). Fig.1 shows the running Higgs boson mass for the top quark mass 180GeV. The intersection of two functions $y = m_H(t)$ and $y = m_Z \exp(t)$ indicates the position of the physical Higgs boson mass m_H , which gives $m_H = 171.92\text{GeV}$. Fig.2 shows the relation of the Higgs boson and top quark masses in which the top quark mass varies from 168GeV to 192GeV according to the experimental data in Eq.(4.29). Corresponding to the variation of top quark mass, the Higgs boson mass varies from 153.42GeV to 191.94GeV.




Fig.1: $t = \log(\mu/m_Z)$ and a curve of $y = m_H(t)$ shows the running Higgs boson mass numerically calculated according to Eq.(4.27). An intersection of two curves indicates the position of the physical Higgs boson mass from which we can read $m_H = 171.92\text{GeV}$.




Fig.2: This figure shows the relationship between the Higgs boson and top quark masses. Corresponding to the variation of top quark mass, the Higgs boson mass varies from 153.42GeV to 191.94GeV.

5 Concluding remarks

The RG analysis has been carried out for the Higgs boson mass by adopting the initial condition induced by $m_H = \sqrt{2}m_W$ which was presented in the new scheme of NCG [10] for the reconstruction of the standard model. This mass relation together with $\sin^2\theta_W = 3/8$ holds under same condition that $g_+ = g_-$ (see Eqs.(3.25) and (3.26) which yields $g^2 = (5/3)g'^2$ with the $SU(2)_L$ and $U(1)_Y$ gauge coupling constants. However, it should be noted that the color gauge coupling constant g_c includes an additional parameter g_s as in Eq.(3.28), and so g_c does not necessarily coincide with the $SU(2)_L$ gauge coupling constant g . Thus, contrary to $SU(5)$ GUT without supersymmetry, the grand unification of coupling constants is not realized in the present scheme. In solving the RG equation we use these relations as initial conditions not inconsistent with the current observation of the grand unification of coupling constants.

The numerical mass of the Higgs boson considerably depends on the top quark mass since the top quark Yukawa coupling constants α_t gives a negative contribution to β_H in the range of $-2.0 \leq t \leq 25$ in Eq.(4.9). We have the initial condition induced by $m_H = \sqrt{2}m_W$ at $t = 25.431$ and the calculation

has performed down to $t = -2$. The larger top quark mass is, the larger Higgs boson mass we have as in Fig.2. If we adopt 180GeV as top quark mass, the Higgs boson mass becomes 171.92GeV which is smaller than that predicted by Shinohara and co-worker. If we use the data reported by Dawson [19], $m_H = 164.01\text{GeV}$ for $m_{top} = 175\text{GeV}$. Generally speaking, the NCG approach provides the Yang-Mills-Higgs Lagrangian with the restrictions on gauge and Higgs quartic coupling constants. Our mass relation $m_H = \sqrt{2}m_W$ is with respect to gauge and Higgs boson masses, whereas other version of NCG [4], [11] including Sogami's approach [13] present the mass relation between the Higgs boson and top quark. This is because the Yukawa coupling constants written in matrix form in the generation space are not contained in the generalized gauge field in our formulation. This aspect is a characteristic point of our approach compared to others. This is due to the introduction of one-form basis d^μ in M_4 and χ in Z_2 instead of γ^μ and γ^5 . This leads to the meaningful Higgs potential even in one fermion generation and the mass relation for the Higgs and gauge bosons. Therefore, we predict relatively small Higgs boson mass as seen in this article.

Calculations have performed within one loop approximation and top quark dominance is assumed in the whole range of $t = \log(\mu/m_Z)$ in the RG equations for β functions. Though there are fairly uncertainties with respect to Yukawa coupling constants for three generations of fermions, it is very interesting to analyze the RG equations for the Higgs boson mass in two loop approximation. This will be attempted in future work.

Acknowledgement

The author would like to express their sincere thanks to J. Iizuka, H. Kase, K. Morita and M. Tanaka for useful suggestion and invaluable discussions on the non-commutative geometry.

References

- [1] A. Connes, p.9 in *The Interface of Mathematics and Particle Physics*, ed. D. G. Quillen, G. B. Segal, and Tsou. S. T., Clarendon Press, Oxford, 1990. See also, Alain Connes and J. Lott, Nucl. Phys. **B** (Proc. Suppl.) **18B**, 57 (1990).
- [2] References in J. Madore and J. Mourad, "Noncommutative Kaluza-Klein Theory", hep-th/9601169.
- [3] A. H. Chamseddine, G. Felder and J. Frölich, Phys. Lett. **B296** (1992), 109; Nucl. Phys. **B395**, 672(1993); A. H. Chamseddine and J. Frölich, Phys. Rev. D **50**, 2893 (1994).
- [4] D. Kastler, Rev. Math. Phys. **5**, 477 (1993); M. Dubois-Violette, Class. Quantum. Grav. **6**, 1709 (1989); R. Coquereaux, G. Esposito-Farese, and G. Vaillant, Nucl. Phys. **B353**, 689 (1991); M. Dubois-Violette, R. Kerner, and J. Madore, J. Math. Phys. **31**, 316 (1990); B. Balakrishna, F. Gürsey and K.C. Wali, Phys. Lett. **B254**, 430 (1991); Phys. Rev. D **46**, 6498 (1992), R. Coquereaux, G. Esposito-Farése and G. Vaillant, Nucl. Phys. **B353**, 689 (1991), R. Coquereaux, G. Esposito-Farese and F. Scheck, Int. Journ. Mod. Phys., **A7**, 6555 (1992); R. Coquereaux, R. Hausling, N. Papadopoulos and F. Scheck, *ibid.* **7**, 2809 (1992).
- [5] A. Sitarz, Phys. Lett., **B308**, 311 (1993), Jour. Geom. Phys. **15**, 123 (1995).
- [6] Y. Okumura, Prog. Theor. Phys., **91**, 959 (1994).
K. Morita and Y. Okumura, Phys. Rev. D, **50**, 1016 (1994).
- [7] Y. Okumura, Phys. Rev. D, **50** (1994), 1026.
- [8] Y. Okumura, Prog. Theor. Phys., **94**, 589 (1995).
- [9] Y. Okumura and M. Morita, Nuovo Cimento A, **109A**, 311 (1996).
- [10] Y. Okumura, Prog. Theor. Phys. **95**, 969 (1996).
- [11] E. Alvarez, J.M. Gracia-Bondía and C.P. Martín, Phys. Lett., **B306**, 55 (1993); *ibid.* **B329**, 259 (1994).
- [12] T. Shinohara, K. Nishida, H. Tanaka and I.S. Sogami, "Renormalization Group Effects on the Mass Relation Predicted by the Standard Model with Generalized Covariant Derivatives, hep-ph/9606278.

- [13] I.S. Sogami, Prog. Theor. Phys. **94**,117 (1995); *ibid.* **95** (1996).
- [14] C.Y. Lee, D.S. Hwang and Y. Ne'eman, "BRST Quantization of Gauge Theory in Noncommutative Geometry: Matrix Derivative Approach", hep-th/9512215.
- [15] Y. Okumura, Phys. Rev. **D54**, 4114 (1996).
- [16] H. Arason, D.J. Castaño, B. Kesthelyi, S. Mikaelian, E.J. Piard, P. Ramond, Phys. Rev. D **46**, 3945 (1992); M.E. Machacek and M.T. Vaughn, Phys. Lett. **103B**, 427 (1981); Nucl. Phys. **B222**, 83 (1983); *ibid.* **B136**, 221 (1984); *ibid.* **B249**, 70 (1985).
- [17] Particle Data Group, R.M. Barnett *et al.*, Phys. Rev. D, **54**, 1 (1994).
- [18] G. Jungman, Phys. Rev. D **46**, 4004 (1992).
- [19] S. Dawson, "DEF '96: The triumph of the standard model", hep-ph/9609340.