

Quark Masses, Chiral Symmetry, and the $U(1)$ Anomaly

Michael Creutz

Physics Department, Brookhaven National Laboratory, Upton, NY 11973; email:
creutz@bnl.gov

I discuss the mass parameters appearing in the gauge theory of the strong interactions, concentrating on the two flavor case. I show how the effect of the CP violating parameter is simply interpreted in terms of the state of the ether via an effective potential for meson fields. For degenerate flavors I show that a first order phase transition is expected at $\beta = \beta_c$: I speculate on the implications of this structure for Wilson's lattice fermions.

I. Introduction

This talk concerns the mass term $m \bar{\psi}\psi$ in the standard theory of quarks and gluons. It is an abridged version of my recent Phys. Rev. article¹. One of my goals is to provide an intuitive picture for the physical meaning of the CP-violating parameter of the strong interactions. This term, often called the term, is usually discussed in terms of topological excitations of the gauge fields. Here, however, I treat it entirely in terms of the chiral symmetries expected in the massless limit of the theory.

I conclude that a first-order transition is expected at $\beta = \beta_c$ when the flavors have a small degenerate mass. This transition can be removed if flavor-breaking is large enough. At the transition, CP is spontaneously broken. I will also remark on the implications for the structure of Wilson's lattice fermions.

This is a subject with a long history, and most of what I say is buried in numerous previous studies. The implications of θ to the fermion mass matrix are well known to low-energy chiral Lagrangian discussions²⁻⁹. The possibility of a first-order phase transition at large θ has been discussed in³. The possibility of a spontaneous breaking of CP was pointed out even before the significance of the parameter θ was appreciated⁴. The relation of θ to lattice Wilson fermions was elucidated some time ago by Seiler and Stamatescu⁵ and was the subject of some recent work of my own¹¹.

The sign of the fermion mass is sometimes regarded as a convention. This is indeed the case for ordinary quantum electrodynamics in four space-time dimensions, where by Furry's theorem¹² there are no triangle diagrams and corresponding anomalies. However, it is explicitly false for the massive Schwinger model of electrodynamics in two space-time dimensions^{13;11}. Furthermore, as the remaining discussion will argue, hadronic physics would change if the sign of one of the quark masses were flipped.




Figure 1: The phase diagram for the one flavor case. The wavy line represents a first-order phase transition, along which $\bar{\psi}_5$ acquires an expectation value. The end point of this transition line is renormalized away from the origin towards negative m_1 .

To start the discussion, consider a change of variables

$$\psi \rightarrow e^{i\theta_5} \psi \quad (1)$$

Since $1 = (\theta_5)^2$, this modifies the fermion mass term to

$$m \bar{\psi} \psi \rightarrow m_1 \bar{\psi} \psi + i m_2 \bar{\psi}_5 \psi_5 \quad (2)$$

where

$$\begin{aligned} m_1 &= m \cos(\theta_5) \\ m_2 &= m \sin(\theta_5) \end{aligned} \quad (3)$$

The kinetic and gauge terms of the quark-gluon action are formally invariant under this transformation. Thus, were one to start with the more general mass term of Eq. (2), one might expect a physical situation independent of θ_5 . However, because of the chiral anomaly, this is not true. The angle θ_5 represents a non-trivial parameter of the strong interactions. Its non-vanishing would give rise to CP violating processes. As such are not observed in hadronic physics, the numerical value of θ_5 must be very small⁶.

If Eq. (1) just represents a change of variables, how can this affect physics? The reason is entwined with the divergences of quantum field theory and the necessity of regularization. Fujikawa¹⁴ has shown how to incorporate the anomaly into the path integral formulation via the fermionic measure, which becomes non-invariant under the above chiral rotation. Under a Pauli-Villars¹⁵ approach θ_5 represents a relative θ_5 rotation between the mass term




Figure 2: The two flavor phase diagram. First-order lines run up and down the m_2 axis. The second order endpoints of these lines are separated by a flavor breaking mass difference. The chiral limit is pinched between these endpoints.

for the fundamental particle and the mass term for a heavy regulator field. On the lattice with Wilson's fermion prescription¹⁶, the doublers play this role of defining the relative chiral phase^{10;11}.

The phase diagram in the $(m_1; m_2)$ plane is strongly dependent on the number of fermion flavors. With a single species, a first-order phase transition line runs down the negative m_1 axis, starting at a non zero value for m_1 . This is sketched in Fig. (1). For two flavors I argue for two first-order phase transition lines, starting near the origin and running up and down the m_2 axis. For degenerate quarks these transitions meet at the chiral limit of vanishing fermion mass, while a small flavor breaking can separate the endpoints of these first-order lines. This is sketched in Fig. (2). With $N_f > 2$ flavors, the $(m_1; m_2)$ plane has N_f first order phase transition lines all pointing at the origin. The conventionally normalized parameter is N_f times the angle to a point in this plane, and these transition lines are each equivalent to going through π .

Whenever the number of flavors is odd, there is a first-order transition running down the negative m_1 axis. Along this line there is a spontaneous breaking of CP, with a natural order parameter being $\bar{\psi}_5 \psi$. This possibility of a spontaneous breakdown was noted some time ago by Dashen⁴ and has reappeared at various times in the lattice context^{17;18}.

I concentrate my discussion on the two flavor case. Here several simplifications make the physics particularly transparent. I then discuss how the one

avor result arises when the other flavor is taken to a large mass. Finally I conjecture on an analogy with heavy doublers and Wilson lattice fermions.

II. The effective potential

I begin by defining eight fields around which the discussion revolves

$$\begin{aligned} &= \frac{c}{ic} \bar{\psi} \psi \\ &\sim = \frac{c}{ic} \bar{\psi} \psi \\ &= \frac{c}{ic} \bar{\psi} \psi \\ &\sim = \frac{c}{c} \bar{\psi} \psi : \end{aligned} \quad (4)$$

The fermion has two isospin components, for which \sim represents the standard Pauli matrices. The factor c is inserted to give the fields their usual dimensions. Its value is not particularly relevant to the qualitative discussion that follows, but one convention is take $c = F = \sqrt{2} f_\pi$ where F is the pion decay constant and the condensate is in the standard form.

Corresponding to each of these quantities is a physical spectrum. In some cases this is dominated by a known particle. There is the familiar triplet of pions around 140 MeV and the eta at 547 MeV. The others are not quite so clean, with a candidate for the isoscalar being the $f_0(980)$ and for the isovector being the $a_0(980)$. I will use that the lightest particle in the channel appears to be heavier than the ρ .

Now consider an effective potential $V(\sigma; \vec{\pi}; \vec{\eta})$ constructed for these fields. I first consider the theory with vanishing quark masses. In the continuum limit, the strong coupling constant is absorbed via the phenomenon of dimensional transmutation¹⁹, and all dimensionless quantities are in principle determined. In the full theory with the quark masses turned back on, the only parameters are those masses and θ .

For the massless theory many of the chiral symmetries become exact. Because of the anomaly, the transformation of Eq. (1), which mixes the σ and $\vec{\pi}$ fields, is not a good symmetry. However favored axial rotations should be valid. For example, the rotation

$$\vec{\pi} \rightarrow e^{i \vec{\pi} \cdot \vec{\tau} / f_\pi} \vec{\pi} \quad (5)$$

mixes with σ

$$\begin{aligned} \sigma &\rightarrow \sigma \cos(\theta) + \vec{\pi} \cdot \vec{\tau} \sin(\theta) / f_\pi \\ \vec{\pi} &\rightarrow \vec{\pi} \cos(\theta) + (\sigma \vec{\tau} + \vec{\pi} \times \vec{\pi}) \sin(\theta) / f_\pi \end{aligned} \quad (6)$$

This transformation also mixes with σ

$$\begin{aligned} \sigma &\rightarrow \sigma \cos(\theta) + \vec{\pi} \cdot \vec{\tau} \sin(\theta) / f_\pi \\ \vec{\pi} &\rightarrow \vec{\pi} \cos(\theta) + (\sigma \vec{\tau} + \vec{\pi} \times \vec{\pi}) \sin(\theta) / f_\pi \end{aligned} \quad (7)$$




Figure 3: The "sombbrero" potential representing the chiral limit of massless quarks.

For the massless theory, the effective potential is invariant under such rotations. In this two flavor case, the consequences can be compactly expressed by going to a vector notation. I define the four component objects $\Phi = (\sigma; \vec{\pi})$ and $\vec{\pi} = (\pi_1; \pi_2; \pi_3)$. The effective potential is a function only of invariants constructed from these vectors. A complete set of invariants is $\Phi^2; \vec{\pi}^2; \Phi \cdot \vec{\pi}$. This separation into two sets of fields is special to the two flavor case, but makes the behavior of the theory particularly transparent.

I now use the experimental fact that chiral symmetry appears to be spontaneously broken. The minimum of the effective potential should not occur for all fields having vanishing expectation. We also know that parity and flavor appear to be good symmetries of the strong interactions, and thus the expectation value of the fields can be chosen in the σ direction. Temporarily ignoring the fields $\vec{\pi}$, the potential should have the canonical "sombbrero" shape, as stereotyped with the form

$$V = \frac{1}{2} (\sigma^2 - v^2)^2 = \frac{1}{2} (\sigma^2 + \vec{\pi}^2 - v^2)^2 \quad (8)$$

Here v is the magnitude of the σ expectation value for σ , and $\frac{1}{2}$ is a coupling strength related to the mass. The normalization convention mentioned below Eq. (4) would have $v = F = 2$. I sketch the generic structure of the potential in Fig. (3). This gives the standard picture of pions as Goldstone bosons associated with fields oscillating along degenerate minima.

Now consider the influence of the fields $\vec{\pi}$ on this potential. Considering small fields, I expand the potential about vanishing

$$V = \frac{1}{2} (\sigma^2 - v^2)^2 + \frac{1}{2} \vec{\pi}^2 (\sigma^2 - v^2)^2 + \dots \quad (9)$$

Being odd under parity, $\vec{\pi}$ appears quadratically.




Figure 4: The effect of M_1 on the effective potential. The ellipse in this and the following figures represents the minima of the effective potential from Fig. (3). The dot represents where the ether settles.

The terms proportional to $\bar{\psi}\psi$ and $\bar{\psi}\gamma_5\psi$ generate masses for the u and d particles. Since $v^2 = v_u^2 + v_d^2$, the $\bar{\psi}\psi$ term contributes equally to each. Substituting $(v;0)$ gives $(\bar{\psi}\psi) = v^2/2$; thus, the $\bar{\psi}\psi$ term breaks the \sim degeneracy. Here is where the observation that the u is lighter than the d comes into play; I have written a minus sign in Eq. (9), thus making the expected sign of positive.

Now I turn on the fermion masses. I consider small masses, and assume they appear as a general linear perturbation of the effective potential

$$V = V_0 + (M_1 \bar{\psi}\psi + M_2 \bar{\psi}\gamma_5\psi) = c \quad (10)$$

Here the four-component objects $M_{1,2}$ represent the possible mass terms. The normalization constant c appears in Eq. (4). The zeroth component of M_1 gives a conventional mass term proportional to $\bar{\psi}\psi$, contributing equally to both flavors. The mass splitting of the up and down quarks appears naturally in the third component of M_2 , multiplying $\bar{\psi}\gamma_5\psi$. The term m_2 of Eq. (2) lies in the zeroth component of M_2 .

The chiral symmetries of the problem now tell us that physics can only depend on invariants. For these I can take M_1^2 , M_2^2 , and $M_1 \cdot M_2$. That there are three parameters is reassuring; there are the quark masses (m_u, m_d) and the CP violating parameter θ . The mapping between these parameterizations is non-linear, the conventional definitions giving

$$\begin{aligned} M_1^2 &= (m_u^2 + m_d^2) = 4 + m_u m_d \cos(\theta) = 2 \\ M_2^2 &= (m_u^2 + m_d^2) = 4 - m_u m_d \cos(\theta) = 2 \\ M_1 \cdot M_2 &= m_u m_d \sin(\theta) = 2 \end{aligned} \quad (11)$$

If one of the quark masses, say m_u , vanishes, then the θ dependence drops out. While this may be a possible way to remove any unwanted CP violation from the strong interactions, having a single quark mass vanish represents a fine tuning which is not obviously more compelling than simply tuning θ to zero. Also, having $m_u = 0$ appears to be phenomenologically untenable^{8,9}.




Figure 5: The effect of M_2 on the effective potential. The dots represent two places where the ether can settle.

I now turn to a physical picture of what the two mass terms M_1 and M_2 do to the "Mexican hat" structure of the massless potential. For M_1 this is easy; it simply tilts the sombrero. This is sketched in Fig. (4). The symmetry breaking is no longer spontaneous, with the tilt selecting the direction for the field to acquire its expectation value. This picture is well known, giving rise to standard relations such as the square of the pion mass being linearly proportional to the quark mass²⁰.

The effect of M_2 is more subtle. This quantity has no direct coupling to the field; so, I must look to higher order. The M_2 term represents a force pulling on the field, and should give an expectation value proportional to the strength, h_i / M_2 . Once ϕ gains an expectation value, it then effects through the ϕ^2 and ϕ^4 terms of the potential in Eq. (9). The ϕ^2 term is a function only of ϕ^2 , and, at least for small M_2 , should not qualitatively change the structure of the symmetry breaking. On the other hand, the ϕ^4 term will warp the shape of our sombrero. As this term is quadratic in ϕ^2 , this warping is quadratic. With ϕ^2 positive, as suggested above, this favors an expectation value of ϕ lying along the vector M_2 , but the sign of this expectation is undetermined. This effect is sketched in Fig. (5).

To summarize, the effect of M_1 is to tilt our Mexican hat, while the effect of M_2 is to install a quadratic warping. The three parameters of the theory are the amount of tilt, the amount of warping, and, finally, the relative angle between these effects. To better understand the interplay of these various phenomena, I now consider two specific situations in more detail.

III. Case A : $M_1 \parallel M_2$

First consider M_1 and M_2 parallel in the four vector sense. This is the situation when we have the two mass terms of Eq. (2) and no explicit breaking of flavor symmetry. Specifically, I take $M_1 = (m_1; \vec{0})$ and $M_2 = (m_2; \vec{0})$. In this case the warping and the tilting are along the same axis.

Suppose I consider m_2 at some non-vanishing fixed value, and study the state of the ether as m_1 is varied. The m_2 term has warped the sombrero,




Figure 6: Varying m_1 at fixed m_2 . A first-order phase transition is expected at $m_1 = 0$. This corresponds to $\beta = 0$. The dots represent places where the chiral can settle.

but if m_1 is large enough, the potential will have a unique minimum in the direction of this pull. As m_1 is reduced in magnitude, the tilt decreases, and eventually the warping generates a second local minimum in the opposite direction. As m_1 passes through zero, this second minimum becomes the lower of the two, and a phase transition occurs exactly at $m_1 = 0$. The transition is first order since the expectation of ϕ jumps discontinuously. This situation is sketched in Fig. (6). From Eq. (11), the transition occurs at $m_u = m_d$ and $\beta = 0$.

As m_2 decreases, the warping decreases, reducing the barrier between the two minima. This makes the transition softer. A small further perturbation in, say, the ϕ_3 direction, will tilt the sombrero a bit to the side. If the warping is small enough, the field can then roll around the preferred side of the hat, thus opening a gap separating the positive m_2 phase transition line from that at negative m_2 . In this way sufficient flavor breaking can remove the first-order phase transition at $\beta = 0$. If I start at $\beta = 0$ with a mass splitting between the up and down quarks, an isoscalar chiral rotation to non-zero ϕ will generate just such a term.

IV. Case B : $M_1 \neq M_2$

I now turn to a situation where M_1 and M_2 are orthogonal. To be specific, take $M_1 = (m_1; 0)$ and $M_2 = (0; 0; 0; m)$, which physically represents a flavor symmetric mass term $m_1 = (m_u + m_d)/2$ combined with a flavor breaking $m = (m_u - m_d)/2$. Now M_2 warps the sombrero downwards in the ϕ_3 direction. A large m_1 would overcome this warping, still giving an ϕ with only ϕ_3 having an expectation value. However, as m_1 decreases in magnitude with a fixed m , there eventually comes a point where the warping dominates the tilting. At this point we expect a new symmetry breaking to occur, with ϕ_3 acquiring an expectation value. This is sketched in Fig. (7). As ϕ_3 is a CP odd operator, this is a spontaneous breaking of CP.

To make this into a proper two dimensional phase diagram, I add an m_3




Figure 7: Varying m_1 at fixed quark mass splitting. A second order phase transition occurs when the tilting is reduced sufficiently for a spontaneous expectation of ϕ_3 to develop. The dots represent places where the ether can settle.




Figure 8: The $(m_1; m_3)$ phase diagram for unequal mass quarks. The wavy line represents a first-order phase transition ending at the second order dots. The light box on the right shows how the one flavor diagram of Fig. (1) is extracted.

piece to the potential. This effectively twists M_1 away from being exactly perpendicular to M_2 . This term explicitly breaks CP and can be expected to remove the transition, just as an applied field removes the phase transition in the Ising model. We thus have a phase diagram in the $(m_1; m_3)$ plane with a first-order transition connecting two symmetrically separated points on the m_1 axis. This is sketched in Fig. (8).

Physically, the endpoints of this transition line are associated with the points where the respective quark masses vanish. The phase transition occurs when the two flavors have masses of opposite sign. Simultaneously flipping the signs of both quark masses can always be done by a flavored chiral rotation, say about the ϕ_3 axis, and thus is a good symmetry of the theory.

Taking one of the flavors to infinite mass provides a convenient way to understand the one flavor situation. As sketched in Fig. (8), this represents looking only at the vicinity of one endpoint of the transition line. In terms of the light species, this transition represents a spontaneous breaking of CP with

a non-vanishing expectation for $\bar{\psi}_5$. In the lattice context the possibility of such a phase was mentioned briefly by Smith¹⁷, and extensively discussed by Aoki and Gocksch¹⁸.

V I. Implications for Wilson's lattice fermions

The Lagrangian for free Wilson lattice fermions is¹⁶

$$L(K; r; M) = \sum_j \bar{\psi}_j (i \not{D} + r) \psi_{j+e} + \sum_j \bar{\psi}_{j+e} (i \not{D} + r) \psi_j + \sum_j (m_1 \bar{\psi}_j \psi_j + i m_2 \bar{\psi}_5 \psi_5) \quad (12)$$

Here j labels the sites of a four dimensional hyper-cubic lattice, μ runs over the space time directions, and e is the unit vector in the μ th direction. I have scaled out all factors of the lattice spacing. The parameter K is called the hopping parameter, and r controls the strength of the so called "Wilson term," which separates out the famous doublers. I have also added an unconventional m_2 type mass term to connect with my earlier discussion.

Being quadratic with only nearest neighbor couplings, the spectrum is easily found by Fourier transformation. Conventionally, a massless fermion is obtained by taking $m_1 = 8K r$, but there are other places where this original particle is massless while other doublers from the naive theory become massless. At $m_1 = 8K r$ one such species does so, at each of $m_1 = 4K r$ there are four massless doublers, and at $m_1 = 0$ find the remaining 6 of the total 16 species present in the naive theory.

I conjecture that these various species should be thought of as flavors. When the gauge fields are turned on, then the full chiral structure should be a natural generalization of the earlier discussion. Thus near $m_1 = 8K r$ I expect a first-order transition to end, much as is indicated in Fig. (1). This may join with numerous other transitions at the intermediate values of m_1 , all of which then finally merge to give a single first-order transition line ending near $m_1 = 8K r$. The situation near 0 and $4K r$ involves larger numbers of flavors, and properly requires a more general analysis. One possible way the lines could join up is shown in Fig. (9a).

For two flavors of Wilson fermions, if we look near to the singularity at $8K r$ we should obtain a picture similar to Fig. (2). However, further away these lines can curve and eventually end in the structure at the other doubling points. One possible picture is sketched in Fig. (9b).




Figure 9: Possible phase diagrams for lattice gauge theory with Wilson fermions. The dashed lines represent first-order phase transitions and the dots represent points where massless excitations should exist. Parts (a) and (b) are for the one and two flavor cases, respectively.

V III. Sum m ary and conclusions

I have presented a physical picture of the parameter in the context of an effective potential for spin-zero bilinears of quark elds. I have argued for a first-order transition at $=$ when all avors are degenerate, and shown how avor breaking can remove this transition.

A number of years ago Tudron and I²³ conjectured on the interplay of the confinement mechanism with χ , and speculated that confinement might make χ unobservable. Recently Schierholz²⁴ argued that keeping confinement in the continuum limit may drive the theory to $\chi = 0$. The connection with present discussion is unclear, but the symmetries seem to indicate no obvious problem with χ being observable. Furthermore, the fact that the χ is lighter than particle candidates in the χ channel suggests that there indeed must be the term of Eq. (9), and it is this term which is directly responsible for the physical dependence on χ .

References

1. M. Creutz, Phys. Rev. D 52, 2951 (1995).
2. G. 't Hooft, Phys. Rev. Lett. 37, 8 (1976); Phys. Rev. D 14 (1976) 3432.
3. E. Witten, Annals of Phys. 128, 363 (1980).
4. R. Dashen, Phys. Rev. D 3, 1879 (1971).
5. J. Gasser and H. Leutwyler, Ann. Phys. 158, 142 (1984); H. Leutwyler, Ann. Phys. 235, 165 (1994); lectures at workshop "Hadrons 1994," Gramado, RS, Brazil (1994) (HEP-PH-9406283).
6. V. Baluni, Phys. Rev. D 19, 2227 (1979); R. Crewther, P. Di Vecchia, G. Veneziano, and I. Witten, Phys. Lett. 88B, 123 (1979).
7. E. Witten, Nucl. Phys. B 223, 422 (1983).
8. J. Donoghue, B. Holstein, and D. Wyler, Phys. Rev. Lett. 69, 3444 (1992); H. Leutwyler, Nucl. Phys. B 337 108 (1990).
9. J. Bijnens, J. P. Rades, and E. de Rafael, Phys. Lett. B 348 226-238 (1995).
10. E. Seiler and I. Stamatescu, Phys. Rev. D 25 (1982) 2177.
11. M. Creutz, Nucl. Phys. B (Proc. Suppl.) 42, 56 (1995).
12. W. Furry, Phys. Rev. 81, 115 (1937).
13. S. Coleman, Annals of Phys. 101, 239 (1976).
14. K. Fujikawa, Phys. Rev. Lett. 42, 1195 (1979).
15. W. Pauli and F. Villars, Rev. Mod. Phys. 21, 433 (1949).
16. K. Wilson, in New Phenomena in Subnuclear Physics, Edited by A. Zichichi (Plenum Press, NY, 1977), p. 24.
17. J. Smith, Nucl. Phys. B 175 307 (1980).
18. S. Aoki, Nucl. Phys. B 314, 79 (1989); S. Aoki and A. Gocksch, Phys. Rev. D 45, 3845 (1992).
19. S. Coleman and E. Weinberg, Phys. Rev. D 7 1888 (1973).
20. M. Gell-Mann, R. Oakes, and B. Renner, Phys. Rev. 175, 2195 (1968).
21. S. Aoki, S. Boetcher, and A. Gocksch, Phys. Lett. B 331, 157 (1994); K. Bitar and P. Vranas, Phys. Rev. D 50, 3406 (1994); Nucl. Phys. B, Proc. Suppl. 34, 661 (1994).
22. M. Creutz, Phys. Rev. Lett. 46, 1441 (1981); M. Creutz and K. J. Moriyarty, Phys. Rev. D 25, 1724 (1982).
23. M. Creutz and T. Tudron, Phys. Rev. D 16 2978 (1977).
24. G. Schierholz, Nucl. Phys. B, Proc. Suppl. 42, 270 (1995).